



## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Ms. Glenda Haage  
Platte Chemical Co.  
P.O. Box 667  
Greeley, CO 80632-0667

DEC 27 2001

Dear Ms. Haage:

Subject: Amendment - Comply with 2000 Diazinon MOA;  
Delete Beans, Cucumbers, Parsley, Parsnips,  
Peas, Peppers, Potatoes, Summer Squash,  
Winter Squash, Sweet Potatoes, Swiss Chard,  
Turnips, Nuisance Pests in Outside Areas,  
Grassland Insects (Roadsides, Ditch Banks,  
Wasteland, Non-crop Areas, & Barrier  
Strips), Lawn Pest Control, and Ornamentals  
except those Grown Outdoors in Nurseries;  
Update First Aid per PR Notice 2001-1; and  
Update CSF/replacement diazinon source  
Clean Crop Diazinon AG500 Insecticide  
EPA Registration Number 34704-41  
Your submission dated June 12, 2001

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
  - a. Delete directions for use for foliar or pre-plant soil application on the crop Beans (including the separate entry for Lima Beans) and on the crop Peas. Under the diazinon MOA, and as stipulated on the labeling for the replacement diazinon manufacturing-use products from which agricultural end-use products may be formulated, the only supported agricultural use pattern for use on beans or peas is for seed treatment, which is not appropriate for this formulation. Also, be sure to delete Beans and Peas from the minimum gallonage and PHI tables that appear under the heading VEGETABLE CROPS.

- b. Since there is no use on lawns for this product, please delete the front panel limitation "THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS".
- c. In the Environmental Hazards, delete the sentences beginning "Because of the migratory..." and "On lawns, if waterfowl..", since these sentences pertain to the use on lawns which you have deleted. Also, delete the phrase "For terrestrial uses," from the sentence beginning "For terrestrial uses, do not apply...", since all the uses on the label are terrestrial uses.
- d. Revise the heading FIRST AID STATEMENTS to FIRST AID and under this heading add "This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase."
- e. Under the FIRST AID advice (and above the hotline numbers) add a Note to Physician as follows:

NOTE TO PHYSICIAN

This product contains diazinon , an organophosphate insecticide that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should only be administered in conjunction with atropine.

- f. Under the heading DIRECTIONS FOR USE, slightly revise the reformulation prohibition to read "Reformulation or repackaging of this product is prohibited."
- g. Under the heading GENERAL INFORMATION, replace "Do not use on food crops grown in greenhouses." with "Do not use in greenhouses."
- h. Under the PHI table for Fruit and Nut Crops, in the definition of PHI replace the semicolon after "harvest" with a period and then delete "may vary based on use pattern selected. See specific directions for use on grapes."
- i. For the crop Caneberries, in the Timing/Special Directions for Leafhoppers delete "Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur.". In the Timing/Special Directions for Dryberry Mites delete "Do not apply within 7 days of harvest or illegal residues may occur.". Then add a NOTES entry for this crop as follows:

**Notes:** (1) Do not apply more than 2 qts. of product per acre per application and no more than 5 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.

- j. For the crop Cherries, in the Timing/Special Directions for Leafhoppers revise "Make 3-4 applications..." to read "Make 3 applications..."

- k. For the crop Strawberries, revise the Timing/Special Directions for Aphids and Two-spotted spider mites to read (in its entirety) "Apply as pest occur. Repeat application, if necessary, after 7 days.". Revise the Timing/Special Directions for Cyclamen Mites to include just the first 3 current sentences. Revise the Timing/Special Directions for Strawberry Leafrollers to include just the first current sentence. Then, add a NOTE for this crop as follows:

**Note:** (1) Do not apply more than 1 qt. of product per acre per application and no more than 4 qts. of product per acre per season (2) To avoid illegal residues, allow a minimum of 5 days between the last application and harvest.

- 1. For the crop Walnuts, delete the sentence "Do not make more than 3 applications per season." and the sentence beginning "Do not apply within ..." from the Timing/Special Directions and then add the following Notes for this crop:

**Notes:** (1) Do not apply more than 4 qts. of product per acre per application and no more than 9 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 45 days between the last application and harvest and do not apply after husks open.

- m. Under the minimum gallonage table fore VEGETABLE CROPS delete the footnote "\*Do not use on commercially grown potatoes which will be hand-harvested.". In the paragraph under that location, revise "Do not apply to food crops grown in greenhouses." to read "Do not apply in greenhouses.".
  - n. Under the VEGETABLE CROPS PHI table, since beans and peas are no longer on the label, revise the second note to read "To Protect Bees: Foliar application of this product should not be made to corn during the pollen shed period.".
3. Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The Basic Confidential Statement of Formula (CSF) dated June 12, 2001 which identifies a replacement diazinon source Manufacturing Use Product is acceptable and has been added to this product's registration file.

Sincerely,

George T. LaRocca  
 Product Manager 13  
 Insecticide Branch  
 Registration Division (7505C)

# RESTRICTED USE PESTICIDE

Due to Avian and Aquatic Toxicity  
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

**THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.**



# DIAZINON AG500 Insecticide

ACCEPTED  
with COMMENTS  
to EPA Letter 04-01  
DEC 27 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 34704-41

## Organophosphate Insecticide

For control of certain insects on fruits, nuts, vegetables, and field crops.

### ACTIVE INGREDIENT:

Diazinon O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....48%

INERT INGREDIENT:.....52%

TOTAL 100%

This product contains 4 lbs. Diazinon per gallon.

\*Contains xylene range aromatic solvent.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

EPA REG. NO. 34704-41

EPA EST. NO. 279-CA-1

NET CONTENTS \_\_\_\_\_ GAL. ( \_\_\_\_\_ L)

See Below For Additional Precautionary Statements and Directions For Use

IMPORTANT: Read the entire Directions for Use and the Notice before using this product. If terms are not acceptable, return the unopened product containers at once.

IHT

EXP 06Y01

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not contaminate ornamental fish ponds.

### Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on the EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, chemical-resistant gloves, such as: barrier laminate, butyl rubber, nitrile rubber or viton; and shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. On lawns, if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), apply at least ¼ inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at applications rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

## FIRST AID STATEMENTS

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person or convulsing person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Reformulation or repackaging this product is prohibited.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

# DIAZINON AG500 INSECTICIDE

EPA REG. NO. 34704-41

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI).

Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves, such as: barrier laminate, butyl rubber, nitrile rubber or viton; and shoes plus socks.

**NOTE:** The California Department of Agriculture has set a 5-day reentry for Diazinon on grapes, peaches, and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

## CHEMIGATION APPLICATION THROUGH AN IRRIGATION SYSTEM (CHEMIGATION) - CRANBERRIES ONLY

DIAZINON AG500 may be applied alone or in combination with other pesticides registered for application through sprinkler irrigation systems to cranberries. To insure compatibility, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is compatible.

Do not apply this product through any type of irrigation system to other crops. For chemigation, apply this product to cranberries only through sprinkler irrigation systems. Do not apply this product to cranberries through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect the irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments, if the need arises.

### Chemigation System Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventor (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

### Sprinkler Chemigation

1. The system must contain a functional check valve, a vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from

being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

### Operating Instructions

1. Determine the acreage to be treated by the sprinkler system.
2. Measure the appropriate amount of DIAZINON AG500 needed to treat the acreage to be treated. Refer to the specific directions for use for cranberries for the application rate and the amount of water per acre.
3. Add the pre-measured DIAZINON AG500 to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if DIAZINON AG500 is tank mixed with other pesticides registered for sprinkler chemigation application to cranberries.
4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the diazinon to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

## WORKER SAFETY RULES

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling Diazinon AG500, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment Section of this label.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture or rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER:** Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-303-356-4400 or 1-800-424-9300 day or night.

## GENERAL INFORMATION

DIAZINON AG500 is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use on food crops grown in greenhouses. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

## FRUIT AND NUT CROPS

### REI - 24 hours

DIAZINON AG500 may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The rate of DIAZINON AG500 for aerial spray or concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gallons of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the Rate calls for 1 pt. of DIAZINON AG500 in 100 gallons of water, this amounts to 3-4 pts. of product per acre when applied at the usual dilute ground spray volumes of 300-400 gallons of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the Timing/Special Directions for individual crops.

# DIAZINON AG500 INSECTICIDE

## EPA REG. NO. 34704-41

CROP	Aerial Application		Ground Application	
	MINIMUM Gals./Acre	DILUTE	MINIMUM Gals./Acre	CONCENTRATE
Almonds*	20	100	20	20
Apricots	20	100	20	20
Blackberries**	20	100	20	20
Boysenberries**	20	100	20	20
Cherries	20	100	20	20
Cranberries	20	15	-	-
Dewberries**	20	100	20	20
Grapes	20	100	20	20
Loganberries**	20	100	20	20
Nectarines	20	100	20	20
Peaches	20	100	20	20
Plums	20	100	20	20
Prunes	20	100	20	20
Raspberries**	20	100	20	20
Strawberries	20	100	20	20
Walnuts*	20	100	20	20

\*CA only. \*\*CA, OR, or WA only.

Observe stated time intervals between last application and harvest, as well as intervals between applications and total amount of product that can be applied per season.

Crop	*PHI
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Cranberries	7
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums, Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

\*PHI = Pre-Harvest Interval or the number of days between the last application and harvest; may vary based on use pattern selected. See specific directions for use on grapes.

\*\* = Dormant spray only.

Note: To protect bees, do not apply this product to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Almonds (CA only)	San Jose Scale, Parlatoria Scale	1-1½ pts. + 2- 3 gallons dormant oil or 1-1½ gallons superior type oil in 100 gallons of water	Apply as a dormant spray. Do not apply more than 6 pts. of diazinon or 9 gallons of oil per acre.
Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Crawlers	1 pt. in 100 gallons of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
	Olive Scale Crawlers	½ pt. + 1½ gallons of light medium horticultural oil per 100 gallons of water	Apply when scale crawlers are present.
	Apricot Mealybugs	1 pt. per 100 gallons of water	Apply as a cover spray from petal fall to June. Allow 14 days between applications.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Caneberries (CA, OR, and WA only) (Blackberries, Boysenberries, Dewberries, Logan berries, Raspberries)	Leafhoppers, Thrips, Aphids, Two-spotted Spider Mites	1 qt. per acre	Apply as insects occur. Repeat application after 14 days, if necessary. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur.
	Raspberry Fruitworms, Raspberry Sawflies		Apply in a minimum of 100 gallons of water per acre. (200 gallons per acre maximum). For Raspberry Fruitworms make one application when blossom buds separate. Make a second application immediately before blossoms open.
Cherries	Dryberry Mites	1-2 qts. per acre	Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed but still closed. Apply in a minimum of 100 gallons of water per acre. Do not apply within 7 days of harvest or illegal residues may occur.
	Raspberry Crown Borers, (Raspberry Root Borers)	2 qts. per acre	Apply a single application in a minimum of 100 gallons of water per acre as a trench to the crown and lower canes in the Spring before buds break.
Cherries	Eyespotted Bud Moths, Fruit Tree Leafrollers	1 pt. in 100 gallons of water	Apply in cover sprays as infestations occur beginning at petal fall. Repeat application after 7 days, if necessary.
	Leafhoppers	1 pt. in 100 gallons of water	Make 3-4 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications.
	Cherry Fruitflies	½-1 pt. in 100 gallons of water	Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10 day intervals.
	Black Cherry Aphids	1 pt. in 100 gallons of water	Apply as a cover spray when aphids appear. Repeat application after 10 days, if necessary.
Cranberries	San Jose Scale Crawlers	1 pt. in 100 gallons of water	Apply during the pre-pink stage or when crawlers first appear.
	Cherry Rust Mites	½-1 pt. in 100 gallons of water	Apply to trees as a single supplemental spray after harvest.
Cranberries	Blackheaded Fireworms ( <i>Rhopobota naevana</i> )	2 qts. in a minimum of 15 gallons of water per acre for ground application and up to 400 gallons of water per acre for chemigation.	Apply when pest appears. Repeat application after 14 days, if necessary. A maximum of 6 applications per season is permitted.

Notes: (1) Do not make more than 3 applications per season. (2) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (3) Allow a minimum of 21 days between the last application and harvest.

## DIAZINON AG500 INSECTICIDE EPA REG. NO. 34704-41

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Cranberries Cont'd.	Cranberry Fruitworms	3 qts. in a minimum of 15 gallons of water per acre for ground application and up to 400 gallons of water per acre for chemigation.	Apply as pest appears. Repeat application after 14 days, if necessary. A maximum of 4 applications per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur.

Notes: (1) Do not apply more than 12 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest. (3) This product may be applied to cranberries through sprinkler irrigation. Refer to the section CHEMIGATION for specific chemigation directions for use. (4) Do not use water from irrigated or flooded cranberry beds for drinking purposes or to irrigate crops other than those appearing on EPA approved diazinon labels.

Grapes	Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leafhoppers, Omnivorous Leafrollers, Aphids, Spider Mites	1-2 pts. per acre in a minimum of 100 gallons of water.	Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary.
	<i>Drosophila</i> spp.	6 fl. oz. per acre in a minimum of 100 gallons of water (300 gallons of water maximum)	Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary.

Notes: (1) Do not apply more than 2 pts. of product per acre per application and no more than 10 pts. per acre per season. (2) To avoid illegal residues, allow 28 days between the last application and harvest.

Nectarines	Oriental Fruitmoths	1 pt. in 100 gallons of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight.
	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, Peach Twig Borers	1 pt. in 100 gallons of water	Apply as infestations occur. Repeat application after 14 days, if necessary.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Peaches	Aphids, Brown Mites (Clover Mites), Two spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 pt. in 100 gallons of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
	Oriental Fruitmoths	1 pt. in 100 gallons of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight.
	Peach Twig Borers	1 pt. in 100 gallons of water	Apply as insects occur in May and June. Allow 7 days between applications.
	Leafhoppers	1 pt. in 100 gallons of water	Make 3-4 applications beginning at petal fall and continuing through the growing season. Allow at least 7 days between applications.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Peaches Cont'd.	White Peach Scale	1 1/2-2 pts. in 100 gallons of water	Apply post-harvest timed to coincide with peak crawler and immature scale activity. Under conditions of heavy infestation 2 applications (7 days apart) may be needed. Do not apply before fruit is harvested.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Plums	Brown Mites (Clover Mites), European Red Mites	1 pt. in 100 gallons of water	Apply as needed every 7 days
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	1/2-1 pt. in 100 gallons of water	

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Prunes	Brown Mites (Clover Mites), European Red Mites	1 pt. in 100 gallons of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	1/2-1 pt. in 100 gallons of water	

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Strawberries	Aphids, Two spotted Spider Mites	1 pt. in 100 gallons of water per acre	Apply as pests occur. Repeat application if necessary every 7 days, but do not make more than 3 applications per season. Do not apply within 5 days of harvest or illegal residues may occur.
	Cyclamen Mites	1 qt. in 100 gallons of water per acre	Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat applications after 7 days if necessary. Do not make more than 3 foliar applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Application may be made to plants after harvest.
Walnuts (CA only)	Mole Crickets	1 qt. per acre	Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1-2 days before transplanting and immediately incorporate into the top 1-2 inches of soil.
	Strawberry Leafrollers	3/4-1 pt. in 100 gallons of water	Apply 200 gallons of spray per acre when blossoms show color. Do not apply within 5 days of harvest or illegal residues may occur.
Walnuts (CA only)	Aphids, Spide Mites	1 qt. per acre	Apply as a foliar spray Repeat application in 14 days if necessary. Do not make more than 3 applications per season. Do not apply within 45 days of harvest or after husks open or illegal residues may occur. Make ground applications in a minimum of 100 gallons of water per acre.
	Codling Moths, Scale Crawlers, Walnut Caterpillars	3 qts. per acre	

**DIAZINON AG500 INSECTICIDE**  
**EPA REG. NO. 34704-41**

**VEGETABLE CROPS**

REI - 24 hours

For foliar or soil applications, DIAZINON AG500 should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. DIAZINON AG500 may be applied using ground or aerial application equipment as specified in the following table.

The table indicates the minimum amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application		Aerial Application MINIMUM Gallons/Acre
	MINIMUM Gallons/Acre	MINIMUM Gallons/1,000 Sq Ft.	
Beans	10	2	5
(Lima, Pole, and Snap) (succulent only)			
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabas	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5
Chinese Radish (CA, FL. only)	10	2	5
Collards	10	2	5
Crenshaws	5	1	5
Endive (Escarole)	10	2	5
Ginseng	10	2	5
Honeydew Melons	5	1	5
Kale	10	2	5
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	5
Mustard	10	2	5
Onions (Bulb and Green)	10	2	5
Peas (succulent only)	10	2	5
Persian Melons	5	1	5
Radishes	10	2	5
Spinach	10	2	5
Sweet Corn	20	4	5
Tomatoes	10	2	5
Watermelons	5	1	5

\*Do not use on commercially grown potatoes which will be hand-harvested.

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

**Pre-Harvest Intervals (PHI\*) for Foliar Applications**

Crop	PHI* (Days)
Beans (Lima, Pole, Snap)	7
Beets, Red (Table)	14
Broccoli	7
Brussels Sprouts	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli	10
Chinese Cabbage	10
Chinese Mustard	10
Chinese Radish	10
Collards	10
Crenshaws	3
Endive (Escarole)	14
Ginseng	30
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Mustard	10
Onions (Bulb and Green)	14
Peas	7
Persian Melons	3
Radishes	14
Spinach	14

**Pre-Harvest Intervals (PHI\*) for Foliar Applications, Cont'd.**

Crop	PHI* Days
Sweet Corn	7
Tomatoes	1
Watermelons	3

\*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

Notes: 1) Soil incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see TIMING/SPECIAL DIRECTIONS) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. 2) To Protect Bees: Foliar applications of this product should not be made to peas or beans if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Beans (Lima, Pole or Snap) (succulent only)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworm	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers	1-1½ pts. per acre	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season.  Do not apply within 7 days of harvest or illegal residues may occur.
Beans, Lima (succulent only)	Cucumber Beetles	¾ pt. per acre	Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. Make cucumber beetle application in 100 gallons of water per acre.  Notes: 1) Diazinon will not control organophosphate-resistant leafminers. 2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
	Leafhoppers, Spider Mites	1 pt. per acre	Apply as pests occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season.  Do not apply within 7 days of harvest or illegal residues may occur.  Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.  Note: To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
Beets, Red (Table)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.



## DIAZINON AG500 INSECTICIDE EPA REG. NO. 34704-41

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Beets <i>Red (Table)</i> <i>cont'd.:</i>	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. <b>Note:</b> Diazinon will not control organophosphate-resistant leafminers.
Broccoli	Root Maggots	2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. per 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. <b>Note:</b> Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.  Do not apply within 7 days of harvest or illegal residues may occur. <b>Note:</b> Diazinon will not control organophosphate-resistant leafminers.
	Brussels Sprouts	Root Maggots	2-3 qts. per acre
4-8 oz. in 50 gallons of water			In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. <b>Note:</b> Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Brussels Sprouts <i>cont'd.:</i>	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbageworms	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur.
Cabbage	Root Maggots	2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. in 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. <b>Note:</b> Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage worms, Leafminers, Dipterous	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.  Do not apply within 21 days of harvest, or illegal residues may occur. <b>Note:</b> Diazinon will not control organophosphate-resistant leafminers.
	Carrots	Cutworms	2-4 qts. per acre
Mole Crickets		1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
Wireworms		3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

**DIAZINON AG500 INSECTICIDE**  
**EPA REG. NO. 34704-41**

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Carrots cont'd.:	Aphids	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.
Cauliflower	Root Maggots	2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. in 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Broccoli	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Chinese Mustard (Gai Choy)	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Radish (Japanese Radish or Daikon) (FL and CA only)	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Collards	Cutworms	2-4 qts. per acre.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage worms, Harlequin bugs, Corn Earworms	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
Endive (Escarole)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.

**DIAZINON AG500 INSECTICIDE**  
**EPA REG. NO. 34704-41**

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Ginseng	Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice	¾-1 pt. per acre	Spray when insects first appear. For heavy insect infestations, use 1 pt. per acre. Apply by ground equipment using 10 to 100 gallons of water per acre. Do not apply more than one application per growing season. Do not apply during flowering of three or four year old crops. Do not apply within 30 days of harvest. Do not graze treated areas or feed treated forage to live stock.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Kale	Aphids, Diamondback Moths, Imported Cabbage worms, Harlequin bugs, Corn Earworms	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Lettuce	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians and Hybrids of these) and Watermelons	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	½-1½ pts. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Mustard	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Onions (Bulb and Green)	Onion Maggots	2-4 qts. per acre	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Diazinon will not control organophosphate-resistant onion maggots.
	Onion Thrips	1 pt. per acre	Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into top 4-8 inches of soil.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
Peas (succulent only)	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	¾-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. To avoid illegal residues, allow a minimum of 7 days between the last application and harvest. Vines and hay may be fed to dairy and beef cattle and sheep. To avoid illegal residues, wait 7 days following treatment if forage is to be cut for hay. Notes: 1) Diazinon will not control organophosphate-resistant leafminers. 2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
Radishes	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.

**DIAZINON AG500 INSECTICIDE**  
EPA REG. NO. 34704-41

**DIAZINON AG500 INSECTICIDE**  
EPA REG. NO. 34704-41

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Radishes cont'd.:	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipteropus Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season.  Do not apply within 14 days of harvest or illegal residues may occur.  Note: Diazinon will not control organophosphate-resistant leafminers.
Spinach	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½-1 pt. per acre	Apply as Insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.  Do not apply within 14 days of harvest.  Note: Diazinon will not control organophosphate-resistant leafminers.
Sweet Corn	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Seed Corn Maggots	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Corn Earworms	1-1½ qts. per acre	Apply when silks first appear. Repeat application every 7 days if necessary. Do not make more than 5 applications per season.
	Corn Rootworm (Adults).	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Corn Leaf Aphids	1-2 pts. per acre	Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Grasshoppers, Spider Mites, Fleabeetles	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Sweet Corn cont'd.:	Sap Beetles	2-2½ pts. per acre	Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Tomatoes	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½ pt. per acre	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within one day of harvest or illegal residues may occur
	Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms, Beet Armyworms	¾-1 pt. per acre	Notes: Diazinon will not control organophosphate-resistant leafminers.
	Vinegar Flies ( <i>Drosophila spp.</i> )	½-1½ pts. per acre	

Notes: (1) To protect bees, do not apply foliarly to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be picked 7 days after the last application.

**FIELD CROPS**

**REI - 24 hours**

Soil and foliar applications of DIAZINON AG500 should always be applied as a spray in sufficient water to ensure thorough coverage. For foliar applications, observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

**Pre-harvest Interval (PHI)\* for Foliar Applications**

Crop	PHI* (Days)
Hops	14
Sugar Beets	14

\*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Hops	Aphids, Spider Mites	1 qt. per acre	Apply as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 4 applications per season.  Do not apply within 14 days of harvest.
Sugar Beets	Wireworms	3-4 qts. per acre	Broadcast just before planting. Incorporate into soil 4-6 inches. Do not apply more than 4 qts. of this product to the soil, per season.

**DIAZINON AG500 INSECTICIDE**  
**EPA REG. NO. 34704-41**

REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Sugar Beets cont'd.:	Beet Leafhoppers, Dipterous Leafminers, Aphids	¾-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.  Do not apply within 14 days of harvest.  Note: Diazinon will not control organophosphate- resistant leafminers.
	Grasshoppers	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.  Do not apply within 14 days of harvest.

Note: Sugar beet tops may be fed to beef and dairy animals.

**INSECT CONTROL ON ORNAMENTALS**  
**GROWN OUTDOORS IN NURSERIES**

REI - 12 hours

To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladioli, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Spruce, and Willow, apply the recommended rates indicated below. Try to spray underside of leaves and penetrate dense foliage.

Insects	Amount of DIAZINON AG500 to Use	
	Rate per 3 gallons of water	Rate per 100 gallons of water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omniverous Leaf-tiers, Privet Mites, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mites, Juniper Webworms, Whiteflies	½ fl. oz.*	1 pt.**
Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, Fall Webworms, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars.	1½ fl. oz.*	3 pts.**

\*1 fl. oz. = 2 tablespoons

\*\*16 fl. oz. = 1 pint

**Precaution: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.**

**Mimosa Webworms:** Mix ½ oz. in 3 gallons of water (or 1 pt. in 100 gallons of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

**NOTICE**

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION. ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR

FORMULATED FOR  
**PLATTE CHEMICAL CO.**  
 419 18TH STREET, GREELEY, COLORADO 80631-5852