

ACCEPTED

JUN 4 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act
 as amended by the FIFRA Act of 1972
 34704-55

CLEAN CROP
CHLORPYRIFOS 1/2 G
TURF INSECTICIDE

ACTIVE INGREDIENT:

Chlorpyrifos 10,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)phosphorothioate)..... 0.5%

INERT INGREDIENTS: 99.5%

TOTAL 100.0%

EPA Reg. No. 34704-55

EPA Est. No.

Do Not Ship or Store with Food, Feeds, or Clothing

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only as prescribed on this label.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry. **DO NOT** apply to dachshund's jaws.

STORAGE AND DISPOSAL (12 pt.)

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration or leakage.

STORAGE: Store in safe areas. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Completely empty bag into application equipment. Then dispose of empty

BEST AVAILABLE COPY

bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Apply this product uniformly to home lawns and other ornamental and recreational turf grass areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using this product at the rate of 200 pounds per acre. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. Immediately after application sprinkle the turf with 100 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine amounts of this product and water needed for smaller sized areas. Retract as necessary.

Controls Ants, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Hyperodes (Turfgrass) Weevils*, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs), and Certain White Grubs** Infesting Home Lawns and other Ornamental Turf Areas.

*To control Hyperodes (Turfgrass) weevils use twice the rate of CHLORPYRIFOS 1:2:6 shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

**To control certain white grubs (larvae of European chafers and Japanese beetles) use two to four times the amounts of CHLORPYRIFOS 1:2:6 shown in the table i.e., 10 to 20 pounds per 1,000 sq. ft. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat other grubs (e.g. young and actively feeding near the soil surface) usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

RATES TO USE

Area to be Treated	Pounds of CHLORPYRIFOS 1:2:6	Gallons of Water to Sprinkle on Turf After Treatment
1,000 sq. ft.	5	30
5,000 sq. ft.	25	150
10,000 sq. ft.	50	300

USE INSTRUCTIONS

Seller and Manufacturer warrant that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the manufacturer and seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. Seller and Manufacturer make no other express or implied warranty of fitness or

merchantability. The exclusive remedy of the buyer or user, and the limit of the liability of manufacturer and seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice which may not be varied by any verbal or written agreement.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. 14TH STREET FREMONT, NEBRASKA 68025

BEST AVAILABLE COPY