

11-27-71

ALMONDS: Brown Rot, Shot Hole (Coryneum Blight) - Use 2 pounds per 100 gallons of water in pre-bloom, blossom and petal fall periods. An additional application should be made 5 weeks after petal fall using 2 pounds per 100 gallons of water. Scab - Use 2 pounds in 100 gallons of water. Apply at petal fall and at 5 weeks after petal fall. Do not use later than 5 weeks after petal fall.

APPLES: Scab - Use 1 to 1½ pounds in 100 gallons of water in pre-blossom, blossom and early cover sprays. Do not apply within 7 days of harvest. Apples may be injured by zinc compounds under certain conditions. Butte Rot (Oregon and Washington) - Use 1½ to 2 pounds in 100 gallons of water beginning 12 to 14 days after the calyx stage and repeating as necessary in summer sprays. Perennial Canker (Oregon) - Use 1 to 1½ pounds in 100 gallons of water in pre-blossom and calyx sprays.

APRICOTS: Shot Hole Fungus - Use 2 pounds per 100 gallons of water. Apply in pre-bloom, blossom and petal fall through early cover sprays.

BEANS (SNAP, LIMA): Anthracnose - Use 2 pounds per acre in sufficient water for thorough coverage when plants are small. Increase to 3 pounds per acre on large plants. Make applications at 7 day intervals starting when true leaves appear. Do not apply to snap beans later than 4 days before picking.

BLACKBERRIES: Evergreen Blackberry Canker (Pacific Northwest) - 1½ to 2 pounds between mid-June and early July. Do not make more than 1 application.

BLUEBERRIES: Mummy Berry - 3½ pounds at loose bud scale and 7 days later. No applications later than 3 weeks after full bloom. Use 200 gallons of spray per acre.

CELERY: Early Blight (Cercospora), Late Blight (Sept. a) and Damping-Off - 1½ to 2 pounds or 1 pound of Clean Crop 21RAM plus 1 pound 95% Ferbam Wettable Powder at weekly intervals beginning in the plant bed or at 3 to 5 day intervals during severe conditions. Remove residues by trimming and washing.

CHERRIES: Bitter Rot/Blossom Blight - Use 1½ pounds per 100 gallons of water. Apply in popcorn, full-blossom and petal fall stages of growth. Leaf Spot - Use 1½ pounds per 100 gallons of water. Apply in petal fall, shuck fall stage and approach, 2 weeks after shuck fall. Do not apply later than 10 weeks after shuck fall.

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Downy Mildew, Alternaria Leaf Blight, Anthracnose, and Angular Leaf Spot - 1½ to 2 pounds at 7 to 10 day intervals under conditions favorable to disease, or at 3 to 5 day intervals when disease conditions are great. Increase rate to 3 pounds per 100 gallons of water per acre after plants vine.

NECTARINES: Peach Leaf Curl and Shot Hole - 1½ to 3½ pounds at dormant and popcorn stages. Fruit Brown Rot - 1½ to 2 pounds at petal fall, 2 weeks later, and additional applications at 1 to 2 week intervals beginning near the beginning of last month before picking. Remove residues by wiping or brushing if applied during the last two weeks before picking.

ORNAMENTALS: Azaleas and Camellias (Petal Blight) - Begin at first bloom. Repeat at 7 to 10 day intervals through bloom. Spray flowers and drench mulch and litter beneath bushes. 1½ to 2 pounds at 3 to 5 day intervals as required. Chrysanthemums and Snapdragons (Leaf Spot) - 1½ to 2 pounds every 7 to 10 days or as required. Begin before disease is expected to appear or at first sign of disease.

PEACHES: Peach Blight (Coryneum) and Leaf Curl - Use 2 to 3 pounds in 100 gallons of water in the fall before heavy rains. For Leaf Curl alone, use 2 to 3 pounds in 100 gallons of water as a dormant application prior to bud swell. Brown Rot and Peach Blight (Coryneum) - Use 2 pounds in 100 gallons of water in pre-bloom, blossom and petal fall sprays. May be used in early cover and before harvest. Remove excess residue by wiping or washing.

PEARS: Scab - Use 1 to 1½ pounds in 100 gallons of water in pre-blossom, blossom and early cover sprays. Do not apply within 7 days of harvest. Storage Rot (Oregon) - Use 1 to 1½ pounds in 100 gallons of water as pre-harvest sprays.

PECANS: Scab - 2 pounds four times at 21 to 28 day intervals when tips of small nuts become brown. A fifth application in August may be required when danger of disease is great.

PINE SEEDLINGS: Fusiform Rust in Plant Beds - 2 pounds at 3 to 5 day intervals from emergence to mid-June.

STRAWBERRIES: Leaf Spot - 1½ to 2 pounds when blossom bud shows and repeat when first blossoms open. Gray Mold (Botrytis) - 1½ to 2 pounds at 10 day intervals, starting at early blossom stage and making 3 to 5 sprayings. Remove residues by washing if sprayed as late as 7 days before picking.

TOMATOES: Anthracnose, Early Blight - Use 2 to 3 pounds per 100 gallons of water per acre at 7 to 10 day intervals starting 2 to 4 weeks after plants are set. Under severe disease conditions apply 3 pounds in 180 to 200 gallons per acre at 7 day intervals. Late Blight - Alternate 1½ to 2 pounds with maneb at weekly intervals. Begin application when first fruit forms.

GREENHOUSE TOMATOES: Gray Mold - 1½ to 2 pounds weekly or twice weekly under severe conditions.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025

AC-1000
WETTABLE POWDER
in EPA Approved Formulation

1990

Under the Federal Insecticide, Fungicide and Rodenticide Act, it is a violation of law to apply this product in a manner inconsistent with the instructions and claims contained in the label.

34704-67

BEST AVAILABLE COPY