

JUL 26 1996

Mr. J. Allen Dunlap III
Platte Chemical Co.
150 So. Main St.,
Fremont, NE 68025-5697

Dear Mr. Dunlap:

Subject: Ziram 76WP
EPA Registration No. 34704-67
Application Dated January 10, 1996, Request To
Delete Use Sites from This Registration Due
To Non-Support for Reregistration Under Section
4 of FIFRA

The proposed deletions of use sites: Beans (Snap and Lima), Celery, Cucumbers, Melons, Pumpkins, Squash, Tomatoes, including Greenhouse Tomatoes have been reviewed and your request was published in the Federal Register, Vol. 61: 16780, April 17, 1996. There were no objections to the request; therefore, the proposed deletions of use-sites as named above are acceptable under the Federal Insecticide, Fungicide and Rodenticide Act as amended. A stamped copy of the label that does not show the deleted use-patterns is enclosed for your records.

In future amendments to the subject labeling, you must inform the using public as to when to use the high and low dosages of the range claimed for each use-site of this pesticide product.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

E. Wilson: ABC30:07-25-96

BEST AVAILABLE COPY

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							



ZIRAM 76 WP

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUL 26 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act

as amended, for the pesticide
registered under EPA Reg. No.
34704-67

ACTIVE INGREDIENT:	BY WEIGHT
*Ziram	76.00%
INERT INGREDIENTS:	24.00%
TOTAL	100.00%
*Zinc Dimethyl dithiocarbamate	
Zinc expressed as metallic. . .	16.25%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements.

NET WEIGHT _____ LBS. (_____ KG)

EPA REG. NO. 34704-67

EPA EST. NO. _____

5162

EXP12G95

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

May be harmful if swallowed or inhaled! May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist.

Personal Protective Equipment

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, flush with plenty of water; for eyes, get medical attention.

ENVIRONMENTAL HAZARDS

Observe use limitations given for specific crops.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or

disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label. Do not apply to water used for irrigation, recreational and domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS

TO MIX—FILL the spray tank about 2/3 full of water, and with the agitator running, slowly sift in the required amount of this material. Finish filling the tank with water and let mixing continue in the tank until thoroughly mixed before applying and then with the agitator running apply at once. Do not apply this product through any type of irrigation system.

ALMONDS: Brown Rot, Shot Hole (Coryneum Blight) - Use 2 pounds per 100 gallons of water in prebloom, blossom and petal fall periods. An additional application should be made 5 weeks after petal fall using 2 pounds per 100 gallons of water. Scab - Use 2 pounds in 100 gallons of water. Apply at petal fall and at 5 weeks after petal fall. Do not use later than 5 weeks after petal fall.

APPLES: Scab - Use 1 to 1 1/2 pounds in 100 gallons of water in pre-

393

ZIRAM 76 WP
EPA REG. NO. 34704-67

blossom, blossom and early cover sprays. Do not apply within 7 days of harvest. Apples may be injured by zinc compounds under certain conditions. Bulleye Rot (Oregon and Washington) - Use 1 1/2 to 2 pounds in 100 gallons of water beginning 12 to 14 days after the calyx stage and repeating as necessary in summer sprays. Perennial Canker (Oregon) - Use 1 to 1 1/2 pounds in 100 gallons of water in pre-blossom and calyx sprays.

APRICOTS: Shot Hole Fungus - Use 2 pounds per 100 gallons of water. Apply in pre-bloom, blossom and petal fall through early cover sprays.

BLACKBERRIES: Evergreen Blackberry Canker (Pacific Northwest) - 1 1/2 to 2 pounds between mid-June and early July. Do not make more than 1 application.

BLUEBERRIES: Mummy Berry - 3 1/2 pounds at loose bud scale and 7 days later. No applications later than 3 weeks after full-bloom. Use 200 gallons of spray per acre.

CHERRIES: Brown Rot Blossom Blight - Use 1 1/2 pounds per 100 gallons of water. Apply in popcorn, full blossom and petal fall stages of growth. Leaf Spot - Use 1 1/2 pounds per 100 gallons of water. Apply in petal fall, shuck fall stage and approximately two weeks after shuck fall. Do not apply later than two weeks after shuck fall.

NECTARINES: Peach Leaf Curl and Shot Hole - 1 1/2 to 3 1/4 pounds at dormant and popcorn stages. Fruit Brown Rot- 1 1/2 to 2 pounds at petal fall, 2 weeks later, and additional applications at 1 to 2 week intervals beginning near the beginning of last month before picking. Remove residues by wiping or brushing if applied during the last two weeks before picking.

ORNAMENTALS: Azaleas and Camellias (Petal Blight) - Begin at first bloom. Repeat at 7 to 10 day intervals through bloom. Spray flowers and drench mulch and litter beneath bushes - 1 1/2 to 2 pounds at 3 to 5 day intervals as required. Chrysanthemums and Snapdragons (Leaf Spot) - 1 1/2 to 2 pounds every 7 to 10 days or as required. Begin before disease is expected to appear or at first sign of disease.

PEACHES: Peach Blight (Coryneum) and Leaf Curl - Use 2 to 3 pounds in 100 gallons of water in the fall before heavy rains. For Leaf Curl alone, use 2 to 3 pounds in 100 gallons of water as a dormant application prior to bud swell. Brown Rot and Peach Blight (Coryneum) - Use 2 pounds in 100 gallons of water in pre-bloom, blossom and petal fall sprays. May be used in early cover and before harvest. Remove excess residue by wiping or washing.

PEARS: Scab - Use 1 to 1 1/2 pounds in 100 gallons of water in pre-blossom, blossom and early cover sprays. Do not apply within 7 days of harvest. Storage Rot (Oregon) - Use 1 to 1 1/2 pounds in 100 gallons of water as pre-harvest sprays.

PECANS: Scab - 2 pounds four times at 21 to 28 day intervals when tips of small nuts become brown. A fifth application in August may be required when danger of disease is great.

PINE SEEDLINGS: Fusiform Rust in Plant Beds - 2 pounds at 3 to 5 day intervals from emergence to mid-June.

STRAWBERRIES: Leaf Spot - 1 1/2 to 2 pounds when blossom bud shows and repeat when first blossoms open. Gray Mold (Botrytis) - 1 1/2 to 2 pounds at 10 day intervals, starting at early blossom stage and making 3 to 5 sprayings. Remove residues by washing if sprayed as late as 7 days before picking.

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697