

Precautionary Statements
HAZARDS TO HUMANS &
DOMESTIC ANIMALS

CAUTION: Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water, and get medical attention if irritation persists. May cause irritation of nose, throat or skin. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from food, feedstuffs and domestic water supplies.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by federal, state or local disposal authorities.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of in an approved landfill or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures.

APPLICATION INSTRUCTIONS

Unless otherwise specified for specific tree fruits, dosage rates are given as pints of SULFUR 6 FLOWABLE per 100 gallons water for use in a thorough coverage spray.

APPLE, PEAR - Scab, Pre-Bloom through Calyx sprays--1 1/2 to 3 1/2 pints. Cover sprays--1 to 2 pints. DO NOT apply to D'Anjou or Comice Pears. Powdery Mildew: Pre-Bloom through calyx sprays--2 1/2 to 3 pints. Cover sprays--2 pints. **Pear Rust Mite:** Use 1 quart per 100 gallons to be applied with oil as a delayed dormant or dormant application.

CHERRY, PLUM, PRUNE - Brown Rot, Leafspot: Pink and bloom sprays--1 1/2 to 3 1/2 pints. Petal fall, shuck and cover sprays--1/2 to 1 1/2 pints. Not recommended for application on Deacon and Van varieties of cherries.

RUS (Grapefruit, lemons, limes, oranges, tangelos and tangerines) - Rust Mite--Apply 1 to 4 pints. Do not apply within 4 weeks of an oil application unless applied with a dormant oil to winter-dormant woody plants. Thrips (Early spring treatment for the reduction of Thrips population)--Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

CORN, SORGHUM and SMALL GRAINS - Spider and Common Leaf Spot Mites--Apply 1 gallon per acre in sufficient water to insure thorough coverage when mites appear on bottom 2 leaves. With aerial and concentrate sprayers, use 6 to 20 gallons of water per acre. For other ground sprayers, use a minimum of 20 gallons per acre. Repeat applications may be necessary if conditions persist. Can be used in tank mix combinations with other miticides.

COTTON - Atlantic Mite--2 pints for early season control and up to 1 gallon for mid to late season control.

GRAPES - Powdery Mildew--1 to 2 pints per 100 gallons water, maximum of 8 pints SULFUR 6 FLOWABLE per acre in spring and summer. Make first application when shoots are 6 to 8 inches long. Make second application when shoots are 12 to 15 inches long, and third at about bloom stage.

The timing, rate, and number of applications will vary with local conditions. Consult State Agricultural Service or State Agricultural Experiment Stations in your region for specific information. Follow the **USE PRECAUTIONS** given on this label.

PEACH, NECTARINES - Powdery Mildew, Brown Rot, Leafspot: Pink and bloom sprays--1 1/2 to 3 1/2 pints. Petal fall, shuck and cover sprays--1/2 to 1 1/2 pints. Application to mature nectarines may cause discoloration.

PEANUTS - Rust--2 to 4 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

POTATOES - Powdery Mildew--3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3-week intervals.

STRAWBERRIES - Powdery Mildew--1-2 pints. Apply at first sign of infection and repeat at 3-week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canner's recommendations.

SUGAR BEETS - Powdery Mildew--Apply the recommended rate in 5 to 10 gallons of water per acre by air at first sign of infection. Use 4 pints per acre in the Pacific Northwest, 8 pints per acre in Arizona and 1 to 4 quarts in Colorado, Kansas, Nebraska and Wyoming. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season.

TOMATOES - Tomato Russet Mites--1/2 to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

FOR SPRINKLER IRRIGATION SYSTEM APPLICATION (POTATOES AND SUGAR BEETS) INJECT IN SYSTEM AT OR NEAR IRRIGATION PUMP. Center Pivot System--Apply during last full revolution of the system. Portable or Solid Set Systems--Apply during final 15 minutes of irrigation period. Do not operate irrigation system after application until deposit has thoroughly dried and several days have lapsed. All irrigation systems used for this purpose must have check valve and anti-siphon device to prevent back flow into water source of supply tank and must be equipped with power interlock to insure simultaneous shut off of water pump and injector.

ACCEPTED
APR 24 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. 34704-70

