

**PRECAUTIONARY  
STATEMENTS  
HAZARDS TO HUMANS  
& DOMESTIC ANIMALS  
CAUTION**

Harmful if inhaled, swallowed or absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

**STATEMENT OF  
PRACTICAL TREATMENT**

**If Swallowed:** Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

**If Inhaled:** Remove victim to fresh air. Apply artificial respiration if indicated.

**If On Skin:** Remove contaminated clothing and wash affected areas with soap and water. Seek medical attention if irritation occurs.

**If In Eyes:** Flush eyes with water for at least 15 minutes. Get medical attention.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Apply this product only as specified on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**RE-ENTRY STATEMENT**

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

**CAUTION**

(Insert area or field description) treated with Sulfur on (insert date of application). Do not enter without appropriate protective clothing for 24 hours. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Sulfur and describe his conditions. For further information see the STATEMENT OF PRACTICAL TREATMENT portion of the pesticide label.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly.

**STORAGE:** Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**RECOMMENDATIONS**

Use of this material in a manner other than in accordance with this label may cause serious plant, crop, or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

**APPLICATION INSTRUCTIONS**

Unless otherwise specified for specific tree fruits, dosage rates are given as pints of SULFUR 6 FLOWABLE per 100 gallons water for use in a thorough coverage spray.

When dosage rate specifies a range, the lower rate should be used unless the disease population dictates a higher rate is needed for best control. Certain geographic areas may also have need of higher rates. Prior to application, check with local county extension personnel for area rate recommendations.

**APPLE, PEAR—Scab, Pre-Bloom through Calyx sprays—1½ to 3½ pints. Cover sprays—1 to 2 pints. DO NOT apply to D'Anjou or Comice Pears.**  
**POWDERY MILDEW:** Pre-bloom through calyx sprays—2½ to 3 pints. Cover sprays—2 pints. Pear Rust Mite: Use 1 quart per 100 gallons to be applied with oil as a delayed dormant or dormant application.



# SULFUR 6 FLOWABLE

**ACTIVE INGREDIENT**

Sulfur ..... 52.0%

**INERT INGREDIENTS** ..... 48.0%

**TOTAL** ..... 100.0%

(Contains 6.0 pounds Sulfur per gallon)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED** - Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**IF ON SKIN** - Wash thoroughly with soap and water.

**IF IN EYES** - Wash eyes with plenty of clear water for at least 15 minutes.

See side panels for additional precautionary statements.

NET CONTENTS \_\_\_\_\_ U.S. GALLON(S)

EPA REG. NO. 34704-70

EPA EST. NO. \_\_\_\_\_

\* CLEAN CROP is a Registered T.M. of United Agn Products, Inc.

14544

EXP 20

**CHEFRY, PLUM, PRUNE**—Brown Rot, Leafspot, Pink and bloom sprays— $1\frac{1}{2}$  to  $3\frac{1}{2}$  pints. Petal fall, shuck and cover sprays— $\frac{3}{4}$  to  $1\frac{1}{4}$  pints. Not recommended for application on Deacon and Varn varieties of cherries.

**CITRUS** (Grapefruit, lemons, limes, oranges, tangelos and tangerines)—Rust Mite—Apply 1 to 4 pints. Do not apply within 4 weeks of an oil application, unless applied with a dormant oil to winter-dormant woody plants. Thrips (Early Spring treatment for the reduction of Thrips population)—Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush as made 3 to 4 inch growth. Repeat if necessary. Do not apply within 4 weeks of oil application.

**CORN, SORGHUM AND SMALL GRAINS** (Wheat, barley, oats, rye, and flax)—Spider and Common Leaf Spot Mites—Apply 1 gallon per acre in sufficient water to insure thorough coverage when mites appear on bottom 2 leaves. With aerial and concentrate sprayers use 5 to 20 gallons of water per acre. For other ground sprayers, use a minimum of 20 gallons per acre. Repeat applications may be necessary if conditions persist. Can be used in tank mix combinations with other miticides.

**COTTON**—Atlantic Mite—2 pints for early season control and up to 1 gallon for mid to late season control.

**GRAPES**—Powdery Mildew—1 to 2 pints per 100 gallons water, maximum of 8 pints **SULFUR 6 FLOWABLE** per acre in spring and summer. Make first application when shoots are 6 to 8 inches long. Make second application when shoots are 12 to 15 inches long, and third at about bloom stage. The timing, rate, and number of applications will vary with local conditions. Consult State Agricultural Service or State Agricultural Experiment Stations in your region for specific information. Follow the **PLANT RESPONSE PRECAUTIONS** given on this label.

**PEACH, NECTARINES**—Powdery Mildew, Brown Rot, Leafspot; Pink and bloom sprays— $1\frac{1}{2}$  to  $3\frac{1}{2}$  pints. Petal fall, shuck and cover sprays— $\frac{3}{4}$  to  $1\frac{1}{4}$  pints. Application to mature nectarines may cause discoloration.

**PEANUTS**—Rust—2 to 4 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

**POTATOES**—Powdery Mildew—3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3-week intervals.

**STRAWBERRIES**—Powdery Mildew—1 to 2 pints. Apply at first sign of infection and repeat at 3-week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest

in accordance with local canners' recommendations.

**SUGAR BEETS**—Powdery Mildew—Apply the recommended rate in 5 to 10 gallons of water per acre by air at first sign of infection. Use 4 pints per acre in the Pacific Northwest, 8 pints per acre in Arizona and 1 to 4 quarts in Colorado, Kansas, Nebraska and Wyoming. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season.

**TOMATOES**—Tomatoe Russet Mites— $\frac{1}{2}$  to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

**FOR SPRINKLER IRRIGATION SYSTEM APPLICATION (POTATOES AND SUGAR BEETS) INJECT IN SYSTEM AT OR NEAR IRRIGATION PUMP, Center Pivot System**—Apply during last full revolution of the system. Portable or Solid Set Systems—Apply during final 15 minutes of irrigation period. Do not operate irrigation system after application until deposit has thoroughly dried and several days have lapsed. All irrigation systems used for this purpose must have check valve and anti-siphon device to prevent back flow into water source of supply tank and must be equipped with power interlock to insure simultaneous shut off of water pump and injector.

### PLANT RESPONSE PRECAUTIONS

Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature, sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use **SULFUR 6 FLOWABLE** with oil or within 4 weeks of an oil application unless applied with a dormant oil to winter-dormant woody plants.

### DIRECTIONS FOR MIXING

Before Using, Stir Until Smooth

Pour recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. **DO NOT ALLOW MIXTURE TO STAND.** Failure to maintain agitation will cause **SULFUR 6 FLOWABLE** to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of **SULFUR 6 FLOWABLE** act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is corrosive material. **TO REDUCE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY.**

### WARRANTY

1. The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use

when it is used in accordance with such directions, and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and to residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. **THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

2. This warranty does not extend to, and the Buyer shall be solely responsible for, any and all loss or damage which results from the use of this product in any manner which is inconsistent with the label directions, warnings or cautions.
3. **BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

**BEST AVAILABLE COPY**

FORMULATED FOR  
**PLATTE CHEMICAL COMPANY, INC.**  
150 SO. MAIN STREET  
FREMONT, NEBRASKA 68025