

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
DANGER**

**USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING**  
(1) Wear water proof pants, coat, hat, rubber boots or rubber overshoes. (2) Wear safety goggles. (3) Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under provisions of 30 CFR Part II. (4) Wear heavy duty, natural rubber gloves.

**WORK SAFETY RULES**

Keep children and other persons away from treated areas. Do not eat, drink, or smoke in treated areas. Do not get on treated areas. Do not get clothes or shoes wet with spray. Do not get eyes or mouth wet with spray. If you feel sick or dizzy, STOP working and get fresh air. Call a doctor (Physician) immediately for medical advice. Explain that you are working with parathion and present symptoms. After treatment, please see statement on first aid treatment and first aid instructions on label. Do not go to hospital.

**IMPORTANT!** Before removing your work clothes, wash them with soap and water. Avoid contact with your face. Wash your hands and face with soap and water before eating, drinking, or smoking.

**AFTER WORK,** take off work clothes and shoes. Wash your body with soap and water. Wash your clothes when leaving job. Do not wear contaminated work clothes or shoes. Do not use treated equipment without proper washing. Do not use treated equipment without proper washing. Read and follow all instructions on label and on label for proper use.

**NOTE TO PHYSICIAN**

Antidote: Administer atropine intravenously in doses of **TWO TO FOUR** mg. (0.05 to 0.1 mg./lb.) intramuscularly as soon as possible. Repeat doses as necessary. Administer 1 mg. of atropine intravenously every 15 minutes until the patient is fully atropinized. Apply 2 PAM intravenously in doses of 100 mg. every 15 minutes until the patient is fully atropinized. **DO NOT GIVE MORPHINE OR TRANQUILIZERS** until the patient is fully atropinized. Administer 1 mg. of chlorazepate intravenously every 15 minutes until the patient is fully atropinized. Administer 1 mg. of respiratory depressant intravenously every 15 minutes until the patient is fully atropinized. Administer 1 mg. of atropine intravenously every 15 minutes until the patient is fully atropinized. Administer 1 mg. of chlorazepate intravenously every 15 minutes until the patient is fully atropinized. Administer 1 mg. of respiratory depressant intravenously every 15 minutes until the patient is fully atropinized. **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to water. Do not apply where rain, irrigation, or other weather conditions favor drift from areas treated. Do not contaminate water by washing of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended in this label. Do not apply where these are important resources. Apply this product only as specified on this label. Do not apply to water used for irrigation, recreation and domestic purposes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**PHYSICAL & CHEMICAL HAZARDS**

**WARNING: COMBUSTIBLE**

**KEEP AWAY FROM HEAT AND OPEN FLAME**  
**KEEP CONTAINER CLOSED LEAKING PACKAGES SHOULD BE REMOVED TO A SAFE PLACE**  
**DO NOT USE IN UNDILUTED FORM**  
**NOT FOR USE OR STORAGE IN OR AROUND THE HOME**  
**DO NOT STORE AT TEMPERATURES BELOW 55°F**

**POISON SIGNS (Symptoms)**

Methyl Parathion is a very dangerous poison. It is highly toxic to humans and animals. Symptoms of poisoning with Methyl Parathion include: headache, dizziness, blurred vision, pin point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

**POST TREATED AREA**

Consult your State Agricultural Experiment Station for information on post-treatment procedures.

**RESTRICTED USE PESTICIDE  
FOR RETAIL SALE TO AND USE ONLY BY  
CERTIFIED APPLICATORS OR PERSONS  
UNDER THEIR DIRECT SUPERVISION AND  
ONLY FOR THOSE USES COVERED BY THE  
CERTIFIED APPLICATORS CERTIFICATION**

**CLEAN CROP**

**METHYL PARATHION  
7.5**

<b>ACTIVE INGREDIENTS:</b>	
O,O-dimethyl O-p-nitrophenyl phosphorodithioate	7.50%
Related Products	0.00%
Xylene Range Aromatic Solvent	18.50%
<b>INERT INGREDIENTS:</b>	73.50%
<b>Methyl Parathion</b>	<b>7.50%</b>
<b>TOTAL</b>	<b>100.00%</b>

Contains 7.5 lbs. Methyl Parathion per Gallon

**POISON**  
See side panels  
for antidote  
and precautions



**DANGER**  
Keep out of reach  
of children



**STATEMENT OF PRACTICAL TREATMENT**

If Breathing Has Stopped: Administer artificial respiration. If Swallowed: Administer 100 mg. of 2 PAM intravenously every 15 minutes until the patient is fully atropinized. In Case of Contact: Wash with soap and water. Do not use treated equipment without proper washing.