PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER-Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. May be fatal, if swallowed. Avoid breathing vapors.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

STATEMENT OF PRACTICAL TREATMENT

If in eyes flush with plenty of water and get medical attention. If on skin flush with planty of water for at least 15 minutes and get medical attention if irritation persists. Wash contaminated clothing before reuse. Discard contaminated shoes. If swallowed, induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.


For selective control of broadlest weeds in turfgrasses such as those found on lawns and golf courses. Not for use on Centipede, Carpetgrass, St. Augustine, Bentgrass or Dicondra turf or where desirable clovers are present. Newly seeded areas should not be treated until after second mowing. Treated areas, may be reseaded 3 to 4 weeks after treatment. Directions:

Apply as a foliar apray for the control of broadleaf weeds, including dandelion, clovers, henbit, plantain (buckhorn and broadleaf), wild gertic and wild onlon. Best results are obtained when weeds are young and actively growing in the spring and fall. Avoid applications during hot dry weather when weed growth is slow. Do not water for 24 hours after treatment.

Apply at a rate of 2 pints per acre in sufficient spray solution to provide uniform coverage (% to 4 gallons per 100 sq. ft.)

Mixigo Instructions

Area	Rate	Spray Volume
1000 aq. ft.	% oz.	0.5 to 4 gal.
40 00 s q. ft.	3 oz.	2 to 16 gal.
1 acre	2 pints	22 to 174 gal.
	,	



Selective Broadleaf Weed Control in Turfgrasses, Dandellon, Clover, Henbit, Plantains, Wild Onion and Enthur And Many Other Broadleaf Weeds.

DANGER

KEEP OUT OF REACH OF CHILDREN MAY A RESERVED BY THE PROPERTY OF THE PROPE

See Side Panel for Additional Precautionary Statements and Statement of Practical Treatment

NET CONTENTS_ EPA Reg. No. 34704-84 EPA Est. No.

USEFRECAUTIOPS

Avoid use on benaugagress turf in late spring and summe? as discoloration of turf is likely. Do not exceed label rates within the dripline of desirable trees and shrubs. Do not spray exposed roots of ornamental trees and shruffs. Do not allow spray soution of spray drift to contact follage of desirable trees, shrubs, flowers, or vegetables as injury will likely result. Keep from freezing.

FOR CONTAINERS 5 GALLON AND LARGER THE FOILOWING APPLIES

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL (all except household): Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures.

WEEDS CONTROLLED

Pepperweed Spurge

Bedstraw	Henbit
Buckhorn	Knotweed
Burdock	Lambs-
Chickweed	quarters
Clover	Leapedeza
Dendelion	Mallow
Dock	Morning
Ground by	Glory

Piaweed Plantains Pursiane Ragweed Shepherds. Purse Speedwell

Wild Godic Wild Lettuce Wild Onlock Sheep Sorrel Yarrow and many other similar broadleaf weeds

Wild Cerrot

27

