

ACCEPTED

MAR 21 1975

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR AGRICULTURAL PESTS REGISTERED  
EPA REG. NO. 34704-97

# BEN SUL DUST

## DIRECTION FOR USE:

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

THIS PRODUCT IS AN AGRICULTURAL FUNGICIDE FOR THE BELOW INDICATED USES:

Apply when air is calm. Apply when insects or diseases appear. Repeat as necessary. This product is formulated for application by ground or air. Make thorough coverage of plants with a uniform deposit of dust.

**APPLES:** For the control of Scab — Use 10 to 50 lbs. per acre. Begin at pre-pink and continue through cover. Macintosh, Golden Delicious, Jonathan and certain other varieties of Apple may be injured by sulfur.

**ALMONDS:** For control of Scab — Use 40 to 45 lbs. per acre. Begin when disease threatens or in pre-bloom stages. Repeat at 7-day intervals until control is achieved.

**CHERRIES:** For control of Brown Rot, Powdery Mildew — Use 25 to 50 lbs. per acre. Begin in pink stage. Repeat at bloom, petal fall and in cover stages at 7- to 14-day intervals.

**GRAPES:** For control of Powdery Mildew — Use 5 to 20 lbs. per acre. Begin when shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Concord and other labrusca type grapes may be injured by sulfur.

**NECTARINES, PEACHES:** For control of Brown Rot, Powdery Mildew — Use 40 to 50 lbs. per acre. Apply at pink, bloom, petal fall, shuck split, shuck fall and at 10- to 14-day intervals as long as necessary. For Brown Rot of fruit, apply when fruit starts to ripen.

**PLUMS, PRUNES:** For control of Brown Rot, Powdery Mildew — Use 40 to 50 lbs. per acre. Begin at bloom, petal fall, shuck fall, and in cover period as necessary. For Brown Rot of fruit, dust when fruit starts to ripen.

**PEARS:** For control of Scab — Use 40 to 50 lbs. per acre. Apply at pre-bloom, bloom, petal fall and cover periods. Do not use on Anjou or Comice varieties.

**CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines):** For control of Citrus Flat Mite, Citrus Rust Mite, Citrus Thrips and Yuma Spider Mite — Use at least 100 lbs. per acre. Apply when pests first appear and repeat as necessary. Do not apply during or preceding high temperatures. Do not apply within two months of an oil application. FOR USE IN CALIFORNIA ONLY

## (CONTAINS BENTONITE)

ACTIVE INGREDIENT	By Weight
SULFUR .....	81.0%
INERT INGREDIENTS .....	19.0%
TOTAL.....	100.0%

This product is a r correct proportions ing qualities.

Do not permit worl stantial contact wil following indicated Grapes 1, Peaches a

Certain crops and c from sulfur. Do not conditions have prov material to drift ont ation of feed and foc ately prior to periods or higher), as foliag within 3 weeks of an apply when weather c ed. Do not contamin or disposal of waste:

Consult your local Service for additiona

NOTICE TO BUYER— or implied, concernin indicated on the label or handling when su to label instructions.

## CAUTION:

### KEEP OUT OF REACH OF CHILDREN

EPA REGISTRATION NO. 5967-100 AA

EPA EST. 279-CA-2

1.75

MANUFACTURED FOR


### MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

NET WEIGHT 50 BS.

14

# BEN SUL DUST



## IMPORTANT

This product is a mixture of Bentonite and Sulfur in the correct proportions to ensure the best spreading and sticking qualities.

Do not permit workers to enter treated area where substantial contact will be made with treated foliage until following indicated days have lapsed after application: Grapes 1, Peaches and Nectarines 1, Citrus 1.

## CAUTION

Certain crops and other plants are susceptible to injury from sulfur. Do not use Sulfur on any crop unless local conditions have proven its use to be safe. Do not allow this material to drift onto neighboring crops. Avoid contamination of feed and foodstuffs. Do not use during or immediately prior to periods of extreme high temperatures (90° F or higher), as foliage damage may result. Do not apply within 3 weeks of an oil spray or injury may result. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Consult your local Cooperative Agricultural Extension Service for additional recommendations.

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Buyer assumes all risk of use and/or handling when such use and/or handling is contrary to label instructions.

—NON-RETURNABLE—

## (CONTAINS BENTONITE)

ACTIVE INGREDIENT	By Weight
SULFUR .....	81.0%
INERT INGREDIENTS .....	19.0%
TOTAL .....	100.0%

## CAUTION:

**KEEP OUT OF REACH OF CHILDREN**

EPA REGISTRATION NO. 5967-100 AA

EPA EST. 279-CA-2

1-75

MANUFACTURED FOR

**MOYER CHEMICAL COMPANY**

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

NET WEIGHT 50 BS.

**FOR USE:**  
USE ONLY IN ACCORDANCE WITH THE REGULATIONS OF THE APPLICABLE STATE AND FEDERAL AGENCIES.

**FOR USE:**  
Apply by burying with smoke or fumes.  
GENERAL FUNGICIDE FOR

Control insects or diseases on plants. This product is formulated to provide thorough coverage of foliage and fruit.

Use 50 lbs. per acre. Begin at petal fall. Golden Delicious, Jonathan and Jonathon are injured by sulfur.  
Use 45 lbs. per acre. Begin when blossoms open and repeat at 7-day intervals until petal fall.

Powdery Mildew — Use 25 to 50 lbs. per acre. Apply at petal fall and in cover stages.

Use 5 to 20 lbs. per acre. Begin when blossoms open and continue until fruit starts to ripen. Do not use on Anjou or Comice.

Brown Rot, Powdery Mildew — Use 25 to 50 lbs. per acre. Apply at petal fall, bloom, petal fall, shuck split, and as necessary. For Brown Rot use 45 lbs. per acre.

Powdery Mildew — Use 40 to 50 lbs. per acre. Apply at petal fall, bloom, petal fall, shuck split, and as necessary. For Brown Rot use 45 lbs. per acre.

Use 45 lbs. per acre. Apply at petal fall, bloom, petal fall, shuck split, and as necessary. For Brown Rot use 45 lbs. per acre.

Use 45 lbs. per acre. Apply at petal fall, bloom, petal fall, shuck split, and as necessary. For Brown Rot use 45 lbs. per acre.

15