

34704-99 PM15 193

FEB 17 1987

300 / 171875
37 / 4
345 / 156727
173

Ms. Diana G. Williams
Platte Chemical Company
419 18th Street
P.O. Box 667
Greeley, CO 80632

Dear Ms. Williams:

Subject: Amendment - Update Label/Additional Use Rates
Clean Crop Diazinon Dust No. 4
EPA Registration No. 34704-99
Your Applications Dated March 25 and June 24, 1986

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable provided that you make the labeling changes indicated below prior to release for shipment of product bearing the amended labeling.

1. We feel that it would be very difficult for the consumer to measure in 1/6 or 2/5 or 4/5 ounce increments. We suggest that numbers such as these be rounded off. In addition, we do not feel it appropriate to provide application rates based on a per acre basis. ~~This information is generally reserved for agricultural/commercial use products.~~
2. The heading should be revised to read "Directions For Use."
3. ~~The direction for application by air is not appropriate for homeowner use products. Please delete.~~
3. The directions for use for ornamentals should provide additional directions for smaller areas. Perhaps directions based on size of individual or amentals.

2/17/87
THU OK
2/17/87
RFB

91903:Shell:LR-1:KENCO:1/6/87:1/15/87:dej:VO

CONCURRENCES

SYMBOL	SURNAME	DATE					

34704-99 283

2

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

**HAZARDOUS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

Wash with soap and water after handling and before eating or smoking.

Do not store or transport near feed or food.

Do not use on humans or household pets. Keep out of reach of domestic animals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not apply directly to water. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective Information may be obtained from your Cooperative Agricultural Extension Service. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**STATEMENT OF
PRACTICAL TREATMENT**

If swallowed—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled—Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin—Remove contaminated clothing and wash affected areas with soap and water.

If in eyes—Flush eyes with plenty of water. Call a physician if irritation persists.

in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

CAUTION

(Insert area or field description) treated with Diazinon on (insert date of application.) Do not enter treated areas without protective clothing until dusts have settled. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Diazinon and describe his condition. For further information see the "STATEMENT OF PRACTICAL TREATMENT" statement.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

**DIAZINON
DUST**

Controls Aphids and other listed
Vegetables, Fruits and O
as listed in Directions

ACTIVE INGREDIENT

Diazinon [O,O-diethyl O-(2-Isopropyl-4-pyrimidinyl) phosphorothioate].

INERT INGREDIENTS:

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panels For Additional Information

NET WEIGHT _____

EPA REG. NO. 34

EPA EST. NO. _____

tion, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS

Apply when air is calm. Apply when insects appear. Repeat as necessary. This product is formulated for application by ground or air. Make thorough coverage of plants with a uniform deposit of dust.

BEANS (Pole and Snap): For control of Black Bean Aphids, Leafminers, Mexican Bean Beetles - Use 25 to 35 lbs. per acre. Do not apply within 7 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI: For control of Aphids, Diamond Back Moths, Imported Cabbage Worm - Use 20 to 25 lbs. per acre (7½ to 9½ oz. per 1000 sq. ft.). Apply when insects first appear, repeat as necessary. Do not apply within 7 days of harvest on Cabbage or within 5 days of harvest on Cauliflower or Broccoli.

CELERY: For control of Aphids, Flea Beetles - Use 10 to 12 lbs. per acre (3½ to 4½ oz. per 1000 sq. ft.). Apply when insects first appear, repeat as necessary. Do not apply within 10 days of harvest.

CLOVER: For control of Aphids, Leafhoppers - Use 15 lbs. per acre (5½ oz. per 1000 sq. ft.). Do not apply within 7 days of cutting or grazing by animals being finished for slaughter.

CORN: For control of Corn Earworm - Use 30 to 40 lbs. per acre (11 to 14½ oz. per 1000 sq. ft.). Do not apply within 2 days of harvest. Do not feed treated forage to animals being finished for slaughter within 2 days of treatment.

CUCUMBERS: For control of Aphids, Leafhoppers, Thrips - Use 30 to 40 lbs. per acre (11 to 14½ oz. per 1000 sq. ft.). Do not apply within 7 days of harvest.

GRAPES: For control of Pacific Mite, Leafhoppers, - Use 20 to 25 lbs. per acre (7½ to 9½ oz. per 1000 sq. ft.). Do not use within 10 days of harvest.

LETTUCE: For control of Aphids, Dipterous Leafminers - Use 10 to 12 lbs. per acre (3½ to 4½ oz. per 1000 sq. ft.). Apply when insects first appear, repeat as necessary. Do not apply within 10 days of harvest.

LIMA BEANS: For control of Black Bean Aphids, Leafminers, Leafhoppers, Mexican Bean Beetles - Use 25 to 35 lbs. per acre (9½ to 12½ oz. per 1000 sq. ft.). Do not use within 7 days of harvest.

ONIONS (Including Green Onions): For control of Onion Thrips - Use 20 to 30 lbs. per acre (7½ to 11 oz. per 1000 sq. ft.). Do not apply within 10 days of harvest.

PEACHES, NECTARINES: For control of Scale Crawlers, Clover Mite, Pacific Mites - Use 40 to 60 lbs. per acre (14½ to 22 oz. per 1000 sq. ft.). Do not use within 20 days of harvest.

PEPPERS: For control of Aphids - Use 25 to 40 lbs. per acre (9½ to 14½ oz. per 1000 sq. ft.). Apply when insects first appear, repeat as necessary. Do not apply within 5 days of harvest.

SPINACH: For control of Aphids, Dipterous Leafminers - Use 10 to 12 lbs. per acre (3½ to 4½ oz. per 1000 sq. ft.). Apply when insects first appear, repeat as necessary. Do not apply within 10 days of harvest.

STRAWBERRIES: For control of Aphids, Two-Spotted Mites - Use 20 to 30 lbs. per acre (7½ to 11 oz. per 1000 sq. ft.). Apply when insects first appear, repeat as necessary. Thorough underleaf coverage is essential for good control. Do not apply within 5 days of harvest.

TOMATOES: For control of Aphids, Dipterous Leafminers, Cucumber Beetles - Use 30 to 40 lbs. per acre (11 to 14½ oz. per 1000 sq. ft.). Do not apply within 3 days of harvest. If applied within 1 day of harvest - do not use more than 15 to 20 lbs. per acre (5½ to 7½ oz. per 1000 sq. ft.).

ORNAMENTALS (Azalea, Camellia, Carnation, Chrysanthemum, Gladiol, Rose): For control of Aphids, Carnation Bud Mites, Carnation Shoot Mites, Whiteflies - Use 35 to 50 lbs. per acre (12½ to 18½ oz. per 1000 sq. ft.). Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, as injury to plants may occur.

WARRANTY

Platte Chemical Co., Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. The product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or material in or on the soil or crop. Platte Chemical Co., Inc. makes no other express or implied warranty of fitness or merchantability. The exclusive remedy of the user or buyer, and the limit of the liability of Platte Chemical Co., Inc. or any other seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice, which may not be varied by any verbal or written agreement.

FORMULATED FOR:

PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025