

GRASS AND GRASS HAY (0): Aphids, Leafhoppers, and Armyworm - Use 30 to 40 pounds per acre.

LETTUCE (Head 7) (Leaf 14): Aphids, Cabbage Looper, Leafhoppers, Mites - Use 30 to 40 pounds per acre.

MINT (7): Adult Fleabeetles, Caterpillars - Use 20 pounds per acre.

ONIONS (3): Thrips, Onion maggots - Use 30 to 40 pounds per acre.

PASTURE (0), RANGE GRASS (0): Armyworm, Aphids, Leafhoppers - Use 30 to 40 pounds per acre.

PEACHES (7): Plum Curculio, Oriental Fruit Moth, Mites - Use 1/2 to 1 pound per tree.

PEANUTS (0): Leafhoppers - Use 20 pounds per acre.

PEARS (1): Codling Moth, Plum Curculio, Leafhoppers, Flea bug, Mites, Pear Psylla - Use 1/2 to 1 pound per tree.

PEAS: Pea Aphid, Pea Weevil - Use 25 pounds per acre. Do not apply within 7 days of harvest if vines are to be used for feed, if vines are not to be fed may be applied within 3 days of harvest.

PECANS (0): Pecan Nut Casebearer - Use 80 pounds per acre. Apply when first brood larvae begin to enter nuts and repeat after 7 days if necessary. Pecan Bud Moth - Apply 8 pounds per acre by airplane. Apply 4 to 5 times, 2 to 3 weeks apart when insects appear.

PEPPERS (3): Pepper Maggots - Use 30 pounds per acre.

PEPPERMINT (7), SPEARMINT (7): Aphids, Adult Flea Beetles, Climbing Cutworms, Leafhoppers, Cabbage Loopers, Strawberry Root Weevil - Use 20 pounds per acre.

POTATOES (0): Aphids, False Chinch Bug, Leafhoppers, Mealybug - Use 25 to 30 pounds per acre.

SOYBEANS (3): Mexican Bean Beetle - Use 30 to 40 pounds per acre.

STRAWBERRIES (3): Aphids, Field Crickets, Strawberry Leafhopper, Potato Leafhopper, Lygus Bugs, Spittlebug, Twospotted Spider Mites, Thrips, Whitefly - Use 40 pounds per acre.

SUGAR BEETS (3): Aphids, Mites - Use 25 pounds per acre.

SWEET POTATOES (3): Leafhoppers - Use 20 to 25 pounds per acre.

TOBACCO: Aphids: Use 20 to 30 lbs. per acre. No time limitation.

TOMATOES (1): Aphids, Drosophila, Twospotted Spider Mites, Tomato Russet Mite - Use 30 to 40 pounds per acre.

VETCH (0): Pea Aphid, Vetch Bruchid - Use 25 pounds per acre.

WALNUTS (0): Aphids - Use 32 to 48 pounds per acre.

WATERMELONS (1): Cucumber Beetle, Aphids, Leafhoppers, Spider Mite - use 25 to 35 pounds per acre.

#### ORNAMENTALS

Aphids, Spider Mites: Make thorough, full coverage application. Repeat applications as necessary.  
Note: Applications of this material may cause injury

on certain ferns, such as Boston, Maidenhair and Pteris, as well as on certain species of Crassula and on Canada Junipers.

#### ANIMALS DIRECTIONS

POULTRY: Northern Fowl Mite, Poultry Lice: Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary. Treatment of poultry for Northern Fowl Mite should be a supplemental treatment to premise treatment.

PREMISE TREATMENT: Apply 1 lb /50 to 60 sq. ft. of later and floor space and to nests, roosts, and adjacent area.

HOGS: Lice. Treat animals thoroughly. Repeat in 10 days and thereafter as needed.

DOGS AND CATS: Fleas: Direct Application: Apply sufficient dust to treat animal thoroughly. Use any garden type duster, puff duster or shaker-type can. If animal has flea rash, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatment. If the animal is confined to an area free of fleas, one treatment may be sufficient. Note: Do not treat kittens or puppies under 4 weeks of age.

BEEF CATTLE AND NON-LACTATING DAIRY CATTLE: Lice: Direct Application: Treat animal thoroughly. Repeat in 10 days and thereafter as needed. Horn Fly: Use 3 tbsp. per animal. Apply on back and neck of the animal and repeat applications at 10 to 14 day intervals.

Note: Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE: Horn Fly: Direct Application: Use 3 tbsp. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Repeat applications at 10 to 14 day intervals.

NOTE: Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers, and watering troughs. Do not use spray or dip dairy animals in malathion.

HORSES: Horn Flies, Lice, Ticks - Apply by means of a hand or power duster. Treat animal thoroughly and repeat as needed.

GOATS, SHEEP: Keds, Lice, Ticks - Apply by means of a hand or power duster. Treat animal thoroughly and repeat as needed. Do not apply to milk goats. Do not treat animals under one month of age.

#### NOTICE

Pleno warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Pleno. In no case shall Pleno be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Pleno makes no warranties of MERCHANTABILITY or FITNESS for a particular purpose nor any other express or implied warranty except as stated above.

BEST AVAILABLE COPY

2 92

u