

DIRECTIONS FOR USE

Do not use this product in a manner inconsistent with its labeling.

a. Dosages given are suggested only and may not give satisfactory results. Where dosage ranges are indicated, apply larger dosages to spacing, proportionately less for smaller plants and wider spacing. See label for details.

b. Mexican bean beetle and leafhoppers. Apply 25 to 35 lbs. per acre and apply within 24 hours of harvest.

c. CABBAGE, COLLARDS, KALE, MUSTARD and TURNIPS (in home garden and imported cabbage worm. Apply 25 to 50 lbs. per acre every time leaves appear. Do not apply to broccoli or turnips within 3 days of harvest; cauliflower, collards, kale or mustard within 7 days of harvest.

d. (in home garden) to control aphids and cucumber beetle adults. Apply as needed. Do not apply to cucumbers or squash within 1 day of harvest.

e. BEANS and COWPEAS for the control of aphids, apply 25 lbs. per acre as needed. Do not apply within 3 days of harvest. If vines are to be used for silage, apply within 7 days of harvest.

f. For the control of aphids, apply 20 to 30 lbs. per acre to foliage as needed. Repeat every 3 days of harvest.

g. For control of spider mites, apply 25 to 40 lbs. per acre to foliage as needed. Repeat every 1 day of harvest.

h. For control of aphids, lygus bugs, potato leafhoppers, spider mites, and spittle bugs, apply 25 to 40 lbs. per acre to foliage as needed. Do not apply to strawberries within 3 days of harvest. Cyclamen mite is a problem.

i. For control of aphids, apply 1/4 to 1 lb. per 100 square yards of bed. Apply with thorough watering. Do not apply within 3 days of transplanting.

j. For control of aphids, apply 25 to 40 lbs. per acre to foliage, even in home gardens, using sufficient dust to obtain light but thorough, even coverage. Repeat as needed.

k. For control of spider mites, make 3 applications 7 days apart. Repeat at 10-day intervals as necessary.

l. For control of lace bugs; apply as needed.

m. For control of leafhoppers and thrip; apply at 3-day intervals as needed.

n. For control of aphids; apply as needed.

o. For control of aphids in bloom may cause injury to blooms. Do not apply to Hibiscus, Begonia, Maiden Hair, and Pteris, as well as some species of Crassula.

p. For control of aphids on Grapes, Lemons, Limes, Tangelos, and Tangerines for the control of brown scale and aphids, apply with thorough light dusting. Repeat after plants are in bloom. Do not apply within 7 days of harvest.

q. For control of aphids on young or weak animals. Do not apply in conjunction with oral drenches, organic phosphate treatments. Do not contaminate feed, water, feed milking equipment.

r. For control of lice, dust thoroughly into the animals coat. Application should be made into the mouth. If dog is confined, one application may suffice. If animal is in poor condition, application may be repeated in 3 weeks. Do not use on weakened condition. To prevent reinfestation from such areas as pens, move manure and debris, and dust at rate of 2 lbs. per 1,000 square feet as necessary.

s. For control of lice on great animals thoroughly and repeat after 2 or 3 weeks if needed. Do so in this manner.

t. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

u. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

v. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

w. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

x. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

y. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

z. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

aa. For control of horn flies, apply 1 1/2 oz. per animal to the back, neck and flanks. Repeat at 3-day intervals as needed.

the original container and placed in a locked storage area. Dispose of by wrapping the original container in several layers of newspaper in a reuse bag. Discard bag in trash.





MALATHION* 5% DUST INSECTICIDE

ACTIVE INGREDIENT	BY WEIGHT
Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate	5.0%
INERT INGREDIENTS	95.0%
TOTAL	100.0%

*CYTHION[®] insecticide, "The Premium Grade Malathion"[®]

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panels for Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

- If Swallowed - Call a physician or Poison Control Center immediately. Drink on or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
- If Inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
- If On Skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in Eyes - Flush eyes with plenty of water. Call a physician if irritation persists.
- Note to Physician - This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAIA may be effective as an adjunct to atropine.

EPA REG. NO. 34704-105

EPA EST. NO. _____

NET WEIGHT 50 POUNDS

* Cythion is a Registered T.M. of American Cyanamid Co.
 † Premium Grade Malathion is a Registered T.M. of American Cyanamid Co.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
 150 SO. MAIN STREET FREMONT, NEBRASKA 68025

BEST AVAILABLE COPY

Under Section 3 of the Federal Insecticide, Fungicide, and Rodenticide Act, 28 USC 1361, 7 USC 1361, and Executive Order 11737, hereby registered under EPA Reg. No. 34704-105

JAN 23 1989