

## DIRECTIONS FOR LIVESTOCK USE

Apply as a complete coverage spray. Begin applications when flies first appear and repeat at 2 weeks intervals, or as needed, to control adult imitations. Do not spray on feed, food containers, and water.

**CATTLE (BEEF AND NON-MILKING)**—Use 2.0 gallons per 100 pounds of water. Spray calves per 100 pounds of water. Spray cows per 500 pounds of water. Spray heifers per 500 pounds of water. Spray steers per 500 pounds of water. Spray yearlings per 500 pounds of water. Spray adult cows per 100 pounds of water. Spray adult heifers per 100 pounds of water. Spray adult steers per 100 pounds of water. Spray adult yearlings per 100 pounds of water. Repeat every 2 weeks if necessary.

**HOGS**—Use 2.0 gallons per 100 pounds of water. Spray adult hogs per 100 pounds of water. Spray adult sows per 100 pounds of water. Spray adult boars per 100 pounds of water. Repeat every 2 weeks if necessary.

**SHEEP/GOATS**—Use 2.0 gallons per 100 pounds of water. Spray adult sheep per 100 pounds of water. Spray adult goats per 100 pounds of water. Repeat every 2 weeks if necessary.

**POULTRY—DIRECT APPLICATION**—Use 2.0 gallons per 100 pounds of water. Spray adult chickens per 100 pounds of water. Spray adult turkeys per 100 pounds of water. Repeat every 2 weeks if necessary.

**POULTRY HOUSE**—Use 2.0 gallons per 100 pounds of water. Spray poultry house per 100 pounds of water. Repeat every 2 weeks if necessary.

**ROOST PAINT**—Use 2.0 gallons per 100 pounds of water. Spray roost paint per 100 pounds of water. Repeat every 2 weeks if necessary.

## DIRECTIONS FOR MOSQUITO CONTROL

Use 2.0 gallons per 100 pounds of water. Spray mosquitoes per 100 pounds of water. Repeat every 2 weeks if necessary.

## DIRECTIONS FOR FLY CONTROL USE

Use in and around buildings which house domestic animals, animal yards and around homes. Apply to surfaces where fly adult or congregate, such as walls, stanchions, barns, fences and garbage cans. Repeat application as required. Eliminate fly breeders. Wash floors with freshly white washed surfaces for period of 14 days. Do not contaminate drinking water.

**WATER DILUTED FLY SPRAY**—Use 2.0 gallons per 100 pounds of water. Spray the diluted mix at the rate of 1.0 gallon per 1000 square feet on painted surfaces. Use 1.0 gallon per 1000 square feet on unpainted surfaces.

**BAIT SPRAYS**—Use 2.0 gallons of Maathon Maathon per 1000 square feet of area of mixed surfaces. Use 2.0 gallons of Maathon Maathon per 1000 square feet of area of mixed surfaces. Use 2.0 gallons of Maathon Maathon per 1000 square feet of area of mixed surfaces. Use 2.0 gallons of Maathon Maathon per 1000 square feet of area of mixed surfaces.

Do not spray on surfaces which are treated with other insecticides. Do not spray on surfaces which are treated with other insecticides. Do not spray on surfaces which are treated with other insecticides. Do not spray on surfaces which are treated with other insecticides.

Do not spray on surfaces which are treated with other insecticides. Do not spray on surfaces which are treated with other insecticides. Do not spray on surfaces which are treated with other insecticides. Do not spray on surfaces which are treated with other insecticides.

## DISPOSAL WARNING

### DESTROY CONTAINERS WHEN EMPTY

Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers.

### DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers.

Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers.

**BEST AVAILABLE COPY**



**BLUEBERRIES (1)** Japanese Beetle — Use 1½ pints per acre  
Cherry Fruitworm, Cranberry Fruitworm — Use 1 ½ pints per 200  
gallons per acre. Plum Curculio, Sharp nosed Leafhopper — Use  
2 ½ to 3 2 pints per acre

**BROCCOLI (3)**, BRUSSELS SPROUTS (7), CAULIFLOWER (7),  
CABBAGE (7), KALE (7), ROMAINE (7), HOBBLEHEAD (7), LETTUCE  
(13), MUSTARD GREENS (7), FAN LEAF (7), FAN BEET (7),  
RAGINIES (7), ROTABAGAS (7), SALSIFY (7), WILD CHARD (7),  
TURNIPS (3) Aphids, Imported Cabbag Worm, Cabbage Looper  
— Use 1 to 2 pints per acre

**CANTALOUPE (1)** Cucumber Beetle — Use 2 pints per acre  
Leafhoppers, Pickworm — Use 1 to 2 pints per acre. Aphids,  
Spider Mites — Use 1½ pints per 100 gallons

**CARROTS (7)** Aphids — Use 1½ to 2 pints per 100 gallons  
Leafhoppers — Use 2½ pints per acre

**CELERY (7)** Spider Mites, Aphids — Use 1½ pints per acre

**CHERRIES (3)** Bark Cherry Aphid, Fruit Tree Leafminer — Use 1½  
pints. Cherry Fruit Fly, European Fruit Moth — Use 1 pint. Injury  
may occur on certain varieties of Sweet Cherries particularly in the  
Northwest

**CHESTNUTS (10)** Mites — Use ½ pint

**CITRUS — GRAPEFRUIT (2), LEMON (1), LIME (2), ORANGES  
(7), TANGERINE (5)** Citrus Fruit Aphid, Citrus Red Scale, Purple  
Scale, Black Scale, Citrus Scale, Citrus Leafminer — Use 1½ pints  
Red Scale, Purple Scale — Use 1½ pints per 100 gallons of water  
for light infestation and 2 pints per 100 gallons for moderate and  
heavy infestation. Thrips — Use 1½ pints per 100 gallons of water  
per acre. Do not apply during bloom

**CUCUMBERS (10)** Aphids, Leafminers, Cucumber Beetle — Use 1½ to 2 pints  
per acre. Apply for aphids in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

**ALFALFA (10)** Grasshoppers, Pea Aphid, Spotted Alfalfa Aphid,  
Lygus Bugs, Leafhoppers, Alfalfa Weevil Larvae — Use 1½ to 2½  
pints per acre. Armyworms — Use 2 to 2½ pints per acre. Apply  
to Alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

**ALMONDS (10)** Aphids, Branch Twig Borer, Red Scale, Pear Mite — Use  
1 2 to 2 ½ pints per acre

**APPLES (13)** Codling Moth — Use 2 pints. Eye-spotted Bud Moth,  
Furrowed Green Apple Aphid — Use 1 pint. Tent Caterpillars — Use  
1½ to 2 pints. Mealybug — Use 1 to 2 pints. Green Apple Aphid,  
Russet Apple Aphid — Use 1½ pints. For Codling Moth control apply  
at pre-bloom and every 10 to 14 days thereafter until control is  
achieved. Applications of this material may cause injury of McIntosh  
and related varieties of apples

**APPLES (10)** Orange Tortrix, Soft Scale — Use 2 pints

**ASPARAGUS (5)** Asparagus Aphid, Asparagus Beetle, Thrips —  
Use 1½ to 2 pints per acre

**BANANA (1), CANTALOUPE (1), RYE (7), WHEAT (7)** Young Grasshopper,  
Green Leafhopper, Green Aphid — Use 1½ to 2 pints per acre  
Armyworms — Use 2 pints per acre

**BAYONNETTE (1)** Aphids, Spider Bees — Use 2 to 2 ½ pints per  
acre. Mites, Pear Thrips, Leafhoppers — Use 1 2 to 2 ½ pints  
per acre. Armyworm, Pear Thrips — California — Use 1 2 to 2 ½ pints  
per acre

**BERRIES (2)** Aphids — Use 1½ to 2 pints per acre

**BROCCOLI (3)** Aphids, Imported Cabbag Worm, Cabbage Looper — Use 1 to 2  
pints per acre. Leafminers, Aphids — Use 1 to 2 pints per acre  
Cucumber Beetle, Pear Thrips — Use 1 2 to 2 ½ pints per  
acre

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

Apply to alfalfa in bloom only in the evening or early morning when bees  
are not working in the field or are not hanging on the outside of  
hives

WEST VALLEY 1987

