

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION: Harmful If Swallowed:** Avoid contact with skin. Wash thoroughly after using. Avoid breathing of dust. Avoid contamination of feed and foodstuff. Keep out of reach of children. Keep container closed and away from heat and open flame.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

### PHYSICAL & CHEMICAL HAZARDS

Keep container closed and away from heat and open flame.

### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed:** Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled:** Remove victim to fresh air. Apply artificial respiration if indicated.

**If On Skin:** Remove contaminated clothing and wash affected areas with soap and water.

**If In Eyes:** Flush eyes with plenty of water. Call a physician if irritation persists.


**Note to Physician:** This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

- 1. Prohibitions** — Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2. Pesticide Disposal**—Pesticide, spray mixture, or rinse water that cannot be used, according to label instructions, must be disposed of according to applicable Federal, State, or Local procedures.
- 3. Container Disposal**—Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and Local authorities.
- 4. General** — Consult Federal, State or Local Disposal Authorities for approved alternative procedures.



### ACTIVE INGREDIENT:

Malathion\*

\*O, O-dimethyl dithiophosphate of diethyl

mercaptosuccinate . . . . . 6%

INERT INGREDIENT: . . . . . 94%

TOTAL 100%

**Keep Out Of Reach Of Children**  
**CAUTION**

(See side panel for additional precautions)

### FOR INSECT CONTROL ON STORED GRAINS

(Barley, Field Corn, Pop Corn, Grain Sorghum, Oats, Rice, Rye, Sunflowers, Wheat)

### SEEDS

(Field or Garden)

### PEANUTS

NOT FOR USE IN OR AROUND THE HOME

EPA Reg. No. 34704-110

EPA Est. No. 34704-CO-1

Net Contents

\* CLEAN CROP is a registered trademark of Union Carbide Corporation.

Prior to storing treated grain, seeds, or peanuts, clean the storage area (walls and floors) and the truck or wagon used to transport and convey the grain.

Treatment as outlined below will control confused flour beetle, flat grain beetle, grain borer, maize weevil, red flour beetle, saw toothed grain beetle, and Indian meal moth.

It is important to achieve thorough coverage of seeds, or peanuts when going into storage. The storage may result in insect infestation.

**GRAIN, SEEDS, OR PEANUTS CONTROL TREATMENT METHOD** The most effective method of applying 6% MALATHION GRAIN PROTECTOR is to spray the truck or wagon prior to binning. Spray the truck or wagon in this manner: Measure the depth (all in feet) and multiply by the number of bushels.

Apply 6% MALATHION GRAIN PROTECTOR to seeds, or peanuts at the rate given on this label when crop is being loaded or turned into bins.

Spread the PROTECTOR evenly over the seeds, or peanuts in the truck or wagon.

PROTECTOR will become mixed as it is loaded into the auger hopper. Further mixing of grain, seeds, or peanuts is augmented.

Mixing is completed as the PROTECTOR is loaded into the cone shaped hopper. PEANUTS ARE STORED. This treatment will control Indian meal moth.

Apply the PROTECTOR to clean grain, seeds, or peanuts as outlined above. Apply a second treatment at 2-month intervals.

Platte warrants that this product is safe when used as stated on such label only when used in accordance with the risks inherently associated with the use of this product because of such factors as weight, handling of this product. All such risks are beyond the control of Platte. In no event shall Platte be liable for any particular purpose nor any other use.



FORMULATED FOR **PLATTE CHEMICAL CO., Inc.**  
150 SO. MAIN STREET FREMONT, NEBRASKA 68402

# 6% MALATHION GRAIN PROTECTOR

**INGREDIENT:**

Dimethyl dithiophosphate of diethyl  
phthalosuccinate ..... 6%  
**INGREDIENT:** ..... 94%  
**TOTAL** ..... 100%

## Out Of Reach Of Children CAUTION

(See panel for additional precautions)

### INSECT CONTROL ON STORED GRAINS

Corn, Grain Sorghum, Oats, Rice, Rye, Sunflowers, Wheat)

**SEEDS**  
(Field or Garden)

**PEANUTS**

### FOR USE IN OR AROUND THE HOME

Pat. No. 34704-110

Pat. No. 34704-CO-1

**Net Contents**

U.S. Reg. U.S. Pat. & T.M. Office of United States Dept. of Agriculture

**PLATTE CHEMICAL CO., Inc.**  
150 SO. MAIN STREET, FREMONT, NEBRASKA 68102

Prior to storing treated grain, seeds, or peanuts, thoroughly clean the storage area (walls and floor), as well as equipment used to transport and convey the grain, seeds, or peanuts.

Treatment as outlined below will control cereal leaf beetle, confused flour beetle, flat grain beetle, granary weevil, lesser grain borer, maize weevil, red flour beetle, rice weevil, rusty grain beetle, saw-toothed grain beetle and mid in the control of Indian meal moth.

It is important to achieve thorough coverage of the grain, seeds, or peanuts when going into storage. Untreated areas in the storage may result in insect infestations.

**GRAIN, SEEDS, OR PEANUTS GOING INTO STORAGE TREATMENT METHOD:** The most convenient way to apply 6% MALATHION GRAIN PROTECTOR is to treat it when it is in the truck or wagon prior to binning. Determine the amount in the truck or wagon in this manner: multiply the length by width by depth (all in feet) and multiply this total by 0.8 to get the number of bushels.

Apply 6% MALATHION GRAIN PROTECTOR to the grain, seeds, or peanuts at the rate given in the below tables as the crop is being loaded or turned into final storage.

Spread the PROTECTOR evenly over the surface of the grain, seeds, or peanuts in the truck or wagon. Cut the PROTECTOR into the grain, seeds, or peanuts with a shovel. The PROTECTOR will become mixed as it is dumped from the truck into the auger hopper. Further mixing will occur when the grain, seeds, or peanuts is augered or blown into the bin. Mixing is completed as the PROTECTOR and grain, seeds, or peanuts roll down the cone shaped pile in the bin.

**SURFACE TREATMENT AFTER GRAIN, SEEDS OR PEANUTS ARE STORED:** This treatment is an aid in the control of Indian meal moth. Apply the PROTECTOR to the surface of clean grain, seeds, or peanuts as soon as bins are filled and leveled. Apply a second treatment one month later and subsequent treatments at 2 month intervals, as needed.

#### APPLICATION RATES

PEANUTS	
<b>STORAGE TREATMENT</b> 2.6 lbs. per 1,000 sq. ft.	<b>SURFACE TREATMENT</b> 7.2 lbs. per 1,000 sq. ft.

#### APPLICATION RATES

GRAINS, FIELD OR GARDEN SEEDS	
<b>STORAGE TREATMENT</b> 10.4 lbs. per 1,000 bushels	<b>SURFACE TREATMENT</b> 5.2 lbs. per 1,000 sq. ft.

#### IMPORTANT

Use of this product, as directed will be within the tolerance of 8 parts per million for Malathion. TREATED GRAIN IS NOT SUBJECT TO DOWNGRADING AND CAN BE MILLED INTO FLOUR OR FEED USING NORMAL PRODUCTION METHODS.

**STORED ALMONDS (In Shell and Shelled):** For the control of Indian Meal Moth and Merchant Grain Beetle, apply 2.6 lbs. per 10,000 lbs. of Almonds. Thorough coverage of Almonds is important.

**BAGGED CITRUS PULP:** For the control of Almond Moth, Angoumois Grain Beetle, Cigarette Beetle, Confused Flour Beetle, Flat Grain Beetle, Indian Meal Moth, Mediterranean Flour Moth, Red Flour Beetle, and Saw Toothed Grain Beetle. Apply at the rate of 6 1/2 lbs. per 1,000 sq. ft. of exposed bag surface area when bagged citrus pulp is stored. Make 2 separate applications initially when bagged citrus pulp is stored. Once each month thereafter throughout storage period apply 4 1/2 lbs. per 1,000 sq. ft. of exposed bag surface area.

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



34704-110



12702