

Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...

**ENVIRONMENTAL HAZARDS**

Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...  
 Do not feed treated ... Do not feed treated ...

**STATEMENT OF PRACTICAL TREATMENT**

- Swallowed ...
- Inhaled ...
- On Skin ...
- In Eyes ...

**DIRECTIONS FOR USE**

Do not use ... Do not use ...  
 Do not use ... Do not use ...  
 Do not use ... Do not use ...

**PESTICIDE DISPOSAL** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL** Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.  
**HOUSEHOLD USE DISPOSAL** (Do not reuse container bottles as jars) Rinse thoroughly with water, discard in trash. Securely wrap original container in newspaper, if newspaper and discard in trash.

**GENERAL INFORMATION**

MSMA Turf Herbicide is a postemergence herbicide for use on turf and certain non-crop areas. MSMA Turf Herbicide controls a wide range of turf and non-crop weeds, including: ...  
 A partial list of weeds controlled with MSMA Turf Herbicide includes:  
 Bermudagrass, Crabgrass, Field bindweed, Bromegrass, Annual bluegrass, Wild oats, Annual ryegrass, Kentucky bluegrass, Perennial ryegrass, Annual bluegrass, Perennial ryegrass, Annual ryegrass, Kentucky bluegrass, Perennial ryegrass, Annual bluegrass, Perennial ryegrass.

**MIXING INSTRUCTIONS** MSMA Turf Herbicide is a postemergence herbicide for use on turf and certain non-crop areas. MSMA Turf Herbicide controls a wide range of turf and non-crop weeds, including: ...

**MSMA Turf Herbicide**

A Postemergence Herbicide for use on turf & certain non-crop areas

ACTIVE INGREDIENT:

Monosodium Acid Methanearsonate 48.3%  
 INERT INGREDIENTS 51.6%

TOTAL 100.0%

Total Arsenic, all in water-soluble form, expressed as elemental 22.3%

Product Contains 6.0 Pounds of MSMA Per Gallon

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

See Side Panels for Additional Precautionary Statements

EPA REG NO 34704-114

EPA EST NO 34704-MS

NET CONTENTS 1 GALLON

EPA ...  
 EPA ...  
 EPA ...

a low-volume, properly calibrated sprayer having satisfactory pumping and bypass action. Adjust nozzles in a manner to allow maximum coverage of weeds. Adequate coverage is very important for effective weed control. DO NOT apply with hose-end applicators. Do not apply this product through any type of irrigation system.

### DIRECTIONS

**GOLF COURSE and ORNAMENTAL TURFGRASS** MSMA Turf Herbicide can be used for selective control of Bahiagrass, Barnyardgrass, Chickweed, Smooth and Large Crabgrass, Dallisgrass, Nutsedge, Sand bur, and Wood Sorrel with little or no injury to well established, actively growing turfgrasses. Mow turfgrass to a height of 1 to 1½ inches before treatment. Mix at a rate of 2½ to 2¾ pints in 100 gallons of water for application to one acre. For small areas, mix 1 fluid ounce (2 tablespoons) in ½ gallons of water for application to an area of 1,000 sq. ft. Application should be uniform and thorough to adequately wet all undesirable plants.

Two or more repeat treatments at 14-day intervals may be necessary. Make applications during warm weather when temperature is between 80° and 90°. DO NOT water turf for at least 24 hours after application. Turfgrasses may be temporarily discolored. Bermudagrass, Bluegrass, and Zoysiagrass have shown tolerance to properly applied MSMA. Injury may result if applied to bentgrasses and Fescues. DO NOT apply to St. Augustinegrass, Carpetgrass, Centipedegrass, or to Dichondra lawns. DO NOT reseed until 2 weeks after last application.

**NON-CROP** MSMA Turf Herbicide is effective in control of the above listed weeds and many similar weeds on drainage ditch banks, fence rows, sand traps, storage yards and similar non-crop areas. Application should be made when weeds are small and conditions are favorable for grass growth. Mix at a rate of 2½ to 2¾ pints of MSMA in 40 gallons of water for application to one acre. Spray undesirable vegetation thoroughly to point of runoff. Adequate coverage and complete wetting of foliage is important for effective control. Repeat applications may be necessary if regrowth occurs. Use only as spot treatment in Florida.

which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

NOT REVIEWED

In Accordance with PR Notice 82-20  
Based on Draft Labeling Dated

FORMULATED FOR  
**PLATTE CHEMICAL COMPANY, INC.**  
150 SO. MAIN STREET  
FREMONT, NEBRASKA 68025