



034704-00120-050399

Printed: 15:52:23 Monday, 24 May, 1999 # 15 / 3019

Systems Integration Group, Inc.

PM 23

34704-120

5-3-99

1 of 7

MAY 3 1999

Ms. Glenda Haage
 Platte Chemical Company
 P.O. Box 667
 Greeley, CO 80632

Dear Ms. Haage:

Subject: Label Amendment Extending 2,4-D Use in Reduced or No-Tillage Soybeans
 (Pre-plant Only)
 Clean Crop Amine 4 2,4-D Weed Killer
 EPA Reg. No. 34704-120
 Your Application Dated April 8, 1999

The Agency is conditionally approving an amendment to the registration of the above-referenced product under the authority of section 3(c)(7)(B) of the Federal Insecticide, Fungicide Act (FIFRA). This amendment allows use of the subject product on reduced or no-tillage soybeans (pre-plant only) with a maximum permissible level for residues of the herbicide in or on soybeans of 0.1 ppm. This amendment will expire automatically on December 31, 2001. In addition, during the period that this amendment is effective, it will be subject to the conditions listed below:

- 1) This acceptance is based on your certification that the submitted labeling on preplant soybean use is an exact copy of the labeling accepted by the Agency on September 18, 1998. Please note that the acceptance of this labeling pertains to the previously approved preplant soybean use only and does not signify the acceptance of any other labeling revisions. No other label revisions are being reviewed or considered with this action. You should also note that if you fail to satisfy the conditions imposed in this certification, EPA may issue a notice to cancel this amendment under FIFRA section 6(e).
- 2) This conditional registration will expire automatically on December 31, 2001. Sale or distribution of the subject product bearing labeling for this use on reduced or no-tillage soybeans (pre-plant only) after December 31, 2001 will be illegal. The tolerance authorizing residues of the subject product will also expire on December 31, 2001. After that date, sale or distribution of food in interstate commerce containing any residue of the subject product will be a violation of the Federal Food, Drug, and Cosmetic Act.

RD:STANTON:PM Team 23:Rm. 237:CM-2:305-5218:Disk #10:S561302

CONCURRENCES

SYMBOL ▶	7505C						
SURNAME ▶	S. Stanton						
DATE ▶	May 3, 1999						

- 3) Finally, if and when a permanent tolerance is established, EPA will entertain an application to amend the registration of the subject product without any special limitations on the duration of the amendment.

A stamped copy of the labeling is enclosed for your records. Please submit one copy of your final printed labeling before you release the product for shipment.

Sincerely yours,



Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

3 of 7



AMINE 4

2,4-D WEED KILLER

For Selective Broadleaf Weed Control in Certain Crops, Turf and Non-Crop Areas.

ACTIVE INGREDIENT:

*Dimethylamine salt of 2,4-Dichloro-phenoxyacetic acid	46.5%
INERT INGREDIENTS:	53.5%
TOTAL	100.0%

*Equivalent to 38.6% 2,4-D acid or 3.74 pounds per gallon.
*Isomer specific by AOAC Method No. 6.275-6.279 (13th Ed.)

USER SAFETY RECOMMENDATIONS

Users should:
Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

(See Below for Additional Precautionary Statements)

EPA REG. NO. 34704-120

EPA EST. NO. _____

NET CONTENTS 2½ GAL. (9.46 L)

IHT 01P99

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not contaminate water used for irrigation, domestic or spray purposes.

Personal Protective Equipment:

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eye-wear. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washing, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

For containers over 1 gallon and less than 5 gallons in capacity: Mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to other required PPE.

Engineering controls statements:

When handlers use enclosed cabs or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (5-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

For containers of 5 gallons or more in capacity: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

MAY 3 1999

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No:
34704-120**

STATEMENT OF PRACTICAL TREATMENT

If in Eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.
If Swallowed: Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.
If on Skin: Wash with plenty of soap and water. Get medical attention.
If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
Note to Physician: If in eyes, specialized ophthalmologic attention may be necessary. If swallowed, probable mucosal damage may contraindicate gastric lavage. There is no specific antidote; treat symptomatically.
FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 812-851-8180, EXT. 136.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Groundwater Contamination:

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Do not apply when weather conditions favor drift from treated areas. Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not use in or near a greenhouse.

Do not contaminate water used for irrigation or domestic purposes (except as specifically recommended on this label) especially in areas where susceptible plants are grown.

Do not treat irrigation ditches in areas where water will be used to overhead (sprinkler) irrigate susceptible crops.

Do not apply AMINE 4 directly to, or permit mist to drift onto, sensitive crops or ornamental plants which are susceptible to 2,4-D herbicide. Crops contacted by AMINE 4 sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Do not apply when a temperature air inversion exists. Such a condition is characterized by little or no air movement and an increase in air temperature with an increase in height. In humid regions a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. If questions exist pertaining to the existence of an inversion consult with local weather services before making application.

Use coarse sprays to minimize drift. Do not apply with hollow cone type insecticide or other nozzle that produce fine spray droplets.

AMINE 4 2,4-D WEED KILLER

EPA REG. NO. 34704-120

407

Drift from aerial or ground application may be reduced by: (1) applying as near the target as possible in order to obtain coverage; (2) by increasing the volume of spray mix per acre; (3) by decreasing the pounds of pressure at the nozzle tips; (4) by using nozzles which produce a coarse spray pattern; (5) by not applying when wind is blowing toward susceptible crops or valuable plants.

Applications by airplane, ground rigs, and hand dispensers should be carried out only when there is no hazard from drift. Do not spray when the wind is blowing across the area to be sprayed towards susceptible crops or ornamental plants. Violent windstorms may move soil particles. If 2,4-D is on these particles and they are blown onto susceptible plants, visible symptoms may appear. Serious injury is unlikely. The hazard of movement of 2,4-D on dust during violent windstorms is reduced if treated fields are irrigated or if rain occurs shortly after application.

Large amounts of 2,4-D in the soil may temporarily inhibit seed germination. Do not use in any manner other than recommended on the label to avoid possible crop injury or residues at harvest.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI), of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

USE REQUIREMENTS FOR PASTURE, RANGELAND AND NON-CROP AREAS OTHER THAN TURF: Do not enter treatment areas until spray has dried or dust has settled. For early entry to treatment areas, wear eye protection, chemical resistant gloves, long sleeved shirt, long pants, sock and shoes.

TURF USE REQUIREMENTS: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

NOTE: For application to turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes, follow AGRICULTURAL USE REQUIREMENTS on this label.

STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Do not store below temperature of 25° F. If frozen, warm to 70° F. and redissolve before using by rolling or shaking the container. Store in a safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Storage & Disposal Cont'd.:

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPRAY PREPARATIONS

Mix AMINE 4 only with water to obtain adequate coverage, unless otherwise directed on this label. Add about half the water to the mixing tank, then add the AMINE 4 with agitation, and finally the rest of the water with continuing agitation. Note: Adding oil, wetting agent, or other surfactant to the spray may increase effectiveness on weeds, but also may reduce selectivity to crops resulting in crop damage.

APPLICATION PROCEDURES

Generally, the lower dosages given will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply AMINE 4 during warm weather when weeds are young and growing actively. Apply a minimum of 2 gallons of spray solution per acre by air or a minimum of 5 gallons of spray solution per acre by ground unless directed otherwise.

Do not apply this product through any type of irrigation system.

PLANTING IN TREATED AREAS

Labeled Crops: Crops listed as use sites on this or other registered 2,4-D labels may be planted within 29 days of AMINE 4 application. Follow more specific limitations (if listed) provided in the directions for individual crops. Labeled crops may be at risk for crop injury or loss when planted soon after application, especially in the first 14 days.

Degradation factors described below should be considered in weighing this risk.

Other Crops: All other crops may be planted 30 or more days following an application without concern for illegal residues in the planted crop.

However, under certain conditions, there may be a risk of injury to susceptible crops. Degradation factors described below should be considered in weighing this risk. Under normal conditions, any crop may be planted without risk of injury if at least 90 days of soil temperatures above freezing have elapsed since application.

Degradation factors: When planting into treated areas, the risk of crop injury is less if lower rates of product were applied and conditions following application have included warm, moist soil conditions that favor rapid degradation of 2,4-D. Risk is greater if higher rates of product were applied and soil temperatures have been cold and/or soils have been excessively wet or dry in the days following application.

Consult your local Agricultural Extension Service for information about susceptible crops and typical soil conditions in your area.

WEED LIST

The degree of control is dependent upon the species, stage of growth and overall growing conditions. Best results are obtained when weeds are young and actively growing. AMINE 4 may be applied to control the following listed weeds.

Arichoke	Hairy galinsoga	Ragweed (common)
Aster	Heal-all	Rough fleabane
Austrian Fieldcress	Hoary Cress*	Russian Thistle*
Beggarticks*	Horse Tail	Salsify
Bindweed*	Iron Weed*	Sicklepod
Bittersweet	Jerusalem artichoke	Smartweed*
Bitterweed	Jimson Weed	Sowthistle
Blue Lettuce	Knotweed*	Spanishneedles
Broomweed	Lambsquarters	Stinging Nettles
Bull Thistle	Lettuce (wild)	Strawberry (wild)
Burdock	Loco Weed	Sunflower
Canada Thistle*	Mallow*	Tansy Mustard
Catnip	Many flowered aster	Tan Weed
Chicory	Marsh Elder	Toadflax
Cockle	Morningglory	Tumble Weed
Cocklebur	Musk Thistle	Velvet leaf
Coffee Weed	Mustard*	Venice Mallow
Common evening primrose	Nettles**	Vervains*
Croton	Orange hawkweed	Vetch
Dandelion	Parsnip	Wild Carrot
Docks*	Pennywort	Wild Garlic*
Dogbane*	Peppergrass	Wild Lettuce
Fleabane (Daisy)	Pigweed**	Wild Onion*
Flix Weed	Plantains	Wild Parsnips
Frenchweed	Poverty Weed	Wild Radish
Galinsoga	Prickly Lettuce	Wild Rape
Goatsbeard	Primrose	Wild Sweet Potato
Goldenrod*	Puncturevine	Worm Wood
Ground Ivy*	Radish	
	Ragweed	

*These species may require repeated applications or use the higher rate recommended on this label.

**Control of these species in areas which are locally resistant, may not be satisfactory with this product.

AMINE 4 2,4-D WEED KILLER

EPA REG. NO. 34704-120

5 of 7

APPLE AND PEAR ORCHARDS—Non-Bearing trees (well established, one year or older) and Bearing Trees before and after bloom: Use 3 pts. product in 20 to 50 gals. of water per acre of ground sprayed. For band or spot treatment calculate rates according to the actual portion of an acre treated. Apply as a directed spray onto the weeds to point of runoff when weeds are young and actively growing (pre-bud to early stage). A maximum of 2 applications per season may be applied with a minimum retreatment interval of 75 days. Do not apply within 14 days of harvest.

GRAPE VINEYARDS (Established at least 3 years)
For control of field bindweed, morning glory, Canada thistle and other 2,4-D susceptible broadleaf weeds, apply 2 to 3 pts. AMINE 4 in 10 to 80 gallons of water per acre. For band or spot treatment, calculate rates according to the actual portion of an acre treated. Apply AMINE 4 up to the point of bud break only. Apply when weeds are in the bud to early bloom stage and growing vigorously. Use a hooded boom and low pressure flooding nozzles, where appropriate, to minimize potential of nontarget contact.
Precaution: Grapes are extremely sensitive to 2,4-D. Refer questions, regarding rate and application, to an Agricultural Extension Service specialist.

STONE FRUIT AND NUT ORCHARDS (including pistachios)—For control of annual broadleaf weeds in the orchard floor, apply 3 pints product per acre. Apply using coarse sprays and low pressure in sufficient volume of water to obtain thorough wetting of weeds. Do not make more than two applications per year. Do not harvest stone fruit within 40 days of application or harvest nuts within 60 days of application.

Precautions in applying Amine 4 in Orchards and Vineyards
When applying AMINE 4 in orchards or vineyards, apply only after irrigation and allow maximum time before the next irrigation. Do not apply around fruit trees or vines with handgun. Use only flat, fan-typed nozzles and low pressure—20 to 30 lbs. Use a fixed boom application which can be calibrated and will deposit the spray uniformly. Apply precisely and uniformly to prevent damage to the trees and vines and to obtain satisfactory weed control. Do not apply during windy periods or extremely high temperature. Do not use on light sandy soil. Application to bare ground may result in injury. Do not allow spray to drift or contact foliage, fruit, vines, stems, trunk of trees, or exposed roots as injury may result. Trees must be at least 1 year old and in vigorous conditions before application is made. Do not apply during bloom. Do not graze or feed cover crops from treated orchards or vines.

ASPARAGUS
Apply 3 to 4 pts. AMINE 4 in about 60 gallons of water per acre for ground application and 12 gallons per acre for air application. Apply on actively growing weeds, usually in April or May. If spears are present, treat immediately after cutting. Make no more than 2 applications during the harvest season and these should be spaced at least one month apart. Spears contacted by the spray may be malformed and off-flavored. If malformed, spears should be cut immediately and discarded. Post-harvest spraying should be only by ground rig using drop nozzles to avoid spraying the fern.

WEED CONTROL IN SMALL GRAINS NOT UNDERSEEDED WITH A LEGUME (Barley, Oats, Rye, Wheat) See Table for recommended use rates. Spray after grain begins tillering and before the boot stage (usually 4 to 8 inches tall) and weeds are small. Do not apply before the tiller stage nor from early boot through the milk stage. To control large weeds that will interfere with harvest or to suppress perennial weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.
Note: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock.

WEED CONTROL IN CORN: See Table for recommended use rates.
Preemergence - Apply to soil any time after planting but before corn emerges. Do not use on very light, sandy soil. **Emergence:** Apply just as corn plants are breaking ground. **Postemergence** - Apply to emerged corn. When corn is over 8 inches tall or the fifth leaf collar is visible, whichever occurs first, use drop nozzles to keep spray off corn foliage. Do not apply from 7 to 10 days before tasseling to dough stage. Injury to corn is most likely to occur if Amine 4 is applied when corn is growing rapidly under high temperature and high soil moisture conditions. In such situations, use the low rate of ½ pint per acre. After application, delay cultivation for 8 to 10 days to allow the corn to overcome any temporary brittleness.
NOTE: Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

With Liquid Nitrogen Solutions: For late season control of young smartweeds, cocklebur, annual morningglory and other annual broadleaf weeds less than 1 inch high. Field should be as clean as possible and corn 20 to 30 inches tall. Apply 1 pint with 80 to 120 lbs. Nitrogen per acre. The spray **MUST** be prepared by first adding required amount of liquid nitrogen solution to spray tank. Next dilute 1 pint of AMINE 4 with 2 quarts of clean

water for each acre to be treated with one tankful. Start the tank agitator and **SLOWLY** add the diluted 2,4-D solution. Spray immediately, maintaining continuous agitation until spray tank is empty. Direct the spray to lower 3" to 4" of corn stalk.

Use spray equipment designated to handle corrosive liquid nitrogen solutions. After spraying, remove any remaining solution and rinse spray rig thoroughly with water. Mix only one tank at a time. Do not spray during or immediately following cold weather.

WEED CONTROL IN SORGHUM (MILO): See Table for recommended use rates. Treat only after the sorghum is 6 inches high and preferably before it is 15 inches high. Do not treat during the boot, tasseling, or early dough stages. Reduce spray drift by keeping the boom and spray nozzles as low as possible. If crop is taller than 8 inches, use drop nozzles to keep the spray off the leaves. Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply AMINE 4 under these conditions, use no more than ½ pint per acre.
Note: Hybrids vary in tolerance in 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

WEED CONTROL IN RICE: See Table for recommended use rates. Apply in the late tillering stage of rice development, at the time of first joint development (first to second green ring), usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed ½ inch, at early seedling, early panicle, boot, flowering, or early heading growth stages. **NOTE:** Some rice varieties under certain conditions can be injured by 2,4-D. Therefore before spraying consult local Extension Service or university specialists for appropriate rates and timing of 2,4-D sprays.

Amount of AMINE 4 to use in crops:

CROP	DOSAGE PER ACRE	
	Normal rates (usually safe for crops)	Higher rate for special situations ² (More likely to injure crop)
SMALL GRAINS		
Spring postemergence wheat, barley, rye	¾ to 1½ pints	2 to 3 pints
oats	¾ to 1 pint	1½ to 2 pints
Preharvest (dough stage)		
wheat, barley, oats	1 to 2 pints	2 to 3 pints
CORN¹		
Preemergence	2 to 4 pints	
Emergence ¹	1 pint	1½ pints
Postemergence ¹		
up to 8 inches tall	¾ to 1 pint	
8 inches to tasseling (use only directed spray)	1 pint	1½ to 2½ pints
SORGHUM (Milo)¹		
Postemergence		
6 to 8 inches tall	2/3 to 1 pint	
8 to 15 inches tall (use only directed spray)	1 pint	1½ to 2 pints
RICE	1½ to 2 pints	2 to 3 pints

¹ Corn and sorghum varieties vary in tolerance to 2,4-D, some are easily injured. Before spraying, get information on 2,4-D tolerance of specific varieties and spray only those known to be resistant to 2,4-D injury. If plants are more than 8 inches tall, use directed spray and keep spray off corn and sorghum foliage.

² These higher rates may be needed to handle difficult weed problems in certain areas such as under dry conditions especially in western areas. However, do not use unless possible crop injury will be acceptable. Consult State Agricultural Experiment Station or Extension Service weed specialists for recommendations or suggestions to fit local conditions.

WEED CONTROL IN SUGARCANE: Use 1 to 2 quarts of product in 5 to 25 gallons of water per acre as a pre- or postemergence spray in the spring after cane emerges and through lay-by. Consult local Agricultural Experiment Station or Extension Service Weed Specialist on specific use of this product. Do not apply within 6 weeks of harvest.

SUGARCANE-HAWAII ONLY
Apply 1 to 3 pts. Amine 4 per acre per application as required, but not to exceed a total of 8 qts. per acre per crop. Do not apply within 6 weeks of harvest.

FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS IN SOYBEANS (Preplant only):

General Information
AMINE 4 is a phenoxy-type herbicide that provides postemergence control of many susceptible annual and perennial broadleaf weeds. AMINE 4 may be applied prior to planting soybeans to provide foliar burn-down control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. AMINE 4 should only be applied preplant to soybeans in situations such as reduced tillage production systems, where emerged weeds are present. Apply only according to the application instructions given below. Do not use any tillage operations between application of AMINE 4 and planting of soybeans.

AMINE 4 2,4-D WEED KILLER

EPA REG. NO. 34704-120

Mixing Instructions

Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of AMINE 4 on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.

Application Procedures

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of spray mixture per acre for ground equipment.

Application Timing and Use Rates

Maximum Rate Per Acre	When to apply (Days prior to planting soybeans)
1 pint	NOT LESS THAN 15 DAYS
2 pints	NOT LESS THAN 30 DAYS

WEEDS CONTROLLED

alfalfa*	garlic, wild*	ragweed, common
bindweed*	horseweed or mare's tail	ragweed, giant
bullnettle	ironweed	sheep's spurge
bittercress, smallflowered	lamb's quarters, common	smartweed, Pennsylvania*
buttercup, smallflowered	lettuce, prickly	sowthistle, annual
Carolina perennium	morningglory, annual	speedwell
onchocitil, common	mouse tail	thistle, Canada*
and rough	mustard, wild	thistle, bull
clover, red*	onion, wild	velvetleaf
cocklebur, common	pennycress, field	vetch, hairy*
caneholon*	peppergrass*	Virginia copperhead
dock, curly*	plantains	
eveningprimrose, cutleaf	purslane, common	

*These species are only partially controlled.

In general, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage at the time of treatment. The response of individual weeds species to AMINE 4 is variable. Consult your local county or state Agricultural Extension Service or crop consultant for advice.

Application Restrictions and Precautions

Important Notice: Unacceptable injury to soybeans planted in fields previously treated with AMINE 4 may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present at the time of application. Do not apply AMINE 4 as described on this label unless you are prepared to accept soybean injury, including loss of stand and yield. Apply a maximum of one application per growing season regardless of the treatment rate.

Do not use on sandy soils with less than 1% organic matter.

Do not replant fields treated with AMINE 4 in the same growing season with crops other than those labeled for use with AMINE 4.

Do not apply AMINE 4 when weather conditions such as temperature air inversions or wind favor drift from treated areas to susceptible plants. **Livestock Grazing Restriction:** Do not feed hay, forage or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

In fields previously treated with AMINE 4, plant soybean seed as deep as practical or at least 1 inch deep. Adjust the planter, if necessary, to ensure that planted seed is completely covered.

STRAWBERRIES

To control broadleaf weeds in established strawberry plantings, apply 2 to 3 pts. AMINE 4 in 25 to 50 gallons of water per acre. Apply in early spring when strawberries are dormant or immediately after the last picking. Do not apply unless possible injury to the crop is acceptable. Follow recommendations of State Extension Weed or Horticultural Specialist in your area.

AQUATIC WEED CONTROL (ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, rivers, and streams that are quiescent or slow moving)

Use 4 to 8 pts. AMINE 4 per acre to control weeds including water hyacinth. Spray the weed mass only. Use 8 pts. per acre when plants are matured or when the weed mass is dense. Spray when weeds are actively growing. Repeat as necessary to kill regrowth.

Surface Application: Use power sprayers operated with a boom or spray gun mounted on a boat, tractor, or truck. Thorough wetting of foliage is essential for maximum control. Use 100 to 400 gals. per acre of spray mixture. Special precautions, such as the use of lower pressure, large nozzles, and thickening agents should be taken to avoid spray drift in areas of sensitive crops.

Air Application: Use drift control spray equipment or thickening agents mixed into the spray solution. Apply 8 pts. AMINE 4 per acre through standard boom systems with a minimum of 5 gallons of spray mixture per acre.

NOTICE TO APPLICATORS

State and Local coordination: Before application, coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for such use.

Fish Toxicity: To avoid fish kill from decaying plant material, do not treat more than one half the lake or pond at one time. For large bodies of weed infested waters leave buffer strips of at least 100 feet wide and delay treatment of these strips for 4 to 5 weeks or until the dead vegetation has decomposed.

Irrigation: Delay the use of treated water for irrigation for three weeks after treatment unless an approved assay shows that the water does not contain more than 0.1 ppm 2,4-D acid. Do not treat irrigation ditches in areas where water will be used to overhead sprinkler irrigate susceptible crops.

Potable Water: Delay the use of treated water for domestic purposes for a period of three weeks or until such time as an approved assay shows that the water contains no more than 0.1 ppm 2,4-D acid.

FALLOWLAND AND CROP STUBBLE

Apply 1 to 4 pts. AMINE 4 per acre on annual broadleaf weeds and up to 6 pts. per acre on established perennial species. Apply to actively growing weeds. See Planting In Treated Areas section. Do not graze dairy animals on treated areas within 7 days after application. Do not graze meat animals on treated areas within 3 days before slaughter. Do not cut treated grass for hay within 30 days after application.

FOREST SITE PREPARATION

To control alder, susceptible broadleaf weeds, and susceptible woody plants before planting forest seedlings, apply 4 to 8 pts. AMINE 4 in 5 to 25 gallons of water, per acre. To provide uniform uptake of product, apply when sufficient foliage exist.

FOREST CONIFER RELEASE

To control alder, susceptible broadleaf weeds, and susceptible woody plants in conifer plantations, apply 2 to 6 pts. AMINE 4 per acre in a minimum of 5 gallons spray mixture per acre. For best results, apply in the spring before budbreak or after budset in late summer to help reduce risk of conifer injury. Certain conifer species are less tolerant to 2,4-D and injury will occur with application. Consult your local university or Agricultural Extension Service specialist for more specific information on rates and timing of applications.

FORESTRY-TREES INJECTION

For controlling species such as alder, aspen, birch, blackgum, cherry, oak, poplar spp., sweetgum, and tulip poplar, make injections or cuts around the tree or stem, using one injection or cut per inch of trunk diameter. For resistant species such as hickory, injection cuts should touch. For best results, injections should be made during the growing season, May 15 to October 15. **For Concentrate Injections or Stump Treatments:** Use 1 to 2 ml. of undiluted AMINE 4 per injection. The injection bit must penetrate the inner bark.

GRASS PASTURES, RANGELAND, AND CONSERVATION RESERVE PROGRAM

Apply 2 to 4 pts. AMINE 4 per acre, when weeds are small and actively growing and prior to bud stage. Do not apply to newly seeded areas until grass is well established. Do not apply to grass in the early boot through milk stage if grass seed production is desired. Use lower rates on annuals or use higher rate on perennials or when weeds are taller.

Bentgrass and legumes may be injured by this treatment. When using on Conservation Reserve Land, follow all applicable state and Federal regulations. Follow the most severe grazing restrictions imposed either by the pesticide label or by USDA Acreage Conservation Reserve Program, whichever is longest. Do not graze dairy cattle in treated areas for 7 days after application. Do not cut forage for hay within 30 days after application. Do not permit meat animals being finished for slaughter to forage treated fields within 3 days of slaughter.

GRASS SEED CROPS: Use 1 to 4 pints per acre in spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to the milk stage. Spray seedling grass only after the five-leaf stage, using ¾ to 1 pint per acre to control small seedling weeds. After the grass is well established, higher rates of up to 4 pints can be used to control hard-to-kill annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth.

Note: Do not use on bentgrass unless grass injury can be tolerated. Do not graze dairy animals on treated areas within 7 days after application. Do not graze meat animals on treated areas within 3 days before slaughter. Do not cut treated grass for hay within 30 days after application.

NON-CROPLAND (FENCEROWS, HEDGEROWS, ROADSIDES, DRAINAGE DITCHES, ROADSIDES ADJACENT TO ORCHARDS, RIGHTS-OF-WAYS, UTILITY POWER LINES, RAILROADS, AND OTHER NON-CROP AREAS)

Treat annual broadleaf weeds, when young and actively growing, with 2 to 4 pts. AMINE 4 per acre. Apply 4 to 8 pts. AMINE 4 per acre for control of biennial and perennial broadleaf weeds. Do not apply to newly seeded area until grass is well established. Bentgrass, clover, legumes and dichondria may be injured by this treatment. Do not graze dairy animals for 7 days following application. Use sufficient gallonage for thorough and uniform coverage.

ORNAMENTAL AND RECREATIONAL TURF

For weed control on golf courses, cemeteries, parks, and lawns, apply 2 to 4 pts. AMINE 4 per acre when weeds are young and actively growing. Do

AMINE 4 2,4-D WEED KILLER
EPA REG. NO. 34704-120

GIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

7 of 7

not apply to newly seeded areas until grass is well established. Use sufficient gallonage for thorough and uniform coverage. Do not apply more than 2 broadcast applications per year per treatment site.

PINE RELEASE: To control hardwoods, such as Oak, Hickory, Maple, Pecan, Elm, Sumac, and Hawthorn in Southern pine stands, use AMINE 4 undiluted in a concentrate tree injector calibrated to apply 0.75 ml. per injection. Space injections 2' apart, edge to edge, completely around the tree and close to the base. The injector bit must penetrate the inner bark. On hard-to-kill species such as Hickory, Dogwood, Red Maple, Blue Beech and Ash, make injections 1' to 1½' apart, edge to edge. Treatment may be made at any time of year.

CONTROL OF SOUTHERN WILD ROSE: On rangelands, roadsides, and fence-rows, use 1 gallon of AMINE 4 plus 4 to 8 fluid ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required. On rangeland, apply a maximum of 2 quarts of AMINE 4 per acre per application. Do not graze dairy animals on treated areas within 7 days after application. Do graze meat animals on treated areas within 3 days before slaughter. Do not cut treated grass for hay within 30 days after application.

WEEDS AND BRUSH ON IRRIGATION CANAL DITCHBANKS—
Seventeen Western States
Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska,
New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South
Dakota, Texas, Utah, Washington and Wyoming.

For control of annual and perennial broadleaf weeds, apply 1 to 2 quarts of AMINE 4 per acre in approximately 20 to 100 gallons of water per acre. Treat when weeds are young and actively growing before the bud or early bloom stage. For harder-to-control weeds a repeat spray after 3 to 4 weeks using the same rates may be needed for maximum results. Apply no more than two treatments per season.

For woody brush and patches of perennial broadleaf weeds, mix one gallon of AMINE 4 in 150 gallons of water. Wet foliage thoroughly using about one gallon of solution per square foot.

Spraying instructions: Low pressure (10 to 40 psi) power spray equipment should be used and mounted on a truck, tractor or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is calm, 5 mph or less. Do not use on small canals (less than 10 fts) where water will be used for drinking purposes. Boom spraying onto water surface must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than two-foot overspray onto water with an average of less than one-foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

Water within treated banks should not be fished.
 Do not graze dairy animals on treated areas within 7 days after application.
 Do not graze meat animals on treated areas within 3 days before slaughter.
 Do not cut treated grass for hay within 30 days after application.

SPOT TREATMENT IN NON-CROP AREAS: To control broadleaf weeds in small areas with hand sprayer, use ¼ pint of AMINE 4 in 3 gallons of water and spray to thoroughly wet all foliage.

SMALL QUANTITY DILUTION TABLE

To spray small areas use the following dilution table.

If Dosage on Label shows:	Use this Amount for each Gallon of water
2 pints (1 quart)	¾ ounces (4 teaspoons)/1,000 sq. ft.
3 pints (1½ quarts)	1¼ ounces (2½ tablespoons)/1,000 sq. ft.
4 pints (2 quarts)	1½ ounces (3 tablespoons)/1,000 sq. ft.
6 pints (3 quarts)	2¼ ounces (4½ tablespoons)/1,000 sq. ft.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELI-

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697