

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

14

ASHINGTON, D.C. 2046

MAY 3 1 1994

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

William M. Mahlburg Agent for: PLATTE CHEMICAL CO., INC. P. O. Box 667 Greeley, CO 80632

Subject: Label Amendment Submission of 04/13/94 in Response to PR Notice 93-7 EPA Reg. No. 34704-130 CLEAN CROP MCP AMINE 4

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

# WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

# WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.



Page 2

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL) Office of Pesticide Programs (7505C) U.S. Environmental Protection Agency 401 M Street, SW Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL) Office of Pesticide Programs Room 266A, Crystal Mall 2 1921 Jefferson Davis Highway Arlington, VA 22202

Sincerely, V Jim Tompkin, Deputy Chief **Registration Support Branch** Registration Division (7505W) ľ

Attachment

.

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division


1

William M. Mahlburg WILLIAM M. MAHLBURG AGENT FOR: PLATTE CHEMICAL CO., INC. - BOX 667 GREELEY CO 80632

Comment for: EPA Reg Nr.34704-130 CLEAN CROP MCP AMINE 4

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

Remove the statement "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application." from its current position within the Agricultural Use Box and place it above the Agricultural Use Box.



## **MCP AMINE 4**

EPA REG. NO. 34704-130

## STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function property.

STORAGE: Do not store below temperature of 25°F. If frozen, warm to 70°F, and redissolve before using by rolling or shaking the container. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to tabel instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or pulicture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### **GENERAL INFORMATION**

This product is an amine salt formulation of MCP (2-Methyl-4chlorophenoxyacetic acid), a selective herbicide similar to 2,4-D. Certain crops, such as flax, oats, and rice appear to be more tolerant of MCP than 2,4-D, and some weeds such as Mustard and Canada thistle are more easily controlled by MCP than 2,4-D.

Generally, the lower dosages listed will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Use moderate spray pressure (20 pounds) and enough spray volume for good coverage, usually 5 to 20 gallons per acre by ground equipment and at least 2 to 5 gallons per acre by air. Higher spray volumes may be useful in some situations to provide better spray coverage of weed foliage and to reduce spray drift. The use of adjuvants such as wetting agents, may increase herbicidal effectiveness but may also decrease selectivity.

### Do not apply this product through any type of irrigation system.

Agitation is helpful to assure uniform mixing, but is not necessary after the product has been completely dispersed. Use of this product may be governed in certain areas by local regulations. MCP produces best results when applied to weeds growing rapidly. Thorough coverage is important. Lack of moisture may cause less effective control. Some perennial weeds may require repeat treatment for control. Treat such weeds alloud stage if this occurs during a period when crop is not unduly susceptible.

ATTENTION—Extreme care must be used when applying MCP to prevent injuring desirable plants and crops MCP should not be applied in the vicinity of susceptible crops such as cotton, tomatoes, vegelables, grapes, fruit trees, soybeans, tobacco and beans. Do not allow spray mist to drift on to such plants, since even minute quantities of spray may cause severe injury. Use low pressures when spraying to produce coarse sprays which are less likely to drift. Do not spray on windy days. Do not use in greenhouses. Large amounts of MCP in the soil may temporarily inhibit seed germination. Never use more MCP than recommended, Do not use in any manner other than recommended on the tabel to avoid possible crop injury or residues at harvest.

It is best to use a separate sprayer for application of MCP herbicides. If the same sprayer must be used for other crop spraying, it should be flushed thoroughly with water immediately after use. Next, fill equipment with water containing one volume household ammonia to 100 volumes of water, and atlow to stand for one day. Then flush equipment thoroughly, including hoses and nozzles, several times with water.

Read and carefully observe the cautionary statements and all other information appearing on labeling of all products used in mixtures. Do not contaminate viater used for irrigation, domestic or spray purposes. Do not contaminate seed, feed, foodstuffs, fertilizers, insecticides or fungicides.

Do not apply where spray drift may be a problem due to proximity of susceptible crops or other desirable plants. Read and follow all Use Precautions given on this tabel.

PREPARING THE SPRAY: Mix CLEAN CROP MCP AMINE 4 with water, unless otherwise directed on this label. Fill the mixing tank or spray tank about half full with water, then add the required amount of this product. with agitation. NOTE: Adding oil, wetting agent or other surfactant to the spray may reduce selectivity to crops, possibly resulting in crop injury.

#### Some Weeds Controlled By MCP

Galinsoga

Goatsbeard

Hemonettle

Jimsonweed

Lambsquarter

Pepperweed

**Prickly lettuce** 

Pennycress

Ragweed

Red Stern

Marsh eider

Mustards

Pigweed

Plantain

Henbit

Indigo

Barberry Beggartick Buckhom Bulrushes Burcucumber Burdock Buttercup Canada thistle Carpetweed Catsear Chickweed Cocklebur Croton Daisy Dandelion Dock

Sedge Shepherdspurse Sicklepod Stinkweed Sunflower Tie vine Water plantam White brush White brush White lop or Hoary Cress Wild carrot Wild radish Yellow rocket

WEED CONTROL IN SMALL GRAINS: NOTE: Do not forage or graze > meat or deiry animals on treated areas within 7 days of slaughter.

SELECTIVE SPRAYING: Wheat, Oats, Barley and Rye-Apply as a water mix spray by ground sprayer or airplane. Use 1/2 to 1 pint per acre for the more susceptible weeds after crop has reached the 3 to 4 leaf stage up to boot stage. Use up to 3 pints per acre for the less susceptible weeds after crop has tillered and up to early boot stage. Do not spray from boot to dough stage. For small grains underseeded with legumes, see the following paragraph.

SMALL GRAINS: Underseeded with Alfalta, Birdsfoot, Tretoit, Lespedeza, Red and White Clover: For emergency control of serious infestations of mustards, yellow rocket and other susceptible broadleaf weeds, apply 1/4 to 1/2 pint in not more than 5 to 10 gallons of watter per acre. Make application after cereal is well tillered (4 to 8 inches tail) and before reaching boot stage. The nurse crop and weeds should provide a protecting canopy which, together with the use of low gallonage applied at low pressure, will reduce the risk of damage to the legumes. To control certain weed species, 1 pint per acre may be needed. Do not use where other small seeded legumes (vetch, sweet clover, trefoil, aifalfa) are seeded unless injury to them can be tolerated.

FLAX: Use 1/2 pint CLEAN CROP MCP AMINE 4. Apply by ground sprayer or by airplane. Use sufficient water to give uniform and adequate coverage. Apply only when weeds are up and when flax is 4 to 8 inches high and before it comes into bud. Treatment atter early bud stage may result in severe damage. If Canada thistle is present it may be necessary to go as high as 3/4 to 1 pint per acre to prevent seed head production. Some injury to the flax may result. Do not lorage or graze meat animals on treated areas within 7 days of slaughter.

WEED CONTROL IN RICE: For control of certain broadleaf weeds and sedges, apply 2 to 3 pints of CLEAN CROP MCP AMINE 4 in 10 to 30 gallons of water per acre. Treat when rice is fully tillered and 6 to 8 inches above water, usually 35 to 65 days after planting. Do not treat when rice is in the early seedling, boot or early heading stages, or when temperatures are over 90°F. Some rice varieties may be less tolerant to MCP than others. Consult your Agricultural Experiment Station or Extension Service Weed Specialists for appropriate rates and timing of MCP spray on the tolerance of rice varieties. Do not grow crayfish or catfish in treated rice fields.

WEED CONTROL IN PEAS: Use on peas to control susceptible weeds only where recommended by local agricultural Extension Service or Experiment Station specialists. Use 1/4 to 1/5 pint of CLEAN CROP MCP AMINE 4 in at least 15 gations of water per acre when peas are 3 to 7 inches tail, before first floworing and weeds are small. Higher rates of 1/2 to 3/4 pimt (4-6 ounces MCP) per acre may be used to improve control of difficult eds but crop injury is more likely to occur. Do not apply if peas are tailer than 7 inches or when they are stressed for lack of soil moisture. Do not apply when temperatures are over 90°F. NOTE: MCP can cause injury and defayed inaturity in the pea crop.

GRASSES GROWN FOR SEED: Use 1 to 2 pints per acre in sufficient water to give adequate coverage. Use the higher rate where weed stands are heavy. In established grasses apply in spring before head comes into boot and on seedling grass after grass has tillered.

NOTE: In weed control in grasses, repeat treatment may be needed for less susceptible weeds. While clover and other legumes may be temporarily injured or killed. In some areas, bent, butfalo, carpet and St. Augustwe grasses may also be injured by the treatment.

WEED CONTROL IN ESTABLISHED ALFALFA: For control of yellow rocket and other susceptible annual weeds such as pennycress or fanweed, use 1 pini per acre of CLEAN CROP MCP AMINE 4. Apply in late ( fall following frosts when the legumes are dormant. The temperature at the time of spraying should be above 40°F.



## **MCP AMINE 4**

### EPA REG. NO. 34704-130

### WEED CONTROL IN GRASS PASTURES &

GRASSLAND: Use 2 to 4 pints per acre of CLEAN CROP MCP AMINE 4. Apply when weeds are small and actively growing for best results. Use the higher rates for whitetop, Canada thistle, buttercup and other more difficult to control weeds. Legumes may be injured or killed. Do not use this treatment where legumes, especially alfalfa, are present and desirable. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage where grass seed production is desired. Do not graze dairy animals on treated grass until 7 days after treatment. Do not graze meat animals in treated areas within 7 days of slaughter. Do not harvest grass for hay less than 21 days after application.

CONTROL OF WHITEBRUSH ON RANGELAND: Use 3 pints of CLEAN CROP MCP AMINE 4 with 1 gallon of diesel oil and enough water for good spray coverage, usually 6 to 10 gallons per acre by aircraft. Apply in spring or fall when plant foliage is well developed and actively growing. Spraying during blown is recommended but not immediately after Shedding of blossoms. Retreatments may be needed in succeeding years. NOTE: First mix this product in the water, then add the oil with agitation. Spray should be kept agitated during spraying to avoid separation in the tank.

LAWNS, GOLF COURSES AND SIMILAR TURF: For controlling whitetop, dandelions, and similar weeds apply 1 to 2 pints per acre with sufficient water to provide good coverage. Treatment may reduce stands of clover which may be present. Do not apply to newly seeded lawns until grass has become wett established. MCP may injure hent or other creeping grasses. For best result treat thistles when in the early bud stage.

NON-CROP SPRAYING: CANADA THISTLE, WHITE TOP AND MEADOW BUTTERCUP (In Non-Crop Areas such as Roadsides, Fence Rows, Rights-of-Way and Similar Places): For spot treatment, use 1/4 pint to 3 to 4 gallons of water, or 3/4 gallon per acre in sufficient water to give coverage for more extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on fall regrowth. Spray European barberry during any season. Thoroughly wet the foliage, base of the canes and the ground line. Do not forage or graze livestock on treated areas within 7 days of treatment. Local conditions may affect use of this chemical. Consult State agricultural extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

### NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFO' IMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REA-SONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PROD-UCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BE-YOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FIT-NESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

BEST AVAILABLE COPY

FORMULATED FOR PLATTE CHEMICAL CO. 150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697