



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460

Reg. Number:	Date of Issua

34704-130

6 JUN 2008

NOTICE OF PESTICIDE:

_ Registration

<u>x</u> Reregistration (under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:
MCP Amine 4

Name and Address of Registrant (include ZIP Code):

Loveland Products, Inc.

P. O. Box 1286

Greeley, CO 80632-1286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

A Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

- B. Make the following label changes:
 - Under Precautionary Statements";

Add "Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals." Also, delete "Avoid breathing vapor . . . domestic or spray purposes."

"First Aid":

Add "Note to Physicians - Probable mucosal damage may contraindicate the use of gastric lavage."

3. Change the "Personal Protective Equipment (PPE)" to the following:

vanne J. Miller

Mixers, loaders, applicators, flaggers and other handlers must wear:

- Long- sleeved shirt and long pants,
- · Protective eyewear (goggles or face shield),
- · Shoes plus socks, and

Signature of Approving Official:

Date:

6 JUN 2001

Joanne I. Miller ()
Product Manager (23)
Herbicide Branch

Registration Division (7505P)

Page 2 EPA Reg. No. 34704-130

- Chemical- resistant gloves made of any waterproof materials, meet the requirements of EPA selection Category A, when mixing, loading, or using any hand-held equipment.
- Use the following respirator requirements:
 - The mixers and loaders supporting aerial application to rangelands, pasture lands, or non-crop land must wear a National Institute for Occupational Safety and Health (NIOSH) approved half-mask respirator equipped with any N, R, P or HE series particulate air filters and has the approval number prefix (84A-XXX).
 - It is recommended that you require that respirator wearers be fit tested, and trained in the proper use, care, cleaning, inspection, and storage of the respirator. Refer to the Occupational Safety and Health Administration (OSHA) standard on respiratory protection (29 CFR 1910.134) for additional information.
- 4. Under "User Safety Recommendations";

Change "Remove clothing immediately" to "Remove clothing/PPE immediately."

- 5. Under "Storage and Disposal"
 - · Change "Storage" to "Pesticide Storage."
 - Move "Do not reuse containers . . . to function properly" to the Container Disposal subsection.
 - Delete "Open dumping is prohibited."
- 6. Under "Engineering Control Statement";

Add "Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d) (6)]."

Under "Non-Agricultural Use Requirements";

Change "Do not enter treated area until spray have dried." to "Do not enter or allow others to enter treated area until spray have dried."

8. Under "Control of Whitebrush on Rangeland";

Add "Do not apply more than 3. pints of product (1.5lb. ae) per acre per year" to the rangeland application instruction.

9. Under "General Information"; (page 3)

Remove "such as," "similar places" and "similar turf."

10. Under "Environmental Hazards":

Remove "This pesticide may be toxic to fish, aquatic invertebrates, and aquatic plants."

11. Add the following statement to the front panel of the label if this product is intended for Residential Use:

"Intended for use by applicators who are authorized/licensed by the state for this type of application."

Page 3 EPA Reg. No. 34704-130

12. Under "Warranty Disclaimer";

Add "To the extent consistent with applicable law" before "Unintended consequences, including but not limited to . . . assume by buyer or user."

13. Under "Limitation of Liability";

Add "To the extent consistent with applicable law" before "Manufacturer or seller shall not be liable to buyer . . . sold by the manufacturer or seller to buyer" and "Buyer's or user's exclusive remedy, . . . or its customers."

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure



FOR BROADLEAF WEED CONTROL IN SMALL GRAINS, PEAS, ALFALFA, FLAX, GRASSLAND AND NON-CROP AREAS.

ACTIVE INGREDIENT: 2-Methyl-4-chlorophenoxyacetic acid, dimethylamine salt 52.0%* INERT INGREDIENTS: TOTAL 100.0%

*Equivalent to 42.5% 2-Methyl-4-chlorophenoxyacetic acid or 4.0 pounds per gallon.

ONLY FOR SALE TO, USE AND STORAGE BY AGRICULTURAL PERSONNEL OR COMMERCIAL APPLICATORS.

KEEP OUT OF REACH OF CHILDREN DANGER—PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes,
j	then continue rinsing eye:
	Call a poison control center or doctor for treatment advice.
lf on skin	Take off contaminated clothing.
or clothing:	Rinse skin immediately with plenty of water for 15-20 minutes.
	 Call a poison control center or doctor for treatment advice.
If swallowed:	Call a poison control center or doctor immediately for treatment advice.
}	Have a person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by the poison control center or doctor.
	 Do not give anything by mouth to an unconscious person.
If Inhaled:	Move person to fresh air.
	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-301-7976.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

(See Below For Additional Precautionary Statements)

EPA REG. NO. 34704-130

EPA EST. NO. 37507-MT-1

NET CONTENTS _____ GALS. (

EXP 10B06

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

PELIGRO

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin. Do not contaminate water used for irrigation, domestic or spray purposes.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category "F" on an EPA chemical-resistant category selection chart.

Mixers, loaders, applicators, flaggers and other handlers must wear: Longsleeved shirt and long pants, shoes plus socks, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or Vinton gloves when mixing, loading, or using any hand-held equipment, and protective evewear.

Additional PPE requirements for mixers and loaders supporting aerial applications to rangelands, pasture lands, or noncropland. These mixers/loaders also must wear: A chemical resistant apron, and a NIOSH-approved respirator with a dust/mist filter with a MSHA/NIOSH approved number prefix TC-21C or any N2, R, P,

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

See engineering controls for additional requirements.

Engineering controls:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment wash waters or rinsate. Drift or run-off may adversely affect nontarget plants.

Do not contaminate water when disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Groundwater Contamination:

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

MCP AMI 4 EPA REG. NO. 34704-130

Cleaning of Equipment: When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Extreme care must be used when applying MCP to prevent injuring desirable plants and crops. MCP should not be used in susceptible crops such as cotton, tomatoes, vegetables, grapes, fruit trees, soybeans, tobacco, or beans. Do not allow spray mist to drift onto such plants, since even minute quantities of spray may cause severe injury. Use low pressures when spraying to produce coarse sprays which are less likely to drift. Avoid use of small diameter spray nozzles. Do not use in greenhouses.

Large amounts of MCP in the soil may temporarily inhibit seed germination. Do not use in any manner other than recommended on the label to avoid possible crop injury or residues at harvest.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Keep people and pets out of the area during application. Do not allow people or pets to enter the area until sprays have dried.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application.

Do not enter treatment areas until sprays have dried.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Do not store below temperature of 25° F. If frozen, warm to 70° F. and redissolve before using by rolling or shaking the container. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This product is an amine salt formulation of MCP (2-Methyl-4-chlorophenoxyacetic acid), a selective herbicide similar to 2,4-D. Certain crops, such as flax, oats, and rice appear to be more tolerant of MCP than 2,4-D, and some weeds such as Mustard and Canada thistle are more easily controlled by MCP than 2,4-D.

Generally, the lower dosages listed will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Use moderate spray pressure (20 pounds) and enough spray volume for good coverage, usually 5 to 20

gallons per acre by groun ament and at least 2 to 5 gallons per acre by air, Higher spray volumes may seful in some situations to provide better spray coverage of weed foliage and to reduce spray drift. The use of adjuvants such as wetting agents, may increase herbicidal effectiveness but may also decrease selectivity.

Do not apply this product through any type of irrigation system.

Agitation is helpful to assure uniform mixing, but is not necessary after the product has been completely dispersed. Use of this product may be governed in certain areas by local regulations. MCP produces best results when applied to weeds growing rapidly. Thorough coverage is important. Lack of moisture may cause less effective control. Some perennial weeds may require repeat treatment for control. Treat such weeds at bud stage if this occurs during a period when crop is not unduly susceptible.

ATTENTION—Extreme care must be used when applying MCP to prevent injuring desirable plants and crops. MCP should not be applied in the vicinity of susceptible crops such as cotton, tomatoes, vegetables, grapes, fruit trees, soybeans, tobacco and beans. Do not allow spray mist to drift on to such plants, since even minute quantities of spray may cause severe injury. Use low pressures when spraying to produce coarse sprays which are less likely to drift. Do not spray on windy days. Do not use in greenhouses. Large amounts of MCP in the soil may temporarily inhibit seed germination. Never use more MCP than recommended. Do not use in any manner other than recommended on the label to avoid possible crop injury or residues at harvest.

It is best to use a separate sprayer for application of MCP herbicides. If the same sprayer must be used for other crop spraying, it should be flushed thoroughly with water immediately after use. Next, fill equipment with water containing one volume of household ammonia to 100 volumes of water, and allow to stand for one day. Then flush equipment thoroughly, including hoses and nozzles, several times with water.

Read and carefully observe the cautionary statements and all other information appearing on labeling of all products used in mixtures. Do not contaminate water used for irrigation, domestic or spray purposes. Do not contaminate seed, feed, foodstuffs, fertilizers, insecticides or fungicides.

Do not apply where spray drift may be a problem due to proximity of susceptible crops or other desirable plants. Read and follow all Use Precautions given on this label.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site. For groundboom applications; do not apply with a nozzle height greater than 4 feet above the crop canopy.

Aerial Applications: The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater that 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

PREPARING THE SPRAY: Mix MCP AMINE 4 with water, unless otherwise directed on this label. Fill the mixing tank or spray tank about half full with water, then add the required amount of this product with agitation. NOTE: Adding oil, wetting agent or other surfactant to the spray may reduce selectivity to crops, possibly resulting in crop injury.

Some Weeds Controlled By MCP

	Some weeds Controlled by	IVICE
Barberry	Galinsoga	Sedge
Beggartick	Goatsbeard	Shepherdspurs
Buckhorn	Hempnettle	Sicklepod
Bulrushes	Henbit	Stinkweed
Burcucumber	Indigo	Sunflower
Burdock	Jimsonweed	Tie vine
Buttercup	Lambsquarter	Water plantain
Canada thistle	Marsh elder	Mustards
Carpetweed	Pepperweed	White brush
Catsear	Pigweed	White top or
Chickweed	Plantain	Hoary Cress
Cocklebur,	Prickly lettuce	Wild carrot
Croton	Pennycress	Wild radish
Daisy	Ragweed	Yellow rocket
Dandelion	Red Stem	

WEED CONTROL IN SMALL GRAINS: NOTE: Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment.

Dock

MCP A NE 4 EPA REG. NO. 34704-130

SELECTIVE SPRAYING: Wheat, Oats, Barley and Rye—Apply as a water mix spray by ground sprayer or airplane. Use ½ to 1 pint per acre for the more susceptible weeds after crop has reached the 3 to 4 leaf stage up to boot stage. Use up to 1½ pints per acre for the less susceptible weeds after crop has tillered and up to early boot stage. Do not spray from boot to dough stage. For small grains underseeded with legumes, see the following paragraph. Do not apply more than 0.75 lbs ae/acre per year (1.5 pints of product/acre/year).

SMALL GRAINS: Underseeded with Alfalfa, Birdsfoot, Trefoil, Lespedeza, Red and White Clover: For emergency control of serious infestations of mustards, yellow rocket and other susceptible broadleaf weeds, apply ¼ to ½ pint and no more than 5 to 10 gallons of water per acre. Make application after cereal is well tillered (4 to 8 inches tall) and before reaching boot stage. The nurse crop and weeds should provide a protecting canopy, which together with the use of low gallonage applied at low pressure, will reduce the risk of damage to the legumes. To control certain weed species, 1 pint per acre may be needed. Do not use where other small seeded legumes (vetch, sweet clover, trefoil, alfalfa) are seeded unless injury to them can be tolerated.

FLAX: Use ¼ to ½ pint MCP AMINE 4. Apply by ground sprayer or by airplane. Use sufficient water to give uniform and adequate coverage. Apply only when weeds are up and when flax is 4 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not apply more than 0.25 lbs ae per acre per year (0.5 pints of product/acre/year).

WEED CONTROL IN PEAS IN THE PACIFIC NORTHWEST: For use only in the Pacific Northwest. Use on peas to control susceptible weeds only where recommended by local agricultural Extension Service or Experiment Station specialists. Use ½ to ½ pint of MCP AMINE 4 in at least 15 gallons of water per acre when peas are 3 to 7 inches tall, before first flowering and weeds are small. Higher rates of ½ to ¾ pint (4-6 ounces MCP) per acre may be used to improve control of difficult weeds but crop injury is more likely to occur. Do not apply if peas are taller than 7 inches or when they are stressed for lack of soil moisture. Do not apply when temperatures are over 90° F. NOTE: MCP can cause injury and delayed maturity in the pea crop.

GRASSES GROWN FOR SEED: Use 1 to 1½ pints per acre in sufficient water to give adequate coverage. Use the higher rate where weed stands are heavy. In established grasses apply in spring before head comes into boot and on seedling grass after grass has tillered. Do not apply more than 1.5 lb ae per acre per year (3 pints of product per acre per year). Do not make more than 2 applications per year with a minimum retreatment interval of 21 days.

NOTE: In weed control in grasses, repeat treatment may be needed for less susceptible weeds. White clover and other legumes may be temporarily injured or killed. In some areas, bent, buffalo, carpet and St. Augustine grasses may also be injured by the treatment.

WEED CONTROL IN ESTABLISHED ALFALFA: For control of yellow rocket and other susceptible annual weeds such as pennycress or fanweed, use 1 pint per acre of MCP AMINE 4. Apply in late fall following frosts when the legumes are dormant. The temperature at the time of spraying should be above 40° F.

WEED CONTROL IN GRASS PASTURES & GRASSLAND: Use 2 to 3 pints per acre of MCP AMINE 4. Apply when weeds are small and actively growing for best results. Use the higher rates for whitetop, Canada thistle, buttercup and other more difficult to control weeds. Legumes may be injured or killed. Do not use this treatment where legumes, especially alfalfa, are present and desirable. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage where grass seed production is desired. Do not graze dairy animals on treated grass until 7 days after treatment. Do not graze meat animals in treated areas within 7 days of slaughter. Do not harvest grass for hay less than 21 days after application. Do not apply more than 1.5 lb ae per acre per year (3 pints of product per acre per year). Do not make more than 2 applications per year with a minimum retreatment interval of 21 days.

CONTROL OF WHITEBRUSH ON RANGELAND: Use 3 pints of MCP AMINE 4 with 1 gallon of diesel oil and enough water for good spray cover

age, usually 6 to 10 gallons per acre by aircraft. Apply in spring or fall when plant foliage is well developed and actively growing. Spraying during bloom is recommended but not immediately after shedding of blossoms. Retreatments may be needed in succeeding years. NOTE: First mix this product in the water, then add the oil with agitation. Spray should be kept agitated during spraying to avoid separation in the tank.

LAWNS, GOLF COURSES, SOD FARMS AND SIMILAR TURF: For controlling whitetop, dandelions, and similar weeds apply 1 to 1½ pints per acre with sufficient water to provide good coverage. Treatment may reduce stands of clover which may be present. Do not apply to newly seeded lawns until grass has become well established. MCP may injure bent or other creeping grasses. For best result treat thistles when in the early bud stage. For residential turf and other uses under this heading: Do not apply more than 1.5 lb ae per acre per year (3 pints of product per acre per year). Do not make more than 2 applications per year with a minimum retreatment interval of 21 days.

NON-CROP SPRAYING: CANADA THISTLE, WHITE TOP AND MEADOW BUT-TERCUP (In Non-Crop Areas such as Roadsides, Fence Rows, Rights-of-Way and Similar Places): For reatment, use ¼ pint to 3 to 4 gallons of water, or ¾ gallon per acre in sufficie... water to give coverage for more extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on fail regrowth. Spray European barberry during any season. Thoroughly wet the foliage, base of the canes and the ground line. Do not forage or graze livestock on treated areas within 7 days of treatment. Local conditions may affect use of this chemical. Consult State agricultural extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages. Do not apply more than 3.0 lb ae per acre per year (3 quarts product per acre per year) for soot treatment.

Broadcast treatment: Do not apply more than 1.5 lb ae per acre per year (3 pints of product per acre per year). Do not make more than 2 applications per year with a minimum retreatment interval of 21 days.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVE-LAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, ATTHE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80632-1286