

34704-131

1/2



MCPP GREEN

**For Ornamental Turfgrass Weed Control.
Only for Sale To, Use and Storage by
Lawncare and Landscaping Personnel or
Commercial Applicators.**

ACTIVE INGREDIENT:

Potassium salt of 2-(2-methyl-4-chlorophenoxy) propionic acid*	48.3%
INERT INGREDIENTS:	51.7%
TOTAL	100.0%

*Equivalent to 41.2% of 2-(2-methyl-4-chlorophenoxy) propionic acid or 4 lbs. of MCPP acid per U.S. Gallon at 68° F.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements
EPA REG. NO. 34704-131
EPA EST. NO. 37507-MT-1
NET CONTENTS 2½ GALS. (9.46 L)
IHT EXP 08P03

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed or absorbed through the skin. If swallowed, induce vomiting immediately by giving two glasses of water and stick finger down throat. Never give anything by mouth to an unconscious person. Seek medical attention.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15–20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-301-7976.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Drift or run-off may adversely affect nontarget plants. For terrestrial uses,

do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Groundwater Contamination:
Most cases of groundwater contamination involving phenoxy herbicides such as MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Cleaning of Equipment:
When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Use care to avoid spray contact or drift to susceptible plants such as beans and other legumes, cotton, flowers, grapes, ornamentals, vegetables and other plants. Do not permit spray mist containing this product to drift onto them, since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods. Do not spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays to minimize drift. Spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 8 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine droplet spray. Flush sprayer out on suitable non-crop area after use. Do not use the spray equipment for applying other materials to susceptible crops as injury may result.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not reenter or permit workers to reenter treated areas until sprays have dried.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly.
STORAGE: Do not store below temperatures of 32°F. When stored at temperatures below freezing, it may be necessary to warm contents to 45°F and mix thoroughly before using. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Read entire label before using this product.

GENERAL INFORMATION

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. This herbicide may be used on ornamental turf grasses with greater safety than 2,4-D. This herbicide is safe to use on Washington, Toronto and Seaside bentgrasses, Kentucky and Meroe bluegrasses. Apply on a warm calm day when weeds are growing vigorously. Use any suitable sprayer or

MCPP GREEN

EPA REG. NO. 34704-131

watering can with fine nose to give a uniform cover of the weeds. Do not apply this product through any type of irrigation system. Although this herbicide is relatively slow in action it gives a superior result on the more difficult to control weeds such as:

More Susceptible Weeds
Controlled by 2½ pints per acre.
Clover
Common chickweed
Lambquarters
Mouse-ear chickweed

Pigweed
Plantain
Ragweed

Less Susceptible Weeds
Controlled by 4 pints per acre. Repeat applications at half the recommended dosage may give better control.
Black medic
Knotweed
Morningglory

Purslane
Speedwell
Yarrow

Note: Local conditions, crop varieties and application regulations vary and may affect use of this herbicide. Consult local agricultural experiment station or extension service weed specialists and state regulatory agencies for recommendations in your area.

Preparation of the Spray: Fill half the spray tank with water; then add the required amount of this product. Mix thoroughly, adding the remainder of water during the operation.

Rate of Use	Per 1,000 sq. ft.	Per acre
Established greens	1 fluid oz. in 2½ gallons of water	2½ pints in 20-40 gallons of water
Established lawns, golf fairways	1½ fluid oz. in 2½ gallons of water	4 pints

Time of Application: To obtain the best results this herbicide must be applied when the weeds are most susceptible and conditions are such that no damage to the grasses will occur. Apply when the weeds are growing vigorously. During periods of drought weeds become more resistant to herbicides. Under the same conditions fine grasses are more susceptible to this herbicide particularly when temperatures exceed 90F. This chemical is absorbed by the foliage of broadleaf weeds and is translocated throughout the whole plant to roots, etc. where it eventually causes death of the plant. Do not cut or mow the lawn 2 to 3 days before or after treatment so that maximum uptake and translocation can occur. Do not spray if rain is likely otherwise the chemical may be washed off before the absorption has taken place. For the same reason do not water the lawn for at least 24 hours after application.

Better control of the weeds classified as Less Susceptible may be obtained with repeat applications of half the recommended dose carried out at not less than monthly intervals.

This product is not recommended for St. Augustinegrass and Centipedegrass, as it can cause damage to these species. Do not use this product on newly sown lawns or turf for six months after planting. Avoid drift onto other plants and do not use the first cuttings after treatment as a mulch for flowers and vegetables.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.
P.O. BOX 667, GREELEY, COLORADO 80632-0667

ACCEPTED
OCT - 2 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
registered under
EPA Reg. No. 34704-131