May cause allergic skin reactions. Do not get in eyes, Viear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately faunder clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

If swellowed—Call a physician or Poson Control Center immediately. Drink one or two glasses of water and induce wornting by touching the back of throat with finger. Repeat until vomit fluid is clear Do not Induce vomiting or give anything by mouth to an unconscirus person.

If Inhaled—Remove victim to fresh air. Apply artificial respiration if indicated.

If on aldn-Remove contaminated clothing and wash affected area with soap and water.

If In eyes—immediately flush eyes with plenty of water for at least 15 minutes. Call a physician.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition. Alropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. On tot apply directly to water or wellands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. On not contaminate water by clearung of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension, Service.

ENDANGERED SPECIES RESTRICTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of their habitats have been identified by county:

Slackweier Darter

Alabama (Lauderdale, Limestone and Madison) Tennessee (Lawrence and Wayne) New Mexico (Chaves and

Pecos gambusia

Eddy) Texas (Reeves, Jelf Davis and Pecos)

Freshwater Mussels

Tennessee (Hancock, Claiborne, Hawkins and Sullayan)

Virginia (Smyth, Scott, Lee, Washington and Grayson

Commanche Springs Puplish Texas (Reeves and Jeff Davis)

t Texas (Reeves and Jeff Davis)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be truated with captan. Consult your regional U.S. Fish and Widdlife Service Mide (Endangered Species, Specialist) or State Wildlife Agency if there is any Count us to the areas where the outdoor one of captan is prohibited.

DIRECTIONS FOR USE

It is a violation of Federal Law to rise this 'product in a manner inconsistent with its labe'r 2.

CONTAINER DISPOSAL: Oo not reuse empty container Wrap container and put in trash.

Wear natural or synthetic rubber, viryl, or plastic gloves impermeable to captan when using. When using wear long pants and long-slewed shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn white handling this product must be laundered separately from other clothing before reuse.

For disease control, begin applications before diseases appear and repeat at 7-10 day intervals. Use a dependable pressure sprayer or hand duster, applying to top and bottom leaf surfaces and all narts of fruit.

APPLES: Apply 8 oz. as a dust or mix 8 oz. in 4 gals. water per tree for the control of Scab, Black Rot, Brook Spot, Sooty Blotch, Apple Maggot, Codling Moth, Curculio, and Tent Caterptilars. Apply in delayed dormant, pre-bloom, petal fall and 3 to 4 cover sprays. Cover sprays should be at 7 to 10 day intervals following petal fall. Do not apply within 7 days of harvest. May injure longe on some straintes.

PEACHES, APRICOTS: Apply 8 oz. as a dust, or mix 8 oz. in 4 gals, of water per tree. For control of Brown Rot, Scab, and Curcutio, apply in delayed dormant, pre-bloom, petal fall, shucksplit, and 2 cover sprays at 7-10 day intervals and a pre-harvest spray 3 weeks before harvest. Do not apply within 21 days of harvest.

PEARS: Apply 8 oz. as a dust or mix 8 oz. in 4 gals, of water per tree. For the controt of Scab, Fruit Spot, Pear Stub, and Codling Moth. Spray at 7-10 day intervals following petal tall. Do not apply within 7 days of harvest.

STRAWBERRIES: Apply 2 oz. as a dust or mix 2 oz. in 1 gal, of water per 200 sq. ft. for the control of Botrytis Rot, and Tarnished Plant Bug. For Root Weeval control, application should be made before planting and worked into the top 6-8 inches of soil. Do not apply later than 14 days before harvest.

NOTICE OF WARRANTY

Plate warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such tabel only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of the product. Crop injury, ineffectiveness, or other unlintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantabrity or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Methoxychlor Mixture RQ NA2761

ORM-A

FORMULATED FOR

PLATTE CHEMICAL COMPANY, INC. 150 SO. MAIN STREET FREMONT, NEBRASKA 68025

FRUIT GUARD

ACTIVE INGREDIENTS:	
Malathion (O,O-Dimethyl dithiophosphate	
of diethyl mercaptosuccinate)	5.0%
Methoxychior Technical*	15.0%
Captan (N-Trichloromethylthio-4-cyclo-	
hexene-1,2-dicarboximide)	9.5%
Related Derivatives	
INERT INGREDIENTS:	
	100.0%

HOME GARDEN USE

*15% Methoxychlor technical is equivalent to 13.2%.

2,2-Bis (p-Methoxyphenyl)-1,1, 1-Trichloroethane and 1.8%

other isomers and reaction products.

KEEP OUT OF REACH OF CHILDREN DANGER

See Side Panel for Additional Precautionary Statements.

EPA REG. NO. 34704-141

EPA EST. NO. _____

NET WEIGHT _____ POUNDS

14349

* CLEAN CROP is a Registered T.M. of United Agri Products, Inc.

EXP10