

CAPTAN-STREPTOMYCIN

7.5-01

POTATO SEED PIECE PROTECTANT

ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	7.33%
Related Derivatives	0.17%
*Streptomycin	0.01%

BY WEIGHT

INERT INGREDIENTS:	92.49%
TOTAL	100.00%

*Streptomycin from streptomycin sulfate (Agricultural grade)

**KEEP OUT OF REACH OF CHILDREN
DANGER**

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician.
If swallowed: Call physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. Do not give anything by mouth to an unconscious person.
If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.
If on skin: Remove contaminated clothing and wash affected area with soap and water.

See Side Panel for Additional Precautionary Statements

FPA REG. NO. 34704-156

EPA EST. NO. 38091-ID-1

NET WEIGHT 40 POUNDS

© CLEAN CROP is a Registered T.M. of United Agri Products, Inc.

17142

EXP-23

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. **NOTE TO USER:** Federal Government and most states have seed laws and regulations requiring an appropriate caution statement be affixed to bags containing treated seed such as: "Treated Seed—Do Not use for food or feed."

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Protect from excessive heat. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

After work: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

This material is especially formulated for potato seed piece treatment.

POTATO (Seed Piece Treatment): For use as an aid in the control of Tuber Soft Rot, Black Leg and surface borne Verticillium Wilt. Dust at 10 ounces per bushel cut seed pieces (16 ounces or 1 pound to 100 pounds cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 to 2 days, the treated pieces should be stored for 2 or 3 days in open crates before bagging.

Do not use treated seed for food or feed purposes.

NOTICE

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of the product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Not licensed for use or export outside of the United States, its Territories and Possessions, Canada or Mexico.

U.S. Pat. Nos. 2,533,770—2,553,771.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025