

11/2 1794 x 8 3/4

LINDANE 25W



FOR INSECT CONTROL ON CERTAIN FLOWERS,
VEGETABLES, TREES, SHRUBS
AND FARM ANIMALS.

ALSO FOR USE IN CERTAIN FARM BUILDINGS TO
CONTROL LISTED INSECTS.

ACTIVE INGREDIENT:		
Lindane (Gamma isomer of benzene hexachloride)	25.0%
INERT INGREDIENTS:	75.0%
	TOTAL	100.0%

**KEEP OUT OF REACH OF CHILDREN
WARNING AVISO**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le expliquen lo hayo sido explicado simplemente.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

ACCEPTED

EPA REG. NO. 34704-174

DEC 17 1990

EPA EST. NO. _____

POUNDS

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 34704-174

02790

mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Wash hands thoroughly after using and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not use on cats or other household pets nor on poultry or humans.

This product should not be applied under conditions which might result in a toxic residue, or residual odor or taste remaining on foodstuffs or edible portions of treated crops or forage at harvest time. Do not use on forage crops to be fed to dairy animals or animals being finished for slaughter. Do not use in poultry houses or dairy barns. Do not treat forage which is to be sold commercially or shipped interstate. Do not use at higher concentrations than recommended on this label. Applicators must wear water resistant hat, lightweight protective suit or coveralls, unlined waterproof gloves (natural rubber, neoprene, or polyethylene), and unlined lightweight boots. MSHA/OSHA approved respirators are required for applications. In enclosed areas such as crawl spaces. Mixer/loaders must wear goggles or a face shield, waterproof apron, and unlined waterproof gloves. Avoid exposure to children. Do not allow children to handle or apply this product. Do not allow children or pets in treated areas until surfaces are dry.

WARNING
(Insert area or field description) treated with Lindane on (insert date of application.) Do not enter without appropriate protective clothing for 24 hours. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Lindane and describe his condition. For further information see the "STATEMENT OF PRACTICAL TREATMENT" portion of the pesticide label.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment listed under "PRECAUTIONARY STATEMENT" when handling open containers.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

APPLICATIONS

Do not apply this product through any type of irrigation system. Aerial application of Lindane is prohibited.

FARM BUILDINGS

Except poultry houses, dairy barns, milk rooms, FLIES, MOSQUITOES, ROACHES, ANTS, BEDBUGS, TICKS: For residual control in farm buildings, use 1 lb. per 10 gal. water. Before applying spray, remove livestock and cover feed, utensils, and feed and water troughs. Do not con-

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, birds, and other wildlife. Do not apply directly to water or wetlands. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by disposing of equipment washwaters. This pesticide is highly toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

STATEMENT OF PRACTICAL TREATMENT

If swallowed—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled—Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin—Remove contaminated clothing and wash affected area with soap and water.

If in eyes—Flush eyes with plenty of water. Call a physician if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive reentry intervals for various

quarters of animals, and rums frequented by pests. Repeat as necessary.

FARM ANIMALS

Except Dairy Animals

Do not use on lactating dairy animals. Note: Spray pressure should be delivered at 200 psi in order to penetrate the animals' hair to the skin.

SARCOPTIC MANGE MITES, LICE, TICKS: Do not treat animals in cold, stormy weather. Do not treat overheated or sick animals. Make sure animals are well watered prior to being dipped to prevent them from drinking the treated dip. Do not spray on animals within 30 days of slaughter. Do not dip animals within 60 days of slaughter. Do not use sprays containing more than 0.06%. Do not use on young animals including calves under 3 months of age. Other immature animals may be injured by treatment. Spray or dip application to young animals (over 3 months of age) should be limited to 0.03% concentration of Lindane.

SARCOPTIC MANGE MITES ON BEEF CATTLE: 1 to 1 1/2 lbs. per 100 gals. water (0.03% to 0.045%) and repeat at 7-10 day intervals as needed. Use lower dosage on animals under 6 months. Make thorough application.

LICE and TICKS ON BEEF CATTLE, SHEEP, HORSES and GOATS: 1 lb. per 100 gals. water (0.03%). Use as spray or dip, thoroughly wetting animals. Repeat treatments at 7 to 10-day intervals may be necessary to prevent reinfestation.

SARCOPTIC MANGE MITES and LICE ON HOGS: 2 lbs. per 100 gals. water (0.06%). Complete coverage including inside of ears is necessary. Apply two sprays one week apart.

Do not treat sows within two weeks of farrowing nor at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

Use dip solutions must be disposed of in accordance with the Resource Conservation and Recovery Act (RCRA). If the applicator generates more than 1000 kg of used dip solution per month or more than 1000 kg used dip solution in combination with other hazardous waste, the material must be treated as a hazardous waste subject to part C of RCRA. Any user who wishes to treat, store or dispose of hazardous waste must obtain a permit to serve as a hazardous waste facility pursuant to RCRA.

FLOWERS AND VEGETABLES

For vegetable crops, apply foliage the designated amount of product in 100 to 200 gals. of water per acre. **CUCUMBERS, CANTALOUPE, SQUASH AND PUMPKINS:** For control of cucumber beetles, thrips - use 1.2 to 2 lbs. per acre. Aphids, nematodes, leaf miner - use 1.5 to 2 lbs. per acre. Remove excess residues from cucumbers and squash. Do not use more than 2 lbs. of this product per acre. Do not apply to cucumbers and squash within one day of harvest.

SPINACH: Aphids, leaf miners - use 1.5 to 3.0 lbs. per

GLADIOLUS: Thrips—1 lb. per 100 gals. of water applied as a foliar spray.

ORNAMENTAL TREES AND SHRUBS

FLOWERING FRUITS, ELM, BEECH, MAPLE, SYCAMORE, BIRCH: Aphids—1 lb. per 100 gals. water. Apply when new growth is about 2 inches long. For application to ornamental trees only.

AZALEA, RHODODENDRON, OAK AND SYCAMORE: Lace bugs—1 lb. per 100 gals. water. Apply just after hatching when small nymphs can be seen. Second spray 10 days later. For application to ornamentals only.

PINES (ornamentals only): Pine bark aphid - 1 lb. per 100 gals. water. Apply when immature aphids are present.

NOTICE

DISCLAIMER—It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Platte Chemical Co. Platte Chemical Co. warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages.

LINDANE MIXTURE RQ NA2761

ORM-A



FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025