



BETASAN
7-G

MODE OF APPLICATION

The most effective program for control of crabgrass, annual bluegrass and other annual weeds is a fall application of BETASAN 7-G. The rate of application should be based on the type of lawn and the degree of infestation. A second application in early spring will provide additional control.

QUICK-LAWN BETA-LAWN - For the best results, follow the instructions on the label for the use of these products.

<ul style="list-style-type: none"> Crabgrass Annual bluegrass Common bermudagrass Parrot grass 	<ul style="list-style-type: none"> Crabgrass Annual bluegrass Parrot grass Parrot grass 	<ul style="list-style-type: none"> Crabgrass Annual bluegrass Parrot grass Parrot grass
--	---	---

DICHONDRA LAWNS: BETASAN 7-G can be applied to dichondra lawns at the time of seeding or immediately after.

ORNAMENTALS AND GROUND COVERS: BETASAN 7-G should be applied only to the following listed ornamentals and ground covers.

<ul style="list-style-type: none"> Alyssum Arter Bachelor's-Button Catclaw Candy-Tuft Chrysanthemum Daylily Drumstick Freesia Hebe Jasmin Juniper Parrotwood Shrub Spirea Yucca Campanula Garland 	<ul style="list-style-type: none"> Coral Bell Ornamental Primrose Glaucous Narcissus Juniper Monterey Cypress Monterey Pine Myrtle Privet Hypericum Ivy Plant Ivy Trichysandra 	<ul style="list-style-type: none"> Stock Sweet Pea Wallflower Zinnia Ranunculus Tulip Pyreantha Sandankwa Star Jasmine Todora Xyrisma Periwinkle Sedum Wild Strawberry
---	--	--

KEEP OUT OF REACH OF CHILDREN
WARNING

STATEMENT OF PRACTICAL TREATMENT

Call a Physician Immediately! If known exposure occurs or if poisoning is suspected, do not wait for symptoms to develop. Immediately initiate the recommended procedures described below and simultaneously contact a physician, the nearest hospital or Poison Control Center.

Inform the person contacted of the type and extent of exposure, describe the victim's symptoms and follow the advice given.

NOTE: Be sure to advise the person contacted that the active ingredient in this compound is a cholinesterase inhibitor and follow the advice given.

NOTE TO PHYSICIAN: Exposure to the active ingredient may cause cholinesterase inhibition. Atropine injection is an antidote. 2-PAM is also an antidote when administered early and in conjunction with atropine.

IF SWALLOWED: Call a physician or Poison Control Center immediately. Immediately give large amounts of water and induce vomiting by gagging the victim with fingers placed on the back of the victim's throat. Give water until vomiting ceases. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen, preferably with a mask.

IF ON SKIN: Flush with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Wash clothing before reuse. Get medical attention if irritation occurs.

IF IN EYES: Hold eye/lids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention.

CAUTIONARY STATEMENTS

For control of crabgrass, apply BETASAN 7-G at the recommended rate (see RATES OF APPLICATION) any time from fall (September or October) through early spring, for spring and summer control. With late water or an early spring application, the application must be made prior to crabgrass emergence (SEE PRODUCT INFORMATION).

For control of annual bluegrass and other annual weeds, apply BETASAN 7-G at the recommended rate (see RATES OF APPLICATION) before the weeds emerge from the soil.

Where germination of crabgrass, annual bluegrass and other weeds extends into the late summer or early fall, a second application of BETASAN 7-G at the recommended rate should be made four to five months after the initial application.

BETASAN 7-G should be applied with a lawn spreader or any other applicator that will give uniform coverage. A more uniform application can be made by spreading half of the required amount of BETASAN 7-G over the area to be treated, then applying the remainder at right angles or crosswise to the first application.

RATES OF APPLICATION

Apply a maximum of 357 pounds of BETASAN 7-G per acre (8.2 pounds per 1,000 square feet) per year in two or more applications depending on the weed species and degree of infestation.

If weed specialists are unavailable to provide specific recommendations, follow these directions: **CRABGRASS ONLY:** Apply 107 to 143 pounds of BETASAN 7-G uniformly on one acre (2.4-3.2 pounds per 1,000 square feet).

ANNUAL BLUEGRASS, GOOSEGRASS AND OTHER ANNUAL WEEDS: Apply 180 pounds of BETASAN 7-G uniformly on one acre (4.0 pounds per 1,000 square feet).

Sprinkle the area with water for 10 to 15 minutes after application to wash the BETASAN 7-G down into the soil where the weed seeds are located.

When a weed control program involves two to three applications of BETASAN 7-G for crabgrass, goosegrass and annual bluegrass, the rate of application for goosegrass and annual bluegrass may be reduced to 127 pounds to 143 pounds per acre, using the higher rate for heavy infestations.

BEST DOCUMENT AVAILABLE

