



BETASAN®

3-G

- WEEDS CONTROLLED**
- Smooth Crabgrass (*Digitaria ischaemum*)
 - Hardy Crabgrass (*Digitaria sanguinalis*)
 - Redroot Pigweed (*Amaranthus retroflexus*)
 - Barnyardgrass (Watergrass) (*Echinochloa spp.*)
 - Lambsquarters (*Chenopodium album*)
 - Guofo grass (*Elousine indica*)
 - Annual Bluegrass (*Poa annua*)
 - Shepherdspurse (*Capsella bursa-pastoris*)
 - Deathwax (*Ranibit; Lamium spp.*)

DIRECTIONS FOR USE

It's a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- Do not contaminate water, food, or feed by storage or disposal.
- STORAGE:** Store in cool dry place.
- CONTAINER DISPOSAL:** Do not re-use empty bag. Completely empty bag by shaking and tipping sides and bottom to loosen clinging particles. Empty residue into application equipment. With empty bag end up in trash collection.

TIMING OF APPLICATION

GRASS LAWNS: BETASAN 3.6-G should be applied only to well established grass lawns. BETASAN 3.6-G can be used on the following established turf grasses:

Bahia Bermuda grass spp. Centipede grass spp. Perennial Ryegrass spp.	Centipede Fescue spp. Pensacola Bahia Perennial Ryegrass	Poa trivialis Saint Augustine Red Top Zoysia
--	---	---

DICHONDRA LAWNS: BETASAN 3.6-G can be applied to dichondra lawns at the time of seeding or any time thereafter.

BEST DOCUMENT AVAILABLE

ORNAMENTALS AND GROUND COVERS: BETASAN 3.6-G should be applied only to the following well established ornamentals and ground covers.

Alyssum Aster Bachelor's-Button Candula Candy-Tuft Daffodil Dianthus Freesia Abelia Azalea Azara Boxwood Aucuba Campanula Guzmania	Corn Bell Daisy Marigold Pansy Primrose Gladolus Narcissus Juniper Monterey Cypress Monterey Pine Myrtle Hypericum Ivy Plant Ivy	Pachysandra Stock Sweet Pea Wild Flower Zinnia Ranunculus Tulip Pyracantha Succulents Star Jasmine Tobacco Purwinkle Sedum Wild Strawberry
--	---	---

For control of crabgrass, apply BETASAN 3.6-G at the recommended rate (see RATES OF APPLICATION) any time from fall (September or October) through early spring, for spring and summer control. With a late winter or an early spring application, the application must be made prior to crabgrass emergence (see PRODUCT INFORMATION).

For control of annual bluegrass and other annual weeds, apply BETASAN 3.6-G at the recommended rate (see RATES OF APPLICATION) before the weeds emerge from the soil.

Consult weed specialists of the State Agricultural Experiment Station or the State Agricultural Extension Service for additional information, as the timing of application may vary with local conditions.

Where elimination of crabgrass, annual bluegrass, and other weeds extends into the late summer or early fall, a second application of BETASAN 3.6-G at the recommended rate should be made four to five months after the initial application.

BETASAN 3.6-G should be applied with a lawn spreader or any other applicator that will give uniform coverage. A more uniform application can be made by spreading half the required amount of BETASAN 3.6-G over the area to be treated, then applying the remainder at right angles or crosswise. Avoid misses by overlapping slightly.

RATES OF APPLICATION

CRABGRASS ONLY: Apply 12 to 16 pounds of BETASAN 3.6-G uniformly on 2,500 square feet.
ANNUAL BLUEGRASS AND OTHER ANNUAL WEEDS: Apply 20 pounds of BETASAN 3.6-G uniformly on 2,500 square feet.

Sprinkle the area with water for 10 to 15 minutes after application to wash the BETASAN 3.6-G down into the soil where the weed seeds are located.

RESEEDING

Any material that provides effective control of annual grasses may affect new lawn grass seedlings which germinate after treatment. Proper fertilization and watering will encourage the existing turf grass to fill in the bare spots formerly occupied by crabgrass.

The following procedures are suggested in order to minimize any stunting of turf grass which may occur when reseeding into BETASAN-treated areas:

- Do not reseed within 4 months after application.
- Recreate the area to be reseeded by raking thoroughly to scuff the soil surface. Where practical, make the new soil to provide an optimum seed bed.
- When turf grasses or other grasses are ordered, place orders for reseeding sooner than 4 months after BETASAN application. Apply powdered activated charcoal or to reseeding to deactivate BETASAN 3.6-G.

Apply the powdered charcoal at a rate of 7 pounds in 14 gallons of water per 1,000 square feet (30 pounds in 60 gallons of water per acre).

Following application, irrigate the turf with sufficient water to wash the charcoal into the soil. Reseed no later than 7 days after applying the charcoal.

NOTICE

Platte Chemical Company makes no warranty, express or implied, including the warranties of merchantability and fitness for any particular purpose, concerning this material, except those which are contained on the label.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

CALL A PHYSICIAN IMMEDIATELY if a known exposure occurs or poisoning is suspected, do not wait for symptoms to develop. Initiate first aid and the recommended procedures described below and immediately contact a physician, the nearest hospital or a Poison Control Center.

NOTICE TO PHYSICIAN: Exposure to the active ingredient may cause cholinesterase inhibition. Atropine is the antidote of choice. 2-PAM is also indicated when administered early and in conjunction with atropine.

IF SWALLOWED: Call a physician or Poison Control Center immediately. Immediately give large amount of water and induce vomiting by giving 1 cup of water with a bent object placed on the back of the throat. Give fluid until vomit is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen, preferably with a physician's advice.

IF ON SKIN: Flush with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Give medical attention if irritation occurs. Wash clothes before reuse.

IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Give medical attention if irritation persists.

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

This product is sold only for uses stated on its label. No express or implied license is granted to use or to reproduce in any country except as specified. Country: United States of America.

BETASAN® is a registered trademark of Stauffer Chemical Company.

NET WT. _____ LB.

EPA Reg. No. 34704-
EPA Est.

PRECAUTIONARY STATEMENTS

TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Keep off swiftness.
- Keep off swiftness.
- Do not get in eyes, on skin, or on clothing.
- Do not eat, drink, or smoke while handling and before eating or smoking.
- Wash hands and face with soap and water before re-use.
- Wash contaminated clothing with soap and water before re-use.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning, or disposing of containers or wastes. Do not apply when conditions favor drift from treated areas.

GENERAL USE PRECAUTIONS

- READ ALL DIRECTIONS BEFORE USING.
- BETASAN 3.6-G should be used only for recommended uses and at recommended rates.
- Do not use on lawns or other areas where birds or other animals are present.
- Do not use on lawns or other areas until BETASAN 3.6-G has been watered into the soil.
- BETASAN 3.6-G is recommended for use on mineral soils only.
- Do not apply to lawns, ground covers, or established ornamentals before treating with BETASAN 3.6-G.

PRODUCT INFORMATION

BETASAN 3.6-G is recommended for use on crabgrass, annual bluegrass, and other annual weeds on ornamentals and ground covers for control of crabgrass, annual bluegrass, and other annual weeds. For control of annual bluegrass and other annual weeds, BETASAN 3.6-G must be applied before the weeds emerge from the soil. For control of crabgrass, BETASAN 3.6-G can be applied any time from fall through early spring, for spring and summer control.

Remove leaves, dead or dying grass, and other debris before applying BETASAN 3.6-G to turf areas. Any weeds present around ornamentals must be removed or watered into the soil before applying BETASAN 3.6-G.

ACCEPTED
JAN 19 1983
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 34704-210