



BENEFIN[®] WEED & FEED

Active Ingredient

N-butyl-N-ethyl-a,a-trifluoro-2,6-dinitro-p-toluidine* 0.29%

Inert Ingredients (relative to herbicidal content) 99.71%

Total 100.00%

*Balan[®] — the registered trademark for Elanco Products Company benefin

KEEP OUT OF REACH OF CHILDREN CAUTION:

In case of contact for eyes, flush with water for 15 minutes; for skin, wash with plenty of soap and water. Get medical attention if irritation persists. See back panel for additional precautionary statements and statement of practical treatment.

A fertilizer and selective pre-emergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

Bluegrass (perennial)	Bahiagrass
Ryegrass (perennial)	Bermudagrass
Fescue	Centipedegrass

Treats 2,500 sq. ft.

40 LBS NET WT (18.14 Kg)

EPA Establishment No. 34704-0

EPA REG. NO. 34704-214

CLEAN CROP is a Registered T.M. of United Agri Products, Inc.

ACCEPTED
MAY 25 1983
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 34704-214

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Keep out of reach of children. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not apply when weather conditions favor drift from target area. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in dry place. Do not contaminate feed & foodstuffs, seeds, fertilizers or other pesticides.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

*Benefin W-F is a pre-emergence herbicide fertilizer which is applied to established turfgrass for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. Benefin W-F controls these annual weeds by killing their seeds as they germinate. Apply Benefin W-F before these annual weeds begin to germinate since it does not control established weeds.*

APPLICATION RATE

Application rate: This bag (40 lbs.) will treat 2,500 sq. ft. For smaller areas use the conversion table below.

Square feet	Pounds of Benefin W-F
500	8
1,000	16
1,500	24
2,000	32

FREQUENCY AND TIMING OF APPLICATION

Successful pre-emergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply Benefin W-F approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application time does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed performance will be obtained if treated areas are irrigated soon after application. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Benefin W-F.

1 Spring germinating summer annual grasses—crabgrasses, foxtails and barnyardgrass.

A Northern (cool season) turfgrasses

For pre-emergence control, apply in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

B Southern (warm season) turfgrasses

For pre-emergence control, apply in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of these annual weed grass species continues for extended periods of time, a second application of Benefin W-F at the recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

2 Late spring, early summer germinating annual grasses—goosegrass.

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications of Benefin W-F should, therefore, be timed accordingly.

A Northern (cool season) turfgrasses

For pre-emergence control apply in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

When Benefin W-F is applied as directed, the following established turfgrasses are tolerant:

Kentucky bluegrass	Zoysia grass
Perennial ryegrass	Bermudagrass
Centipedegrass	St. Augustinegrass
Fescue	Bahiagrass

Benefin W-F will not harm most nearby established ornamental trees and shrubs.

B Southern (warm season) turfgrasses

For pre-emergence control apply in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3 Northern and southern turfgrass

In areas of the United States where pre-emergence control of both early germinating annual grasses (crabgrass, foxtail and barnyardgrass) and the later germinating goosegrass is desired, two applications of Benefin W-F at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4 *Poa annua* (annual bluegrass)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

A Northern (cool season) turfgrass

For pre-emergence control apply in late summer or early fall and again in late winter or early spring one to two weeks prior to the expected germination of annual bluegrass.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying Benefin W-F. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using Benefin W-F, the amount of seed used should be increased and

P
%

FORMULATED FOR **PLATTE CHEMICAL CO., Inc.**
150 SO. MAIN STREET
FREMONT, NEBRASKA 68029