

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
(AND DOMESTIC ANIMALS)**

**WARNING**

May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing. Causes eye and skin irritation. Applicators who apply this product manually or without the use of a closed-system treatment procedure must wear the following protective clothing during the application process: long-sleeved shirt, long pants, wear clean synthetic rubber gloves and goggles, and a disposable paper dust mask which covers at least one third of the face. This product should be applied in a well-ventilated area. Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking. Routinely shower or bathe after work and wash work clothes with soap and hot water before reusing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to birds and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all unused treated seeds by burial away from bodies of water.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions before using and use only as specified on this label.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to: positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.


**STORAGE:** Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet condition, can affect package strength. Personnel should use caution and equipment listed under PRECAUTIONARY STATEMENTS when handling open containers.

**PESTICIDE DISPOSAL:** Empty container wastes are toxic. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**GENERAL:** Thoroughly mix the recommended amount of this product into the required amount of water for the slurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations. Store all treated seed in a cool dry place.

**IMPORTANT:** Treated seed must not be used for or mixed with food or animal feed or processed for it. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act.



**FOR AGRICULTURAL USE ONLY**

|   |               |
|---|---------------|
| <b>ACTIVE INGREDIENT:</b>                                   | <b>BY WT.</b> |
| <b>LINDANE (Gamma Isomer of Benzene Hexachloride)</b> ..... | <b>25.0%</b>  |
| <b>INERT INGREDIENTS:</b> .....                             | <b>75.0%</b>  |
| <b>TOTAL</b>  | <b>100.0%</b> |

**KEEP OUT OF THE REACH OF CHILDREN**

**WARNING**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED**—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF INHALED**—Remove victim to fresh air. Apply respiration if indicated.

**IF ON SKIN**—Remove contaminated clothing and wash affected areas with soap and water.

**IF IN EYES**—Flush eyes with plenty of water. Call a physician for further treatment.

See side panel for additional precautionary statements.

**STIR CONTENTS THOROUGHLY INTO WATER BEFORE USING.**

**NET CONTENTS**

**EPA REG. NO. 34704-219 EPA EST. NO.**

**WARRANTY**

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of merchantability or fitness for a particular use or for any other express or implied warranty except as stated above.

CLEAN CROP is a Registered T.M. of United Agr. Products, Inc.

964

11Y83 A

(Directions for use continued)

**SPECIAL USE PRECAUTIONS**

**IMPORTANT:** Treated seed must not be used for human consumption or as animal feed. Bags containing treated seed should be labeled.

**TREATED SEED  
DO NOT USE FOR FEED, FOOD,  
OR OIL PURPOSES**

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

**PRODUCT INFORMATION**

LINDANE 25 WP is a microfine wettable powder agricultural insecticide designed especially for the treatment of seeds against wireworms, seed corn maggots, and certain other soil-borne insects. It contains an effective binding agent to help hold the Lindane on the seed during sowing, storing, and planting. It is designed for use with slurry treater slurry treaters.

**SEED TREATMENT DOSAGE RATES**

| TYPE OF SEED                                      | OUNCES PER 100 POUNDS |
|---|-----------------------|
| Alfalfa Clover                                    | 2.02                  |
| Barley  | 4.02                  |
| Beans   |                       |
| Bush Snap and all others except Limas             | 2.02                  |
| Baby Bush Limas                                   | 2.02                  |
| Standard and Fordhook Large Limas                 | 1.02                  |
| Cantaloupe, Cucumber, Watermelon, Squash, Pumpkin | 4.02                  |
| Carrots, Onions                                   | 4.02                  |
| Corn  | 4.02                  |
| Cotton  |                       |
| Acid Delinted                                     | 8.02                  |
| Machine Delinted                                  | 8.02                  |
| Fuzzy, Undelinted                                 | 9.02                  |
| Grass (Safflower, Rye, Sudan)                     | 2.02                  |
| Milo, Sorghum, Okra                               | 4.02                  |
| Lettuce   | 4.02                  |
| Oats  | 2.02                  |
| Peas  | 4.02                  |
| Soybean   | 2.02                  |
| Spinach, Sunflower                                | 4.02                  |
| Sugar Beets                                       | 4.02                  |
| Tomatoes  | 4.02                  |
| Wheat   | 4.02                  |

**NOT REVIEWED**  
 Based on Draft Label for 5/19/84  
 5/19/84

ONLY SEED OF HIGH GERMINATION with good vigor should be treated with Lindane. Poor seed may suffer greater injury. If sown, germinated that seed be treated not over six weeks before planting. What effect prolonged storage has on germination has not been definitely established. Treated seed should be stored in a dry place until planted so that excess moisture may evaporate. If seed is stacked or when wet, heat may develop which impairs germination.

IF SOIL TEMPERATURE IS EXTREMELY HIGH due to hot weather, midsummer planting, or other reasons, reduce recommended dosage by one-half. Where seed decay occurs due to soil organisms combine with a protective fungicide.

**RQ Lindane Mixture  
NA2761**



FORMULATED BY

**PLATTE CHEMICAL COMPANY, INC.**  
180 SO. MAIN STREET  
FREMONT, NEBRASKA 68005

