

34704-220

-1-

date, all remaining inventories of affected products will fall under the Existing Stocks provisions stated in our letter of October 29, 1963.

Sincerely yours,

George T. LaRocca  
Product Manager (15)  
Insecticide-Fodenticide Branch  
Registration Division (9S-767)

Enclosure

P.S. Since this product is applied as a liquid add the following precautionary statements to the precautionary text: "wear clean synthetic rubber gloves and goggles".

BEST DOCUMENT AVAILABLE

(DIRECTIONS FOR USE CONT.)

**SPECIAL USE PRECAUTIONS**

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled.

**TREATED SEED  
DO NOT USE FOR FEED, FOOD,  
OR OIL PURPOSES**

"Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

**PRODUCT INFORMATION**

LINDANE 75 WP is a microfine wettable powder agricultural insecticide designed especially for the treatment of seeds against wireworms, seed corn maggots, and certain other soil-borne insects. It contains an effective binding agent to help hold the Lindane on the seeds during sacking, storing, and planting. It is designed for wet application through slurry treaters.

**SEED TREATMENT DOSAGE RATES**

TYPE OF SEED	OUNCES PER 100 POUNDS
Alfalfa Clover	2.5 oz
Barley	1.5 oz
Beans	
All except Standard & Fordhook Large Limas	2.5 oz
Standard and Fordhook Large Limas	1.5 oz
Broccoli, Brussels Sprouts, Cabbage	
Cauliflower, Radish	2.5 oz
Cantaloupe, Cucumber, Pumpkin	
Squash, Watermelon	1.5 oz
Corn	1.5 oz
Cotton, Acid or Moisture Sensitive	1.5 oz
Cowpeas	1.5 oz
Flax	2.0 oz
Lettuce	1.5 oz
Milo, Sorghum	1.5 oz
Oats	2.5 oz
Okra	1.5 oz
Onions	1.0 oz
Peas	1.0 oz
Rye	2.5 oz
Safflower	2.0 oz
Sorghum	1.5 oz
Soybeans	1.5 oz
Spinach	1.0 oz
Sudan Grass	1.5 oz
Sugar Beets	4.0 oz
Sunflower	1.5 oz
Tomatoes	2.5 oz
Wheat	1.5 oz

Only seed that is free of mold and other fungi should be treated with Lindane. If seed has been previously treated, it is suggested that seed be treated no later than six weeks before planting. What effect prolonged storage has on germination has not been definitely established. Treated seed should be stored in a dry place until planted so that excess moisture may evaporate. If seed is stacked when wet, he it may develop which impairs germination.

If soil temperature is extremely high due to hot weather and summer planting, or other reasons, reduce recommended dosages by one half. Where seed decay occurs due to soil organisms, combine with a protective fungicide.

**NOTICE**

**DISCLAIMER -** It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Platte Chemical Company. Platte Chemical Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use thereon described when used in accordance with the directions for use, subject to the risks referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages.

ACCEPTED  
with COMMENTS  
in EPA Letter # 100-100

MAR 10 1984

Under the provisions of the  
Federal Insecticide, Fungicide,  
and Rodenticide Act, this product  
is registered under the name  
of Lindane 75 WP

**PLATTE COMPANY, INC.**  
FREMONT, NEBRASKA 68025



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
(AND DOMESTIC ANIMALS)**

**WARNING**

May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing. Causes eye and skin irritation. Applicators who apply this product manually or without the use of a closed-system treatment procedure must wear the following protective clothing during the application process: long-sleeved shirt, long pants, gloves, and a disposable, paper dust mask which covers at least one-third of the face. This product should be applied in a well-ventilated area. Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking. Routinely shower or bathe after work and wash work clothes with soap and hot water before reusing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seed by burial away from bodies of water.

**DIRECTIONS FOR USE**

It is a violation of State and Federal law to use this product in a manner inconsistent with its labeling. Read all directions before using and use only as specified on this label.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**ORAGE:** Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or excess water that cannot be used according to label directions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL:** Triple rinse immediately and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration if permitted by State and Local Authorities.

**GENERAL:** Thoroughly mix the recommended amount of this product into the required amount of water for the slurry-treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations. Store all treated seeds in a cool dry place.

**IMPORTANT:** Treated seed must not be used for animal feed with food or animal feed, or processed for feed. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store treated seeds beyond planting time. Seeds and containers treated with this product must be labeled in accordance with the requirements of the Federal Seed Act.



**LINDANE  
75 WP  
DYED SEED  
TREATER**

**AGRICULTURAL  
INSECTICIDE**

**WETTABLE  
POWDER**

**ACTIVE INGREDIENT:**

**BY WT.**

**LINDANE (Gamma Isomer of Benzene Hexachloride) 75.0%**

**INERT INGREDIENTS: 25.0%**

**TOTAL 100.0%**

**FOR AGRICULTURAL USE ONLY  
KEEP OUT OF THE REACH OF CHILDREN**

**WARNING**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED**—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.  
**IF INHALED**—Remove victim to fresh air. Apply respiration if indicated.  
**IF ON SKIN**—Wash the affected area with soap and warm water.  
**IF IN EYES**—Immediately flush eyes with running water for at least 15 minutes. Call a physician.

See side panel for additional precautionary statements

**STORE IN A COOL, DRY PLACE.**

**STIR CONTENTS THOROUGHLY INTO  
WATER BEFORE USING.**

**NET CONTENTS U.S. GALLONS**

**EPA REG. NO. 34704-220**

**EPA EST. NO. 34704-**

OLF MICHOLSON is a Registered T.M. of United Agri. Products, Inc.

**RO Lindane Mixture NA2761**

**ORM-A**

FORMULATED BY

**PLATTE CHEMICAL**  
150 SO. MAIN STREET



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

4/22/79  
3/7/79

MAR 23 1979

SEC 115 1.7  
17/79

BEST DOCUMENT AVAILABLE

TO: DIRECTOR, EPCAA  
FROM: SAC, [illegible]  
SUBJECT: [illegible]  
[illegible text]

ON APRIL 12, 1979, THE FOLLOWING INFORMATION WAS RECEIVED FROM [illegible]... [illegible text]... [illegible text]

[illegible text]... [illegible text]... [illegible text]

