

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

~~34704-230~~  
18/11

JUL 19 1995

Mr. J. Allen Dunlap III  
Platte Chemical Company  
P.O. Box 667  
Greeley, CO 80632-0667

Dear Mr. Dunlap:

Subject: Amendment- Revise per Diazinon Registration Standard  
Clean Crop Diazinon G-14  
EPA Registration Number 34704-230  
Your submissions dated December 15, 1993

This amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable, provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Under DIRECTIONS FOR USE change "Federal Law" to "Federal law".

b. Incorporate all the changes in the Agency's letter of March 3, 1994 for the Worker Protection Standard.

c. Under Statement of Practical Treatment If inhaled: add "Get medical attention."

d. Under Statement of Practical Treatment If on skin: change "Get medical attention." to "Remove contaminated clothing and wash promptly with plenty of soap and water. Rinse thoroughly. Get medical attention."

196

CONCURRENCES

SYMBOL	73055						
SURNAME	ZAO						
DATE	7/14/95						

3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Your current Confidential Statement of Formula (CSF) does not list the impurities of toxicological concern separately nor does it include the diazinon certification for FIFRA Sec. 12(a)(1)(C), that the description of this product refers to the composition set forth on the CSF and supporting materials as required by the Diazinon Registration Standard, as well as the certification regarding the determination of the certified limits of the toxicological significant impurities, i.e., reliance on your diazinon supplier's analysis of their technical. Please check with each of your diazinon suppliers in this regard and update your CSF accordingly.

We note that you list two different suppliers for diazinon. For each supplier you must submit separate CSF forms because the impurities of toxicological concern for diazinon are different for each technical.

Sincerely,

George T. LaRocca  
Product Manager (13)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

# RESTRICTED USE PESTICIDE

Due to Avian and Aquatic Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

with COMMENTS  
in EPA Letter Dated



# DIAZINON G-14

Jul 19 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

34704-230

## GRANULAR INSECTICIDE

FOR CONTROL OF CERTAIN SOIL INSECTS INCLUDING WIREWORMS, CUTWORMS, AND OTHER CROP PESTS.

ACTIVE INGREDIENT: BY WEIGHT

Diazinon: O,O-diethyl O-(2-isopropyl-6-methyl 4-pyrimidinyl) phosphorothioate ..... 14.0%

INERT INGREDIENTS: ..... 86.0%

TOTAL 100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA REG. NO. 34704-230

EPA EST. NO. \_\_\_\_\_

NET WEIGHT \_\_\_\_\_ LBS. ( \_\_\_\_\_ KG)

37014

EXPCG:0Y93

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. Do not store near food or feed products. Do not contaminate ornamental fish ponds.

### ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Keep out of lakes, streams, ponds, tidal marshes and estuaries. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

### STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated.

If on skin: Wash promptly with plenty of soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Unless directions under Timing/Special Directions state otherwise, applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, harrow, or other suitable means for greatest effectiveness.

### WORK SAFETY RULES

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling DIAZINON G-14 do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment section of this label. When handling this product, wear the personal protective equipment listed in the following section.

### PERSONAL PROTECTIVE EQUIPMENT

Long-sleeved shirt, long-legged pants, shoes and socks, and chemical-resistant gloves.

Do not apply this product in a way that will contact unprotected workers, either directly or through drift. Only protected handlers may be in the area during application.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry, well-ventilated area away from food or feed.

Pesticide: Pesticide or equipment residue that cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

Container: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles into application equipment. Then dispose of empty bag in a sanitary landfill or incinerate, or burn if allowed by state and local authorities. Stay out of smoke during burning. Do not reuse container.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean up and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-303-356-4400 or 1-800-424-9300 day or night.

### VEGETABLE CROPS

#### Special Precautions To Reduce Risks To Birds

Birds feeding on treated areas may be killed. To reduce risks to birds:

1. Thoroughly incorporate this product into the soil during or immediately after application. Refer to the Soil Incorporation table that follows.
2. Cover or incorporate spilled granules.

DIAZINON G-14 should be applied uniformly as directed for each crop. When a range of rates is given, use the higher rate for heavy infestations.

For application by ground equipment only. Do not apply this product to food crops grown in greenhouses.

Note: Soil Incorporation - Unless directions under Timing/Special Directions state otherwise, for control of pests listed below DIAZINON G-14 should be incorporated into the soil during or immediately after application.

Insect	Depth of Incorporation
Surface cutworms*	2-3 inches
Subterranean cutworms*	3-6 inches
Mole crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

\*If cutworms cannot be identified, incorporate 4-6 inches.

**DIAZINON G-14**  
**EPA REG. NO. 34704-230**

4 9 6

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Beans (Lima, Pole, and Snap) (succulent varieties only)	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Beets, Red (Table)	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Broccoli	Root Maggots	14-21 lbs. per acre	Broadcast and incorporate into soil just before planting or transplanting.
	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Brussels Sprouts	Root Maggots	14-21 lbs. per acre	Broadcast and incorporate into soil just before planting or transplanting.
	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Cabbage	Root Maggots	14-21 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Carrots	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.

Carrots (cont'd.)	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Cauliflower	Root Maggots	14-21 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Celery	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Collards	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Cucumbers	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Endive (Escarole)	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Ginseng	Cutworms, Climbing cutworms, Wireworms, White grubs	14-28 lbs. per acre.	In new beds, broadcast just before planting and incorporate into the top 4-8 inches of soil by tilling with hand-held or powered ground equipment.
		21 lbs. per acre	When soil insect damage appears in established beds, broadcast in the Spring or late Summer through early Fall. Incorporate mechanically with 1/2 inch of irrigation.

NOTE: Do not apply more than once per growing season. Do not apply within 12 months of harvest.

5 9 6

**DIAZINON G-14**

**EPA REG. NO. 34704-230**

Kale	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Lettuce	Cutworms	14-18 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians and Hybrids of these) and Watermelons	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Mustard	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
Onions (Bulb and Green)	Onion Maggots	14-28 lbs. per acre	Broadcast just before planting and incorporate into the top 3-4 inches of soil.  NOTE: Diazinon will not control organophosphate-resistant onion maggots.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Parsley	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Peas (succulent varieties only)	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Peppers	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.

Potatoes	Cutworms (Surface & Subterranean)	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
NOTE: Do not use on commercially grown potatoes which will be hand-harvested.			
Radishes	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
Spinach	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Squash (Summer and Winter)	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Sweet Corn	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Seed Corn Maggots	14-28 lbs. per acre	Broadcast just before planting and incorporate into the soil 2 inches.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Sweet Potatoes	Wireworms, Flea Beetles	21 lbs. per acre	Broadcast and incorporate into soil just before planting.
Swiss Chard	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Tomatoes	Cutworms	14-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.
Turnips	Mole Crickets	7 lbs. per acre	Broadcast and incorporate into plant beds or fields 1-2 days before sowing of seed or transplanting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorporate into soil just before planting.

**FIELD CROPS**

**Special Precautions To Reduce Risks To Birds:**  
 Birds feeding on treated areas may be killed. To reduce risks to birds:

1. Thoroughly incorporate this product into the soil during or immediately after application.
2. Collect or incorporate granules that are spotted during loading or are visible on soil surface in turn areas.

DIAZINON G-14 should be applied uniformly as directed for each crop

6 7 6

**DIAZINON G-14**

**EPA REG. NO. 34704-230**

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Sugar Beets	Sugar Beet Root Maggots	7-14 lbs. per acre or 4.7- 9.4 oz. per 1,000 linear ft. of row.	Apply in a 5-7 inch band (based on 22" row spacing) incorporated 2-3 inches at planting.
	Wireworms	21-28 lbs. per acre	Broadcast and incorpo- rate into the soil 4-8 inches just before planting.
NOTE: Do not apply more than 28 lbs. of DIAZINON G-14 per acre to the soil per season.			

**NOTICE**

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

FORMULATED FOR  
**PLATTE CHEMICAL CO.**  
150 SO. MAIN STREET      FREMONT, NEBRASKA 68025-5697

6