

34704-231

2/27/2002

1/9

RESTRICTED USE PESTICIDE

Due to Avian and Aquatic Toxicity
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.



DIAZINON 500-AG

APPROVED

FEB 27 2002

34704-231

Organophosphate Insecticide

For control of certain insects on fruits, nuts, vegetables, and field crops.

ACTIVE INGREDIENT:

Diazinon [O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]48%

INERT INGREDIENT: *52%

TOTAL 100%

This product contains 4 lbs. Diazinon per gallon.

*Contains xylene range aromatic solvent.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at applications rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

KEEP OUT OF REACH OF CHILDREN CAUTION

EPA REG. NO. 34704-231 EPA EST. NO. 61282-WI-1
NET CONTENTS _____ GAL. (_____ L)

See Below For Additional Precautionary Statements and Directions For Use.

IMPORTANT: Read the entire Directions for Use and the Notice before using this product. If terms are not acceptable, return the unopened product container at once.

IHT

02Y02

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not contaminate ornamental fish ponds.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as: barrier laminate, butyl rubber, nitrile rubber or viton; and shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID STATEMENTS

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person or convulsing person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should only be administered in conjunction with atropine. This product also contains xylene range aromatic solvent which may present an aspiration hazard.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Reformulation or repackaging of this product is prohibited.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other

DIAZINON 500-AG

EPA REG. NO. 34704-231

persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI).

Exception: if the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves, such as: barrier laminate, butyl rubber, nitrile rubber or viton; and shoes plus socks.

Note: The California Department of Agriculture has set a 5-day reentry period for Diazinon on grapes, peaches, and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

APPLICATION THROUGH AN IRRIGATION SYSTEM (CHEMIGATION)

DIAZINON 500-AG may be applied alone or in combination with other pesticides registered for application through sprinkler (including center pivot lateral move, and four side (wheel) roll, traveler, big gun, solid set or hand move) or drip (trickle) irrigation systems (including surface and subsurface). Do not apply this product through any other type of irrigation system. To insure compatibility, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for 5 minutes. If the combination remains mixed, or can be remixed readily, the mixture is compatible.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect the irrigation system used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

Chemigation Systems Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Sprinkler and Drip (Trickle) Chemigation

1. The system must contain a functional check-valve, a vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being

withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not use sprinkler chemigation when wind speed favors drift beyond the area intended for treatment.

Operating Instructions

1. Determine the acreage to be treated by the system.
2. Measure the appropriate amount of DIAZINON 500-AG needed to treat the acreage to be treated. Refer to the specific directions for use for the application rate and the amount of water per acre.
3. Add the pre-measured DIAZINON 500-AG to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if DIAZINON 500-AG is tank mixed with other pesticides registered for sprinkler or drip chemigation.
4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the DIAZINON 500-AG to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

Note: When treatment with DIAZINON 500-AG has been completed, further field irrigation over the treated area should be avoided for 48 hours to prevent washing the product off of the crop.

Apply DIAZINON 500-AG only in sufficient irrigation water to achieve thorough plant coverage. Excess water will wash the product from the plant resulting in poor control.

To avoid a hazard to animals, wells or adjoining crops, do not allow irrigation water to collect or runoff.

WORKER SAFETY RULES

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling this product do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment Section of this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture or rinsate cannot be disposed of by use according to label instructions, contact State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate, or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-303-356-4400 or 1-800-424-9300 day or night.

GENERAL INFORMATION

DIAZINON 500-AG is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use in greenhouses, barns, or dwellings. Do not use on livestock.

In the directions for use which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

FRUIT AND NUT CROPS

REI - 24 hours

DIAZINON 500-AG may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The rate of DIAZINON 500-AG for aerial spray or concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gallons of water

DIAZINON 500-AG

EPA REG. NO. 34704-231

per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the Rate calls for 1 pt. of DIAZINON 500-AG in 100 gallons of water, this amounts to 3-4 pts. of product per acre when applied at the usual dilute ground spray volumes of 300-400 gallons of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the Timing/Special Directions for individual crops.

CROP	Aerial Application MINIMUM Gals./Acre	Ground Application MINIMUM Gals./Acre	
		DILUTE	CONCENTRATE
Almonds*	20	100	20
Apricot	20	100	20
Blackberries**	20	100	20
Boysenberries**	20	100	20
Cherries	20	100	20
Cranberries	20	15	-
Dewberries**	20	100	20
Grapes	20	100	20
Loganberries**	20	100	20
Nectarines	20	100	20
Peaches	20	100	20
Plums	20	100	20
Prunes	20	100	20
Raspberries**	20	100	20
Strawberries	20	100	20
Walnuts*	20	100	20

*CA only.

**CA, OR, or WA only.

Observe stated time intervals between last application and harvest, as well as intervals between applications and total amount of product that can be applied per season.

CROP	*PHI
Almond	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Cranberries	7
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums, Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

*PHI = Pre-Harvest Interval or the number of days to wait between the last application and harvest; may vary based on use pattern selected. See specific directions for use on grapes.

** = Dormant spray only.

Note: To protect bees, do not apply this product to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Almonds (CA only)	San Jose Scale, Parlatoria Scale	1-1½ pts. + 2-3 gallons dormant oil or 1-1½ per acre.	Apply only as a single dormant spray. Do not apply more than 6 pts. of diazinon or 9 gallons of oil
Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Crawlers	1 pt. in 100 gallons of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
	Olive Scale Crawlers	½ pt. + 1½ gallons of light medium horticultural oil per 100 gallons of water	Apply when scale crawlers are present.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Apricots cont'd.	Apricot Mealybugs	1 pt. per 100 gallons of water	Apply as a cover spray from petal fall to June. Allow 14 days between applications.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Caneberries (CA, OR, and WA only); Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Leafhoppers, Thrips, Aphids, Two-spotted Spider Mites, Raspberry Fruitworms, Raspberry Sawflies	1 qt. per acre	Apply as insects occur. Repeat application after 14 days, if necessary. Apply in a minimum of 100 gallons of water per acre. (200 gallons per acre maximum). For Raspberry Fruitworms make one application when blossom buds separate. Make a second application immediately before the blossoms open.
	Dryberry Mites	1-2 qts. per acre	Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed but still closed. Apply in a minimum of 100 gallons of water per acre.
	Raspberry Crown Borers (Raspberry Root Borers)	2 qts. per acre	Apply a single application in a minimum of 100 gallons of water per acre as a drench to the crown and lower canes in the Spring before buds break.

Notes: (1) Do not apply more than 2 qts. of product per acre per application and no more than 5 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.

Cherries	Eyespotted Bud Moths, Fruit Tree Leafrollers	1 pt. in 100 gallons water	Apply in cover sprays as of infestations occur beginning at petal fall. Repeat application after 7 days if necessary.
	Leafhoppers	1 pt. in 100 gallons of water	Make 3 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications.
	Cherry Fruitflies	½-1 pt. in 100 gallons of water	Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10 day intervals.
	Black Cherry Aphids	1 pt. in 100 gallons of water	Apply as a cover spray when aphids appear. Repeat application after 10 days if necessary.
	San Jose Scale Crawlers	1 pt. in 100 gallons of water	Apply during the pre-pink stage or when crawlers first appear.
	Cherry Rust Mites	½-1 pt. in 100 gallons of water	Apply to trees as a single supplemental spray after harvest.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Cranberries	Blackheaded Fireworms (<i>Rhopobota naevana</i>)	2 qts. in a minimum of 15 gallons of water per acre for ground application and up to 400 gallons of water per acre for chemigation.	Apply to larval stage. Pheromone trap captures and sweep monitoring may be used to optimize treatment timing. Repeat application after 14 days, if necessary. A maximum of 6 applications per season is permitted.
	Cranberry Fruitworms	2-3 qts. in a minimum of 15 gallons of water per acre for ground application and up to 400 gallons of water per acre for chemigation.	Cranberry growth stage and/or berry inspection for eggs may be used to optimize treatment timing; consult your local extension agent or pest management advisor. Repeat application every 14 days, as needed. A maximum of 4 applications per season is permitted for the 3 qt. rate. A maximum of 6 applications is permitted for the 2 qt rate

**DIAZINON 500-AG
 EPA REG. NO. 34704-231**

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
<i>Cranberries</i> <i>cont'd.</i>			
Notes: (1) Do not apply more than 12 qts. of product per acre per season (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest. (3) This product may be applied to cranberries through sprinkler irrigation. Refer to the section CHEMIGATION for specific chemigation directions for use. (4) Do not use water from irrigated or flooded cranberry beds for drinking purposes or to irrigated crops other than those appearing on EPA approved diazinon labels.			
Grapes	Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leafrollers, Omnivorous Leafrollers, Aphids, Spider Mites	1-2 pts. per acre in a minimum of 100 gallons of water.	Apply as a thorough cover spray as pest occurs. Repeat application after 7 days, if necessary.
	<i>Drosophila</i> spp.	6 fl. oz. per acre in a minimum of 100 gallons of water (300 gallons of water maximum)	Apply as a thorough cover spray as pest occurs. Repeat application after 7 days, if necessary.
Notes: (1) Do not apply more than 2 pts. of product per acre per application and no more than 10 pts. per acre per season. (2) To avoid illegal residues, allow 28 days between the last application and harvest.			
Nectarines	Oriental Fruitmoths	1 pt. in 100 gallons of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight.
	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, Peach Twig Borers	1 pt. in 100 gallons of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 4 qts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.			
Peaches	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 pt. in 100 gallons of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
	Oriental Fruitmoths	1 pt. in 100 gallons of water	Apply at 100 % petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight.
	Peach Twig Borers	1 pt. in 100 gallons of water	Apply as insects occur in May and June. Allow 7 days between applications.
	Leafhoppers	1 pt. in 100 gallons of water	Make 3-4 applications beginning at petal fall and continuing through the growing season. Allow at least 7 days between applications.
	White Peach Scale	1½-2 pts. in 100 gallons of water	Apply post-harvest timed to coincide with peak crawler and immature scale activity. Under conditions of heavy infestation 2 applications (7 days apart) may be needed. Do not apply before fruit is harvested.

4/9

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
------	------	------	---------------------------

<i>Peaches</i> <i>cont'd.</i>			
Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.			
Plums	Brown Mites (Clover Mites), European Red Mites	1 pt. in 100 gallons of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	½-1 pt. in 100 gallons of water	
Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.			
Prunes	Brown Mites (Clover Mites), European Red Mites	1 pt. in 100 gallons of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	½-1 pt. in 100 gallons of water	
Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.			
Strawberries	Aphids, Two-spotted Spider Mites	1 pt. in 100 gallons of water per acre	Apply as pests occur. Repeat application if necessary after 7 days.
	Cyclamen Mites	1 qt. in 100 gallons of water per acre	Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat applications after 7 days, if necessary. Do not make more than 3 foliar applications per season.
	Mole Crickets	1 qt. per acre	Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1-2 days before transplanting and immediately incorporate into the top 1-2 inches of soil.
	Strawberry Leafrollers	¾-1 pt. in 100 gallons of water	Apply 200 gallons of spray per acre when blossoms show color.
Notes: (1) Do not apply more than 1 qt. of product per acre per application and no more than 4 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 5 days between the last application and harvest.			
Walnuts (CA only)	Aphids, Spider Mites	1 qt. per acre	Apply as a foliar spray. Repeat application in 14 days, if necessary. Make ground applications in a minimum of 100 gals. of water.
	Codling Moths, Scale Crawlers, Walnut Caterpillars	3 qts. per acre	
Notes: (1) Do not apply more than 3 qts. of product per acre per application and no more than 9 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 45 days between the last application and harvest and do not apply after husks open.			

VEGETABLE CROPS

REI - 24 hours

For foliar or soil applications, DIAZINON 500-AG should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. DIAZINON 500-AG may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

DIAZINON 500-AG
EPA REG. NO. 34704-231

CROP	GROUND APPLICATION		AERIAL APPLICATION
	MINIMUM GALLONS ACRE	MINIMUM GALLONS 1000.SQ. FT.	MINIMUM GALLONS ACRE
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabas	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5
Chinese Radish (CA, FL only)	10	2	5
Collards	10	2	5
Crenshaws	5	1	5
Endive (Escarole)	10	2	5
Ginseng	10	2	-
Honeydew Melons	5	1	5
Kale	10	2	5
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	5
Mustard	10	2	5
Onions (Bulb and Green)	10	2	5
Persian Melons	5	1	5
Radishes	10	2	5
Spinach	10	2	5
Sweet Corn	20	4	5
Tomatoes	10	2	5
Watermelons	5	1	5

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply this product in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

Crop	PHI (Days)
Beets, Red (Table)	14
Broccoli	7
Brussels Sprouts	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli	10
Chinese Cabbage	10
Chinese Mustard	10
Chinese Radish	10
Collards	10
Crenshaws	3
Endive (Escarole)	14
Ginseng	30
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Mustard	10
Onions (Bulb and Green)	14
Persian Melons	3
Radishes	14
Spinach	14
Sweet Corn	7
Tomatoes	1
Watermelons	3

*PHI = Pre-Harvest Interval or the number of days to wait between the last application and harvest.

Notes: (1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see TIMING/SPECIAL DIRECTIONS), using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. (2) To Protect Bees: Foliar applications of this product should not be made to peas or beans if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS	
Beets, Red (Table)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.	
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.	
	Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.	
	Broccoli	Root Maggots	2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
			4-8 oz. per 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
		Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
		Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
		Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Brussels Sprouts	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Root Maggots		2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.	
		4-8 oz. in 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.	

5/9

DIAZINON 500-AG
EPA REG. NO. 34704-231

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Brussels Sprouts <i>cont'd.</i>	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage-worms	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur.
Cabbage	Root Maggots	2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. in 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 21 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Broccoli	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
Chinese Mustard (Gai Choy)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
Chinese Radish (Japanese Radish or Daikon) (FL and CA only)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.
		1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.
		1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.
		1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.

CROP PEST RATE TIMING/SPECIAL DIRECTIONS 6/9

Cauliflower	Root Maggots	2-3 qts. per acre	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.	
		4-8 oz. in 50 gallons of water	In transplant water as a drench application when 200-300 gallons of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.	
Cabbage	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.	
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.	
	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, up to 5 times with a minimum of 7 days between applications. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.	
	Chinese Broccoli	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
	Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Mustard (Gai Choy)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.	
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.	
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.	
		½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.	
Chinese Radish (Japanese Radish or Daikon) (FL and CA only)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.	
		1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.	
		1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.	
		1 pt. per acre	Mix in a minimum of 100 gallons of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.	

**DIAZINON 500-AG
 EPA REG. NO. 34704-231**

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Chinese Radish (Japanese Radish or Daikon) (FL and CA only) cont'd.			Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Collards	Cutworms	2-4 qts. per acre.	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage-worms, Harlequin bugs, Corn Earworms	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
Endive (Escarole)	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
	Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice	¾-1 pt. per acre	Spray when insect first appear. For heavy insect infestations, use 1 pt. per acre. Apply by ground equipment using 10 to 100 gallons of water per acre. Do not apply more than one application per growing season. Do not apply during flowering of three or four year old crops. Do not apply within 30 days of harvest. Do not graze treated areas or feed treated forage to livestock.
Kale	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Kale cont'd.	Aphids, Diamondback Moths, Imported Cabbage-worms, Harlequin bugs, Corn Earworms	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur.
Lettuce	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians and Hybrids these) and Watermelons	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	½-1½ pts per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Mustard	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Onions (Bulb & Green)	Onion Maggots	2-4 qts. per acre	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Diazinon will not control organophosphate-resistant onion maggots.
	Onion Thrips	1 pt. per acre	Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.

**DIAZINON 500-AG
 EPA REG. NO. 34704-231**

8/9

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Onions (Bulb & Green) <i>cont'd.</i>	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into top 4-8 inches of soil.
Radishes	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipteropus, Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Spinach	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Sweet Corn	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Seed Corn Maggots	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Corn Earworms	1-1¼ qts. per acre	Apply when silks first appear. Repeat application every 7 days if necessary. Do not make more than 5 applications per season.
	Corn Rootworm (Adults)	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Corn Leaf Aphids	1-2 pts. per acre	Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Grasshoppers, Spider Mites, Fleabeetles	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Sweet Corn <i>cont'd.</i>	Sap Beetles	2-2½ pts. per acre	Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Notes: (1) To protect bees, do not apply foliarly to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be picked 7 days after the last application.

Tomatoes	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½ pt. per acre	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms, Beet Armyworms	¾-1 pt. per acre	Do not apply within one day of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
	Vinegar Flies (<i>Drosophila spp.</i>)	½-1½ pts. per acre	

FIELD CROPS

REI - 24 hours

Soil and foliar applications of DIAZINON 500-AG should always be applied as a spray in sufficient water to ensure thorough coverage. For foliar applications, observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

Pre-harvest Interval (PHI*) for Foliar Applications

Crop	PHI* (Days)
Hops	14
Sugar Beets	14

*PHI = Pre-Harvest Interval or the number of days to wait between the last application and harvest.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Hops	Aphids, Spider Mites	1 qt. per acre	Apply as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 4 applications per season. Do not apply within 14 days of harvest.
Sugar Beets	Wireworms	3-4 qts. per acre	Broadcast just before planting. Incorporate into soil 4-6 inches. Do not apply more than 4 qts. of this product to the soil, per season.
	Beet Leafhoppers, Dipterous Leafminers, Aphids	¾-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
	Grasshoppers	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest.

Note: Sugar beet tops may be fed to beef and dairy animals.

1/9

DIAZINON 500-AG
EPA REG. NO. 34704-231

**INSECT CONTROL ON ORNAMENTALS
GROWN OUTDOORS IN NURSERIES**

REI - 12 hours

To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladioli, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Spruce, and Willow, apply the recommended rates indicated below. Try to spray underside of leaves and penetrate dense foliage. Repeat as necessary.

Insects	Amount of DIAZINON 500-AG to Use	
	Rate per 3 gallons of water	Rate per 100 gallons of water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leafiers, Privet Mites, Scale Crawlers (Cottonycushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Twospotted Spider Mites, Juniper Webworms, Whiteflies	½ fl. oz.*	1 pt.**
Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, Fall Webworms, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars	1½ fl. oz.*	3 pts.**

*1 fl. oz. = 2 tablespoons

**16 fl. oz. = 1 pint

Precaution: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.

Mimosa Webworms: Mix ½ oz. in 3 gallons of water (or 1 pt. in 100 gallons of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.
419 18TH STREET, GREELEY COLORADO 80631-5852