

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**DANGER:** Corrosive, causes irreversible eye damage. Do not get into eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not apply when weather conditions favor drift away from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**RE-ENTRY STATEMENT**

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

**STORAGE:** KEEP FROM FREEZING. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can effect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**USE PRECAUTIONS**

Avoid drift of spray mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Use only lawn-type sprayers. Coarse sprays are less

34704-239



Only for Sale To, Use and Storage by Lawncare and Landscaping Personnel or Commercial Applicators

ACTIVE INGREDIENTS:	
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	30.56%
Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	16.34%
Dimethylamine Salt of Dicamba (3,6-dichloro-p-anisic acid)	2.77%
<b>INERT INGREDIENTS:</b>	<b>50.33%</b>
<b>TOTAL</b>	<b>100.00%</b>

This product contains:  
 2.40 lbs 2,4-Dichlorophenoxyacetic acid per gallon or 25.38%  
 1.30 lbs 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 13.50%  
 0.22 lbs 3,6-dichloro-p-anisic acid per gallon or 2.30%  
 Isomer Specific by AOAC Method 6, D01-5

Selective Broadleaf Weed Control in Turfgrasses.  
(To Control Dandelion, Clover, Henbit, Plantains, Wild Onion and Many Other Broadleaf Weeds.)

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.  
**IF ON SKIN:** Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.  
**IF SWALLOWED:** Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.  
**IF INHALED:** Remove victim to fresh air and apply respiration if indicated.

See side panel for additional precautionary statements

**NET CONTENTS**

EPA REG. NO. 34704-

© E.F.A. CROP is a Registered T.M. of United Agr. Products, Inc.



FORMULATED BY

**PLATTE-CHEMICAL**  
150 SO. MAIN STREET

likely to "wind-drift." Use coarse spray droplets. Avoid fine mists. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 85°F. Seed can be safely sown 3 to 4 weeks after application at recommended rate. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

**WEEDS CONTROLLED BY DI-MEC AMINE**

- |              |                  |
|--------------|------------------|
| Bedstraw     | Peppergrass      |
| Black Medic  | Pigweed          |
| Buckhorn     | Plantain         |
| Burdock      | Poison Ivy       |
| Chicory      | Poison Oak       |
| Chickweed    | Purslane         |
| Clover       | Ragweed          |
| Dandelion    | Sheep Sorrel     |
| Dock         | Shepherd's Purse |
| Ground Ivy   | Speedwell        |
| Heal-All     | Spurge           |
| Henbit       | Wild Carrot      |
| Knotweed     | Wild Garlic      |
| Lambsquarter | Wild Lettuce     |
| Lespedeza    | Wild Onion       |
| Mallow       | Yarrow           |
| Morningglory |                  |

**ACCEPTED**  
 MAY 3 1984  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under EPA Reg. No. 34704-239

**St. Augustine Grass:**

the container agitated, spray entire contents over 33,000 square feet (approx. 3/4 acre). Do not overlap between spray patterns.

Apply DI-MEC AMINE at 2-3 pints in up to 150 gallons of water per acre. (1-2 oz DI-MEC AMINE in 5 gallons water per 1,500 square feet.) Do not spray St. Augustine grass if grass is stressed from heat, drought, etc. Slight turf yellowing should disappear after about one week.

On closely mowed bentgrass (putting and bowling greens) apply DI-MEC AMINE at a maximum rate of one fluid ounce in 5 gal. water per 1,500 square ft. - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

**NOTE**

Care should be taken to avoid overdosing bentgrass and St. Augustine grass or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.


**NOTICE OF WARRANTY**

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

08P83

8285

A



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS  
DANGER**

Corrosive, causes irreversible eye damage. Do not get into eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not apply when weather conditions favor drift away from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**PROHIBITIONS.** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

**STORAGE:** KEEP FROM FREEZING. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

**PESTICIDE DISPOSAL.** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or empty container is prohibited by Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL**

**Metal.** Triple rinse immediately, then offer for recycling or reuse, if available. Do not reuse for food or beverage purposes, including household food or beverage containers. Do not reuse for any other purpose.

**Plastic.** Triple rinse immediately, then offer for recycling or reuse, if available. Do not reuse for food or beverage purposes, including household food or beverage containers. Do not reuse for any other purpose.



**DI-MEC AMINE**

Only for Sale To, Use and Storage by Lawncare and Landscaping Personnel or Commerical Applicators

**ACTIVE INGREDIENTS:**

- \*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid . . . . . 30.56%
- \*\*Dimethylamine Salt of 2-(2-methylchlorophenoxy) propionic acid . . . . . 16.34%
- \*\*\*Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid) . . . . . 2.77%

**INERT INGREDIENTS:** . . . . . 50.33%  
**TOTAL** . . . . . 100.00%

This product contains \*2.44 lbs 2,4-Dichlorophenoxyacetic acid per gallon or 25.38%  
\*\*1.30 lbs 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 13.50%  
\*\*\*0.22 lbs 3,6-dichloro-o-anisic acid per gallon or 2.30%  
\*Isomer Specific by AOAC Method 6, D01-5

**Selective Broadleaf Weed Control In Turfgrasses.  
(To Control Dandelion, Clover, Henbit, Plantains, Wild Onion and Many Other Broadleaf Weeds.)**

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

IF EXPOSED: Wash thoroughly with soap and water. If swallowed: Get medical attention. If inhaled: Get medical attention. If on skin: Wash with soap and water.

See side panel for additional precautionary statements

**NET CONTENTS** . . . . . **GALLONS**  
EPA REG NO 34704-239 . . . . . EPA EST. NO.

PLATTE CHEMICAL COMPANY, INC.

**USE PRECAUTIONS**

Avoid drift of spray mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift." Use coarse spray droplets. Avoid fine mists. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 85°F. Seed can be safely sown 3 to 4 weeks after application at recommended rate. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

**WEEDS CONTROLLED BY DI-MEC AMINE**

- |              |                  |
|--------------|------------------|
| Bedstraw     | Peppergrass      |
| Black Medic  | Pigweed          |
| Buckhorn     | Plantain         |
| Burdock      | Poison Ivy       |
| Chicory      | Poison Oak       |
| Chickweed    | Purslane         |
| Clover       | Ragweed          |
| Dandelion    | Sheep Sorrel     |
| Dock         | Shepherd's Purse |
| Ground Ivy   | Speedwell        |
| Heal-All     | Spurge           |
| Henbit       | Wild Carrot      |
| Knotweed     | Wild Garlic      |
| Lambsquarter | Wild Lettuce     |
| Lespedeza    | Wild Onion       |
| Mallow       | Yarrow           |
| Morningglory |                  |

**ORNAMENTAL LAWNS & TURF**

Coil Season Grasses  
Use DI-MEC AMINE at the rate of 3 pints per acre.

Application Type	Water to Use
<b>Normal application</b>	Use 4-80 gallons of water per acre. 12 gallons water per 1000 sq. ft.
<b>Professional Lawn Maintenance</b>	Use 4-80 gallons of water per 1000 sq. ft. Higher water volume may be used when tank capacity allows. Follow label directions for proper amounts.
<b>Lower volume equipment</b>	Use 4-80 gallons of water per acre. Only application equipment that is capable of spreading