

PM23

34704-239

ACCEPTED

OCT - 7 1998

10/7/98

1/2



MEC-AMINE-D®

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. 34704-239

Only for Sale To, Use and Storage by Lawn-care and Landscaping Personnel or Commercial Applicators

ACTIVE INGREDIENTS:

**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	30.56%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	16.34%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	2.77%
INERT INGREDIENTS:	50.33%
TOTAL	100.00%

This product contains 2.44 lbs. 2,4-Dichlorophenoxyacetic acid per gallon or 25.38%.

**1.30 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 13.50%

***0.22 lbs 3,6-dichloro-o-anisic acid per gallon or 2.30%

*Isomer Specific by AOAC Method 6, D01-5

Selective Broadleaf Weed Control in Turfgrasses. To Control: Dandelion, Clover, Henbit, Plantains, Wild Onion and Many Other Broadleaf Weeds.

KEEP OUT OF REACH OF CHILDREN DANGER

See Below for Additional Precautionary Statements.

EPA REG. NO. 34704-239

EPA EST. NO. 37507-MT-1

NET CONTENTS 2% GALLONS (9.46 L)

IHT

EXP08P98

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive, causes irreversible eye damage. Do not get into eyes, on skin or on clothing. Wear face shield or goggles, long pants, long-sleeved shirt, shoes, socks and chemical resistant gloves when mixing, loading or applying this product. Harmful or fatal if swallowed.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

For containers over 1 gallon and less than 5 gallons in capacity: Persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron.

For containers of 5 gallons or more in capacity: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Groundwater Contamination:

Most cases of groundwater pollution involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and

MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Cleaning of Equipment:

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift away from target area. Do not contaminate domestic or irrigation waters.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.

IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air and apply respiration if indicated.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not reenter or permit workers to reenter treated areas until sprays have dried.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: KEEP FROM FREEZING. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift." Use coarse spray droplets. Avoid fine mists. Do not spray roots of ornamentals and trees. Do not exceed specified

2/2

dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 85°F. Seed can be safely sown 3 to 4 weeks after application at recommended rate. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury to other plants.

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

WEEDS CONTROLLED BY MEC AMINE-D

Bedstraw	Henbit	Purslane
Black Medic	Knotweed	Ragweed
Buckhorn	Lambsquarter	Sheep Sorrel
Burdock	Lespedeza	Shepherd's Purse
Chicory	Mallow	Speedwell
Chickweed	Morningglory	Spurge
Clover	Peppergrass	Wild Carrot
Dandelion	Pigweed	Wild Garlic
Dock	Plantain	Wild Lettuce
Ground Ivy	Poison Ivy	Wild Onion
Heal-All	Poison Oak	Yarrow

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

ORNAMENTAL LAWNS & TURF

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Do not apply this product through any type of irrigation system. Do not allow people (other than applicator) or pets on treatment area during application. The maximum number of broadcast applications per treatment site is 2 per year.

Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Bermudagrass and Zoysiagrass

Apply 3 to 4 pints of MEC AMINE-D per acre with spray volumes ranging from 20 to 220 gallons per acre. The equivalent application rate would be 1.1 to 1.5 fluid ounces of MEC AMINE-D per 1,000 sq. ft. with spray volumes ranging from 0.5 to 5 gallons per 1,000 sq. ft. of turf. For mature weeds, apply 3.25 to 4 pints of MEC AMINE-D per acre.

Cool Season Grasses

Use MEC AMINE-D at the rate of 3 to 4 pints per acre.

Application Type	Water to Use
Normal application:	Use 40-80 gallons of water per acre. (1-2 gallons water per 1,000 square ft.)
Professional Lawn Maintenance:	Use 1 to 5 gallons of water per 1,000 square feet. Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.
Lower volume equipment:	Use as little as 5 gallons water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.
Bentgrass:	On closely mowed bentgrass (putting and bowling greens) apply MEC AMINE-D at a maximum rate of one fluid ounce in 5 gal. water per 1,500 square ft. - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

Warm Season Grasses

Use MEC AMINE-D at the rate of 3 to 4 pints per acre

Application Type	Water to Use
Normal Application:	Use 20 to 80 gallons of water per acre (0.5 to 2 gallons of water per 1,000 sq. ft.)
Professional Lawn Maintenance	Use 0.5 to 5 gallons of water per 1,000 sq. ft. Higher spray volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

USE PRECAUTION

Do not apply this product to St. Augustine grass. Care should be taken to avoid overdosing bentgrass as injury may result. Some hybrid bermudagrass may be sensitive to this product. Contact your local Extension Service Weed Control Specialist for sensitive grasses in your area. Large volumes of spray water (i.e. one fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697

