



TRIFLURALIN 4EC

ACTIVE INGREDIENT:

Trifluralin (4,4-difluoro-2,6-dinitro-N,N-dipropyl-p-toluidine) 44.5%

INERT INGREDIENTS: 55.5%

TOTAL 100%

Contains 4.45 pounds active ingredient per gallon.

A selective herbicide for the pre-emergence control of annual grasses and broadleaf weeds.

EPA REG. NO. 34704-242

CAUTION: KEEP OUT OF REACH OF CHILDREN

See label for additional precautions and directions for use.

FORMULATED FOR: PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025

CLEAN CROP is a Registered T.M. of United Agri Products, Inc.

APR 2 1955
EPA 34704-242

INDEX

2

Precautionary Statements	4
Weeds and Grasses Controlled	5
Directions for Use	8
Crop Recommendations	
Allata (Established)	15
Almonds	34
Apricots	34
Beans	15
Castor	
Dry	
Guar	16
Lima	16
Mungbeans	16
Snap	16
Soybeans	25
Broccoli	17
Brussels Sprouts	17
Cabbage	17
Cantaloupes	21
Carrots	16
Castor Beans	15
Cauliflower	17
Celery	16
Cotton	17
Cole Crops	17
Broccoli	17
Brussels Sprouts	17
Cabbage	17
Cauliflower	17
Citrus Trees	33
Collard Greens	21
Cucumbers	21
Cantaloupes	21
Cucumbers	21
Watermelons	21
Dry Beans	15
English Peas	21
Grapefruits	33
Grapes (Vineyards)	33
Greens	21
Collard	21
Kale	21
Mustard	21
Turnip	21
Guar Beans	16
Hops	21
Kale Greens	21
Lemons	33
Lima Beans	16
Mint	21
Peppermint	21

Spearmint	
Mungbeans	
Mustard Greens	
Mustard for Seed	
Nectarines	
Okra	
Oranges	
Peaches	
Peanuts	
Spanish	
Peas	
Dry	
English	
Southern	
Pecans	
Peppermint	
Peppers	
Plums	
Prunes	
Potatoes	
Safflower	
Snap Beans	
Southern Peas	
Soybeans	
Spanish Peanuts	
Spearmint	
Sugar Beets	
Sugarcane	
Sunflower	
Tangeloes	
Tangerines	
Tomatoes	
Trees	
Almonds	
Apricots	
Citrus	
Grapefruits	
Lemons	
Nectarines	
Oranges	
Peaches	
Pecans	
Plums	
Prunes	
Tangeloes	
Tangerines	
Turnip Greens	
Vineyards (Grapes)	
Walnuts	
Watermelons	
Wheat (winter)	

(Buffalograss)
 (Coloradograss)
 Wild Cane (Sorghum bicolor)
 (Shattercane)
 (See Soybean section for special instructions)

BROADLEAF WEEDS CONTROLLED

Carpelweed (Mollugo verticillata)
 Chickweed (Stellaria media)
 Field Bindweed (Convolvulus arvensis)
 (See Trees and Vineyard section for special instructions)
 Florida pusley (Richardia scabra)
 (Florida purslane)
 (Mexican clover)
 (Pusley)
 Goosefoot (Chenopodium hybridum)
 Henbit (Lamium amplexicaule)
 (Fall application only)
 Knotweed (Polygonum aviculare)
 Kochia (Kochia scoparia)
 (Mexican fireweed)
 Lambsquarters (Chenopodium album)
 Pigweeds (Amaranthus sp.)
 (Carelessweed)
 (Rustrate pigweed)
 (Redroot)
 (Rough pigweed)
 (Sowthistle)
 Puncturevine (Western U.S. only) (Tribulus terrestris)
 (Catchup)
 Purslane (Portulaca oleracea)
 Russian thistle (Salsola kali)
 (Tumbleweed)
 Stinging nettle (Urtica dioica)
 (Nettle)

TRIFLURALIN 4EC will not control certain resistant weeds such as cocklebur, jimsonweed, nutsedge (nutgrass), ragweed, velvetleaf or Venice mallow.

Weeds controlled in soybeans by the TRIFLURALIN 4EC/Sencor® or TRIFLURALIN 4EC/Lexone® tank-mix in addition to those controlled by TRIFLURALIN 4EC alone (See Soybean section for special instructions.)

Jimsonweed (Datura stramonium)
 Mallow, Venice (Hibiscus trionum)
 (Flower-of-an-hour)
 Mustard, wild (Brassica kaber)
 (Charlock)
 (Field mustard)
 Ragweed, common (Ambrosia artemisiifolia)
 Sesbania, hemp (Sesbania exaltata)
 (Coffeebean)
 (Indigo)
 Smartweed, Annual (Polygonum pensylvanicum)
 (Pennsylvania smartweed)
 (Smartweed)
 Prickly sida (Sida spinosa)
 (Teaweed)
 (Spiny sida)
 Velvetleaf (Abutilon theophrasti)
 (Butterprint)
 (Buttonweed)
 (Cottonweed)
 (Elephant's Ear)
 (Indian mallow)
 (Piemarker)

Cocklebur, morning-glory and giant ragweed Control of cocklebur, morning-glory and giant ragweed (horseweed) may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation

Weeds controlled in dry beans and potatoes by the TRIFLURALIN 4EC/Eptam® tank-mix in addition to those controlled by TRIFLURALIN 4EC alone (See Drybean, Potato section for special instructions)

Henbit (Lamium amplexicaule)
 (Spring applications)
 (Shade, black) (Solanum nigrum)
 (Shade, hairy) (Solanum sarachoides)
 Nutsedge (Cyperus sp.)
 (Nutgrass)
 (Purple nutsedge)
 (Yellow nutsedge)
 Oat, wild (Avena fatua)

Weeds controlled in soybeans by the TRIFLURALIN 4EC/Amiben® tank mix in addition to those controlled by TRIFLURALIN 4EC alone (See Soybean section for special instructions.)

Ragweed, Common (Ambrosia artemisiifolia)
 Smartweed, Pennsylvania (Polygonum pensylvanicum)
 Velvetleaf (Butterweed) (Abutilon theophrasti)

TRIFLURALIN 4EC preplant soil incorporated with an Amben application pre-emergence controls the following additional weeds

Coltweed (Sesbania)	<i>Sesbania exaltata</i>
Mustard, Wild	<i>Brassica kaber</i>
Nightshade, Black	<i>Solanum nigrum</i>
Prickly sida (Teaweed)	<i>Sida sp. nora</i>
Ragweed, Common	<i>Ambrosia trifida</i>
Spurge, Annual	<i>Euphorbia cyparissia</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Velvetleaf (Buttonweed)	<i>Abition theophrasti</i>

Weeds controlled in cotton by the TRIFLURALIN 4EC/Caparol* tank mix in addition to those controlled by TRIFLURALIN 4EC alone (See Cotton section for special instructions)

Smartweed	Groundcherry (Annual)
Prickly sida (Teaweed)	Mustard
Annual morning-glory	Malva
Ragweed	Wild oat

The tank mix also controls shallow-germinating seedlings of Cocklebur

Weeds controlled in cotton by the TRIFLURALIN 4EC/Cotoran* tank mix or Cotoran overlaid post plant pre-emergence in addition to those controlled by TRIFLURALIN 4EC alone, where TRIFLURALIN 4EC has been applied as a preplant soil incorporated herbicide in cotton (See Cotton section for special instructions)

Ryegrass	Prickly sida (Teaweed)
Buttonweed	Ragweed
Cocklebur	Sesbania
Goathead	Sicklepod
Groundcherry, Wright	Smartweed
Jimsonweed	Tumbleweed
Morning-glory	

Weeds controlled in cotton by an overlay treatment of Karmex* post plant pre-emergence in fields where TRIFLURALIN 4EC has been applied as a preplant soil incorporated herbicide in addition to those controlled by TRIFLURALIN 4EC alone (See Cotton section for special instructions)

Ragweed	Shepherdspurse
Groundcherry (Annual)	Velvetgrass
Dogfennel	Wild lettuce
Pennycress	Wild mustard
Morning-glory, Annual	

The tank mix of TRIFLURALIN 4EC plus Avadex* BW will control wild oat in peas grown in Idaho, Oregon, and Washington in addition to the weeds controlled by TRIFLURALIN 4EC alone (See Pea section for special instructions)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

PROHIBITIONS—Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE—Avoid freezing. Do not store below temperature of (40°F). If frozen, poor weed control may result. Store in safe manner. Store in original container. Keep container tightly closed when not in use. Reduce stacking height where conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL—Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

TRIFLURALIN 4EC is a pre-emergence herbicide which is mixed (incorporated) into the soil to provide long-lasting control of a wide range of annual grasses and broadleaf weeds. TRIFLURALIN 4EC controls weeds as they germinate but will not control established weeds.

SOIL TEXTURE

One key to getting good results with TRIFLURALIN 4EC is to know your soil texture so that you can apply the correct rate. The amount you apply to your soil will vary with the soil texture. A fine-textured soil requires more TRIFLURALIN 4EC than a coarse-textured soil.

Soil Texture Guide:

Refer to the following guide to determine your soil texture

Coarse** Soils	Sand
	Loamy sand
	Sandy loam
Medium Soils	Loam
	Silty clay loam*
	Silt loam
	Silt
	Sandy clay loam*
Fine** Soils	Clay
	Clay loam
	Silty clay loam*
	Silty clay
	Sandy clay
	Sandy clay loam*

* *Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine-textured soils. If silty clay loam or sandy clay loam soils are predominately sand or silt, they are usually classified as medium-textured soils; if predominately clay, they are usually classified as fine-textured soils.*

** *Soil classification nomenclature for light and heavy-textured soils have been changed to coarse and fine-textured soils at the direction of the Environmental Protection Agency. Throughout this label all previous references to light-textured soils have been changed to coarse-textured soils. All previous references to heavy-textured soils have been changed to fine-textured soils. These new descriptions relate more closely the size of the soil particles in a given classification.*

SOIL PREPARATION

Destroy existing weeds before TRIFLURALIN 4EC application. Chop and thoroughly mix crop residues into the soil to a depth of at least 4 to 6 inches by deep plowing or discing before application. Use machinery that breaks up large clods before application.

APPLICATION

Add the recommended amount of TRIFLURALIN 4EC to clean water in the spray tank during the filling operation. Agitate before spraying. Apply in from 5 to 40 gallons of water per acre (broadcast basis), using any properly calibrated low-pressure herbicide sprayer that will apply the spray uniformly. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to insure proper calibration and uniform application. Apply TRIFLURALIN 4EC to the soil surface and incorporate in the same operation, if possible. Do not apply to soils which are wet or in poor condition. Do not apply to soils which are subject to prolonged periods of flooding.

AERIAL APPLICATION

For best results from aerial application of TRIFLURALIN 4EC, apply to a dry soil surface at a spray volume of from 4 to 10 gallons per acre. Adjust pump pressure, nozzle arrangements, flying speed and flying height to provide a uniform application to the soil surface. Use markers to assure proper application spray widths.

Do not apply TRIFLURALIN 4EC by aircraft when the wind is blowing at a velocity of 5 mph or greater. This will cause drift of spray particles and result in non-uniform application.

INCORPORATION DIRECTIONS

Incorporation Before Planting

For best results TRIFLURALIN 4EC should be incorporated as soon as possible after application. TRIFLURALIN 4EC must be incorporated one time within 24 hours after application. A second incorporation is required with most equipment (see Incorporation Equipment section for specific instructions). If TRIFLURALIN 4EC is applied to a wet, warm soil surface or if the wind velocity is 10 mph or higher variable weed control may result from delaying the first incorporation beyond 24 hours.

Incorporation should place the TRIFLURALIN 4EC into the top 2 to 3 inches of the final seedbed. Generally, incorporation equipment will place the chemical approximately half as deep as the equipment is run. For example, a disc running 4 inches deep will incorporate TRIFLURALIN 4EC approximately 2 inches deep.

Incorporation After Planting

(Check crop list for those crops approved for incorporation after planting.)

When incorporating TRIFLURALIN 4EC after planting or on established row crops use PTO-driven equipment or rolling cultivators. Adjust equipment to till the soil over the seed or throw treated soil toward the crop. Avoid disturbing the seed or mechanically damaging the crop.

Incorporation in Bedded Culture

For effective weed control, TRIFLURALIN 4EC should be incorporated into the top 2 to 3 inches of the final seedbed.

Lock off beds to planting height before application and incorporation on bedded ground. If TRIFLURALIN 4EC is applied and incorporated before bedding, do not furrow out deeper than the depth to which TRIFLURALIN 4EC was incorporated. Furrowing too deep will expose untreated soil and allow weeds to germinate in the bottom of the furrow.

Avoid removal of treated soil from the seedbed before or during the planting operation. This will expose untreated soil and allow weeds to germinate in the drill row.

Incorporation Equipment

Use machinery that mixes TRIFLURALIN 4EC thoroughly with the soil. Shallow incorporation with implements set to cut less than 2 inches deep may result in erratic weed control. Use of incorporation equipment not listed upon the label may result in poor or erratic weed control and/or crop injury. Recommended equipment includes:

Disc set to cut 4 to 6 inches deep and operated in 2 different directions at 4 to 6 mph. A tandem or double-disc operated one time does not provide adequate incorporation.

Field cultivator set to cut 3 to 4 inches deep and operated at 5 mph or more. The field cultivator used alone or in combination with the double-disc will provide effective incorporation providing the following instructions are used:

- 1 Two passes over the field with a field cultivator with the second pass running at an angle to the first. Do not set cultivator to cut deeper than 4 inches, particularly on the second pass, since untreated soil may be turned up.
- 2 Field cultivator used for the first pass and the double-disc used for the second pass.
- 3 Double-disc used for the first pass and the field cultivator used for the second pass.

NOTE: A field cultivator is defined as an implement with 3 to 4 rows of sweeps, spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Chisel points should not be used.

Rolling cultivator set to cut 2 to 4 inches deep and operated 2 times at 6 to 8 mph. Rolling cultivators are adequate for use on coarse and medium-textured soils only (except when used in sugarcane where the rolling cultivator may be used on fine-textured soils).

Bed conditioner (Do-All) set to cut 2 to 4 inches deep and operated one time at 4 to 6 mph. Bed conditioners are adequate for use on coarse and medium-textured soils only.

Mulch treader and other similar disc-type implements set to cut 3 to 4 inches deep and operated at 5 to 8 mph in two different directions.

PTO-driven equipment (tillers, cultivators, hoes) set to cut 2 to 3 inches deep with rotors spaced to provide a clean sweep of the soil and operated one time. PTO-driven equipment should not be operated at a speed greater than 4 mph.

Other equipment, including the flexible tine-tooth harrow (Flextine, Melroe) is also recommended but only for the special programs for which it is specified in this label

CULTIVATION AFTER PLANTING

Soil treated with TRIFLURALIN 4EC may be shallow-cultivated, rotary-hoed or hand-hoed without reducing the weed control activity of TRIFLURALIN 4EC. Do not cultivate deeper than the TRIFLURALIN 4EC treated layer of soil since this may bring untreated soil to the surface and poor weed control may result

REGIONAL USE MAP



All crop recommendations are given on a regional basis. The dividing line between the Eastern and Western United States is that point where the average rainfall per year is a minimum of 20 to 25 inches. Use the recommendation in your region only (refer to map)

CROP RECOMMENDATIONS

Eastern United States

GENERAL

These recommendations are given as the broadcast (overall) rates of TRIFLURALIN 4EC per acre. For band applications, use proportionately less. Apply TRIFLURALIN 4EC any time after January 1 when the soil can be worked. TRIFLURALIN 4EC is not recommended on muck soils. Where a rate range is shown, use the lighter rate for more coarse soils or soils with lower organic matter.

Western United States

GENERAL

These recommendations are given as the broadcast (overall) rates of TRIFLURALIN 4EC per acre. For band application, use proportionately less. TRIFLURALIN 4EC is not recommended for peat soils exceeding 20% organic matter or on any muck soils. Do not exceed recommended rates at any time. Where a rate range is shown, use the lighter rate for more coarse soils or soils with lower organic matter.

**TRIFLURALIN 4EC APPLIED ALONE
AND IN COMBINATION WITH SENCOR
WITH FLUID FERTILIZERS**

GENERAL

TRIFLURALIN 4EC alone and TRIFLURALIN 4EC plus Sencor may be mixed with most fluid (liquid) fertilizer materials. TRIFLURALIN 4EC alone and TRIFLURALIN 4EC plus Sencor with solutions and suspension-type fertilizers has provided weed and grass control as claimed on the respective labels.

All recommendations for TRIFLURALIN 4EC alone or TRIFLURALIN 4EC plus Sencor tank-mix combinations regarding rates per acre, approved crops, incorporation, special instructions, warnings and special precautions must be followed.

Individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

COMPATIBILITY TEST

TRIFLURALIN 4EC alone and TRIFLURALIN 4EC plus Sencor and some fluid fertilizer materials may not combine properly. Small quantities should always be tested before full-scale mixing.

1. Put 1 pint of fertilizer mixture in a quart jar
2. Add 2 teaspoonfuls of TRIFLURALIN 4EC and 2 level teaspoonfuls of Sencor as usage dictates
3. Close jar and shake well.
4. Watch mixture for several seconds, check again 30 minutes later
5. If the mix does not separate, or if agitation is only needed to resuspend the Sencor, the combination may be used. If the mixture separates, gets very thick or syrupy, DO NOT combine for field application.
6. Mixing ability may be improved by adding a compatibility agent. Follow the procedure outlined above and add 1/2 teaspoonful of the compatibility agent in Step 2. Complete the other steps to determine if the compatibility agent solves the problem.

If needed use a compatibility agent cleared for use on growing crops

MIXING

If a compatibility agent is needed, add it to the fluid fertilizer before adding the TRIFLURALIN 4EC alone or TRIFLURALIN 4EC plus Sencor combination. If compatibility is a problem, mix 2 quarts of water with 1 quart of TRIFLURALIN 4EC alone or TRIFLURALIN 4EC plus Sencor combination before pouring it into the fertilizer.

TRIFLURALIN 4EC alone or in combination with Sencor may be poured directly into the fluid fertilizer and mixed thoroughly. Wettable powders should be mixed with the liquid fertilizer before adding TRIFLURALIN 4EC. Continued agitation is needed until application is complete.

APPLICATION

Spread the fertilizer/chemical mixture normally with a properly calibrated applicator. Be certain the material is applied uniformly to the soil surface.

INCORPORATION

Follow normal TRIFLURALIN 4EC incorporation procedures.

**TRIFLURALIN 4EC APPLICATION WITH
DRY BULK FERTILIZERS**

GENERAL

Dry bulk fertilizers may be impregnated or coated with TRIFLURALIN 4EC. Application of dry bulk fertilizers impregnated with TRIFLURALIN 4EC has provided weed and grass control equal to the same rates of TRIFLURALIN 4EC applied in water. All TRIFLURALIN 4EC label recommendations regarding rates per acre, approved crops, incorporation, special instructions, cautions and special precautions must be followed.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the blender and/or company selling the fertilizer and chemical mixture.

LIMITATIONS

Apply a minimum of 100 pounds per acre of dry fertilizer impregnated with TRIFLURALIN 4EC at the recommended rates. Any commonly used dry fertilizers can be used for TRIFLURALIN 4EC impregnation *except* straight coated ammonium nitrate and straight limestone. These materials will not absorb the herbicide. Blends containing mixtures of these materials can be impregnated.

IMPREGNATION

Use any closed drum, belt, ribbon or other commonly used dry bulk fertilizer blender. The nozzle or nozzles used to spray the TRIFLURALIN 4EC on to the fertilizer should be placed to provide uniform spray coverage.

RATES

Check the crop section to determine the rate of TRIFLURALIN 4EC per acre. See the rate table which follows to determine the amount of TRIFLURALIN 4EC to be impregnated on a ton of dry bulk fertilizer based on the amount of fertilizer which will be applied per acre. (See rate chart.)

APPLICATION

Spread the fertilizer/chemical mixture normally with a properly calibrated applicator. Be certain the material is applied uniformly to the soil surface.

INCORPORATION

Follow normal TRIFLURALIN 4EC incorporation procedures.

RATE CHART FOR IMPREGNATING FERTILIZER WITH TRIFLURALIN 4EC

TRIFLURALIN 4EC added to a TON of fertilizer.		
Fertilizer Rate Per Acre	TRIFLURALIN 4EC Rate Per Acre	
	1 pint	1½ pints
200 pounds	10 pts or 5 qts per ton	15 pts or 7½ qts per ton
250 pounds	8 pts or 4 qts per ton	6 qts or 1½ gal per ton
300 pounds	6½ pts or 3½ qts per ton	10 pts or 5 qts per ton
350 pounds	5½ pts or 2¾ qts per ton	9 pts or 1 ¼ gal per ton
400 pounds	5 pts or 2½ qts per ton	7½ pts or 1 gal per ton
450 pounds	4½ pts or 2¼ qts per ton	3½ qts or ¾ gal per ton

RATE CHART FOR IMPREGNATING FERTILIZER WITH TRIFLURALIN 4EC cont'd.

TRIFLURALIN 4EC added to a TON of fertilizer.			
Fertilizer Rate Per Acre	TRIFLURALIN 4EC Rate Per Acre		
	2 pints	3 pints	4 pints
200 pounds	10 qts or 2½ gal per ton	15 qts or 3¾ gal per ton	20 qts or 5 gal per ton
250 pounds	8 qts or 2 gal per ton	12 qts or 3 gal per ton	16 qts or 4 gal per ton
300 pounds	14 pts or 1¾ gal per ton	20 pts or 2½ gal per ton	27 pts or 13½ qt per ton
350 pounds	12 pts or 1½ gal per ton	17 pts or 2¼ gal per ton	23 pts or 2 ⅞ gal per ton
400 pounds	5 qts or 1¼ gal per ton	15 pts or 1 ⅞ gal per ton	10 qts or 2½ gal per ton
450 pounds	4½ qts or 1 ⅞ gal per ton	13½ pts or 1¾ gal per ton	9 qts or 2¼ gal per ton

For rates other than those listed above, use the following formula to calculate the amount of TRIFLURALIN 4EC to be impregnated on a ton of dry bulk fertilizer.

$$\frac{\text{Pints TRIFLURALIN 4EC}}{\text{Per Acre}} \times \frac{1000}{\text{lbs Fertilizer Per Acre}} = \frac{\text{Quarts TRIFLURALIN 4EC}}{\text{Per Ton of Fertilizer}}$$

ALFALFA—Established Alfalfa Only: (Western U.S. Only)

Apply to established alfalfa stands at a broadcast rate per acre of 1½ pints on coarse soils and 2 pints on medium and fine soils. Use incorporation equipment that will insure thorough soil mixing with a minimum of damage to the established alfalfa.

BEANS—Castor Beans:

Apply and incorporate TRIFLURALIN 4EC before planting.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1½ pts	1½-1½ pts
Fine soils	2 pts	1½ pts
Soils with 2 to 5% organic matter	1½-2 pts	1½-2 pts
Soils with 5 to 10% organic matter	2 pts	2 pts

BEANS—Dry Beans (Kidney, Navy, Pinto, Etc.):

Apply and incorporate TRIFLURALIN 4EC before planting.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1½ pts	1½-1½ pts
Fine soils	2 pts	1½ pts
Soils with 2 to 5% organic matter	1½-2 pts	1½-2 pts
Soils with 5 to 10% organic matter	2 pts	2 pts

BEANS—TRIFLURALIN 4EC/Eptam/EPTC Tank-Mix for Dry Beans:
 The TRIFLURALIN 4EC/Eptam/EPTC tank-mix effectively controls henbit, black nightshade and nutsedge (nutgrass) in addition to all of the annual grasses and broadleaf weeds listed on the TRIFLURALIN 4EC label. Follow normal procedures for soil preparation. The TRIFLURALIN 4EC/Eptam/EPTC tank-mix should be applied from 2 days before planting (up to planting in the Eastern US).

Apply at a broadcast rate of 1 pint of TRIFLURALIN 4EC and 1½ pints of Eptam/EPTC 7E per acre or up to the label recommended rate for each herbicide depending on soil texture and weed problem. TRIFLURALIN 4EC at 1 pint per acre alone or combination, should not be used on soils containing 5% or more organic matter. Incorporate immediately after application. Follow normal TRIFLURALIN 4EC procedures for cultivation.

Caution: Read the Eptam/EPTC label before using. Observe all cautions and limitations on labeling of all products used in mixtures. The combination of TRIFLURALIN 4EC and Eptam/EPTC should not be used on soybeans, black-eyed peas (beans), lima beans and other flatpooded beans, except Romano. Do not use the foliage from a crop treated with the TRIFLURALIN 4EC/Eptam tank-mix for feed or for grazing.

BEANS—Fall Application in Dry Beans Grown in Idaho, Oregon and Washington Only:
 For dry beans grown in Idaho, Oregon and Washington. Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 1 pint on coarse soils, 1¼ to 1½ pints on medium soils, and 1½ pints on fine soils.

BEANS—Gusn Beans and Mungbeans:
 Apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse soils and 1½ pints on medium and fine soils.

BEANS—Lima Beans and Snap Beans:
 Apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1½ pints on fine soils.

CARROTS:
 Apply and incorporate TRIFLURALIN 4EC before planting

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1½ pts	1¼-1½ pts
Fine soils	2 pts	1½ pts
Soils with 2 to 5% organic matter	1½-2 pts	1½-2 pts
Soils with 5 to 10% organic matter	2 pts	2 pts

CELERY—Both Direct-seeded and Transplant: (Western U.S. Only)
 Apply and incorporate TRIFLURALIN 4EC before planting or transplanting at a broadcast rate per acre of 1 pint on coarse soils; 1¼ to 1½ pints on medium soils; 1½ pints on fine soils; 1½ to 2 pints on soils with 2 to 5% organic matter, and 2 pints on soils with 5 to 10% organic matter.

COLE CROPS—Broccoli, Brussels Sprouts, Cabbage and Cauliflower:
 For Direct-Seeded cole crops apply and incorporate before planting. (Eastern US—Direct-seeded cole crops have exhibited marginal tolerance to recommended rates of TRIFLURALIN 4EC. Stunting or reduced stands may occur.)

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1 pt	1 pt
Fine soils	1½ pts	1 pt
Soils with 2 to 5% organic matter	1½ pts	—
Soils with 2 to 10% organic matter	—	1½ pts

For Transplant cole crops apply and incorporate TRIFLURALIN 4EC before transplanting. Do not apply TRIFLURALIN 4EC after transplanting.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1½ pts	1¼ to 1½ pts
Fine soils	2 pts	1½ pts
Soils with 2 to 5% organic matter	1½ pts	1½ to 2 pts
Soils with 5 to 10% organic matter	2 pts	2 pts

COTTON—Pre-emergence applications:
 Apply and incorporate TRIFLURALIN 4EC before planting, at planting or immediately after planting using the following broadcast rates per acre.

	Eastern US	Western US
Coarse soils	1 pt.	1 pt
Medium soils	1½ pts	1¼ to 1½ pts
Fine soils	2 pts	1½ pts
Soils with 2 to 5% organic matter	1½ pts.	1½ to 2 pts
Soils with 5 to 10% organic matter	2-2½ pts	2 pts

When incorporating after planting (post-plant), care must be taken not to disturb the seed.

Seedling disease may weaken cotton plants and increase the possibility of damage in TRIFLURALIN 4EC. To control seedling disease, use a good fungicide program.

COTTON—Post-emergence applications:
 Apply TRIFLURALIN 4EC any time up to layby, but not less than 90 days before harvest. Direct layby applications to the soil between the rows and beneath emerged cotton plants. Use the same rates as for a pre-emergence application.

COTTON—Fall Application:
 For cotton grown in Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri, Bootheel, North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee and Texas: Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 2 pints on coarse and medium soils and 2½ pints on fine soils. For cotton grown in Arizona, California and Nevada: Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 1½ pints on coarse soils,

2 pints on medium soils and 2 1/2 pints on fine soils. For cotton grown in states other than those listed above: Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 1 pint on coarse soils; 1 1/2 pints on medium soils, 2 pints on fine soils; 1 1/2 pints on coarse soils with 2 to 5% organic matter, and 2 to 2 1/2 pints on soils with 5.1 to 10% organic matter.

COTTON—Fall panicum control:

For the control of fall panicum in the states of Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia, apply and incorporate TRIFLURALIN 4EC at the broadcast rate of 2 pints per acre on both coarse and medium soils.

Plant cotton after early season adverse weather conditions have passed. Do not plant cotton deeper than 1 inch. Crop injury in the form of delayed growth or reduced yields may occur under adverse cool, wet weather conditions when TRIFLURALIN 4EC is used according to these special recommendations.

COTTON—Rhizome Johnsongrass control:

All Cotton Producing States except Arizona and California. Commercially acceptable control of rhizome Johnsongrass can be obtained with a double-rate TRIFLURALIN 4EC program when applied for 2 years in a row.

Soil Preparation—Proper preparation of the soil before application is very important for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the top of the soil. Then follow with a disc before application to cut the rhizomes into small (2 to 3-inch) pieces. This should also destroy any emerged Johnsongrass.

Application—Choose the one application program that best fits your cultural practices:

Spring Application—Apply TRIFLURALIN 4EC any time in the spring before planting for 2 years in a row at a broadcast rate per acre of 2 pints on coarse soils; 3 pints on medium soils and 4 pints on fine soils, OR

Fall Application—Apply TRIFLURALIN 4EC between October 15 and December 31 for 2 years in a row at the same rates as a spring application for the control of rhizome Johnsongrass.

Incorporation Deep incorporation is essential to good rhizome Johnsongrass control. Incorporate TRIFLURALIN 4EC thoroughly with a disc set to cut 4 to 6 inches deep and operate in 2 different directions at 4 to 6 mph.

Cultivation—Some Johnsongrass plants will escape. Timely cultivations during the crop season are necessary to obtain commercially acceptable control. Commercially acceptable control will not be obtained with only 1 year of double-rate TRIFLURALIN 4EC use.

Precautions: Plant cotton after early season adverse, wet-weather conditions have passed. Crop injury in the form of reduced stands and delayed growth will occur under adverse cool, wet-weather conditions early in the season and may result in delayed maturity and reduced yields when TRIFLURALIN 4EC is used according to these recommendations. High quality seed accompanied by a good fungicide program to control seedling diseases in addition to other recommended cultural and chemical practices should be used to minimize crop injury from TRIFLURALIN 4EC.

In the season following either the 1 or 2-year treatments, plant only those crops for which TRIFLURALIN 4EC has been registered as a preplant treatment or injury may result.

COTTON—More Complete Control of Pigweed and Seedling Johnsongrass in Cotton Grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, South Carolina, Tennessee and southern Virginia:

For more complete control of pigweed and seedling johnsongrass, TRIFLURALIN 4EC may be applied preplant at a broadcast rate per acre of from 1 to 1 1/2 pints on coarse soils, from 1 1/2 to 2 pints on medium soils and 2 pints on fine soils except in the state of Louisiana where 3 pints per acre are recommended on fine soils.

Precaution: Plant cotton after early season adverse weather conditions have passed. Do not plant cotton deeper than 1 1/2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when TRIFLURALIN 4EC is used according to these recommendations.

COTTON—More Complete Weed and Grass Control in Certain Counties Along the Texas Gulf Coast:

For more complete control of those weeds and grasses listed in the TRIFLURALIN 4EC label in the Texas Gulf Coast Counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, TRIFLURALIN 4EC may be applied up to 2 weeks before planting at a broadcast rate of 1 1/2 pints on coarse soils, 2 pints on medium soils and 3 pints on fine soils.

Precaution: Plant cotton after early season adverse weather conditions have passed. Do not plant cotton deeper than 1 1/2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when TRIFLURALIN 4EC is used according to these recommendations.

COTTON—TRIFLURALIN 4EC/Caparol[®] tank mix for cotton grown in California, Arizona, New Mexico, and Texas:

The TRIFLURALIN 4EC/Caparol combination will control certain grasses and broadleaf weeds listed on the TRIFLURALIN 4EC label. This combination will also control shallow-germinating seedlings of cocklebur and coffee-weed.

NOTE: This combination will not control sunflower, rhizome johnsongrass, deep-germinating seedlings of cocklebur and sandbur or established perennials such as Bermudagrass. Follow normal TRIFLURALIN 4EC procedures for soil preparation and incorporation. Apply the tank mix combination to the flat soil surface before discing.

Broadcast Rates Per Acre:	TRIFLURALIN 4EC	Caparol 80W
Coarse soils	1 pt.	2 pounds
Medium soils	1 1/2 pts.	2 1/2 pounds
Fine soils	2 pts.	3 pounds

*Do not use on sands and loamy sands. For band applications use proportionately less. TRIFLURALIN 4EC is not recommended for use on muck soils.

Mixing Directions: Carefully follow the procedures on the Caparol 80W label for making a slurry and adding it to a partially filled tank of water. After the Caparol is thoroughly mixed with the partially filled tank of water, add the TRIFLURALIN 4EC and continue filling. Agitate during the filling and spraying operation.

Avoid leaving the spray mixture in the tank without constant agitation. If by-pass agitation is used, it should terminate at the bottom of the tank to minimize foaming.

Incorporation Directions: The first incorporation of TRIFLURALIN 4EC/Caparol should be immediately following application. A second incorporation is required with most equipment. (See incorporation equipment section for further instructions.)

Precautions: Do not apply more than the recommended rate for your soil texture. The combination of TRIFLURALIN 4EC/Caparol should not be used under the following conditions because crop injury may result in the cut areas of newly leveled fields, in areas of excess salt, and where flooding or the beds is likely to happen. Do not plant cotton in tractor wheel depressions or crop injury may result.

On mulch planted cotton, water back only after cotton seedlings get well established. **Crop Rotations:** Cabbage, okra, melons and peas may be planted in the fall after spring application of TRIFLURALIN 4EC plus Caparol.

Winter barley, winter rye and winter wheat can be planted in the fall also, if they are plowed down and not used for food or feed. Refer to the Caparol label for directions, cautions and precautions.

COTTON—TRIFLURALIN 4EC/Cotoran tank mix except in Arizona and California: The TRIFLURALIN 4EC/Cotoran tank mix effectively controls all the annual grasses and broadleaf weeds listed on the TRIFLURALIN 4EC label plus many additional annual grasses and broadleaf weeds (see Weeds Controlled section). Follow normal TRIFLURALIN 4EC procedures for soil preparation. Apply TRIFLURALIN 4EC/Cotoran tank mix in 15 to 40 gallons of clean water per acre using any properly calibrated low pressure herbicide sprayer that will apply the spray uniformly.

Broadcast Rates Per Acre

	TRIFLURALIN 4EC	Cotoran 80W
Coarse soils	1 pt	1 1/4 pounds
Medium soils	1 1/2 pts	2 pounds
Fine soils	2 pts	2 1/2 pounds

Mixing Directions: Carefully follow the procedures on the Cotoran 80W label for making a Cotoran slurry and adding it to a partially filled tank of water. After the Cotoran is thoroughly mixed with the partially filled tank of water, add the TRIFLURALIN 4EC and continue filling. Agitate continuously throughout the filling and application operations. Follow normal TRIFLURALIN 4EC incorporation procedures. Do not leave spray mixture in tank without constant agitation. If by-pass agitation is used, it should terminate at the bottom of the tank to minimize foaming.

Precautions: Do not use the tank mix in Arizona and California. Do not plant crops other than cotton on the treated land within 6 months after the application of TRIFLURALIN 4EC plus Cotoran or injury may occur.

West Texas Only: Do not use the tank mix of TRIFLURALIN 4EC plus Cotoran on sandy, loamy sand or fine sandy loam soils. Do not use on cotton planted in furrows. **Arkansas, Louisiana, and Mississippi Only:** Use 1 pound Cotoran 80W in tank mix with TRIFLURALIN 4EC on sandy loam soils low in organic matter.

New Mexico: Cotton can be planted the next spring. Do not plant treated areas to crops other than cotton on treated land until 1 year after last application. Do not use on sandy or coarse textured soil of less than 1% organic matter.

Do not feed foliage from treated cotton plants or gin trash to livestock. The tank mix of TRIFLURALIN 4EC plus Cotoran is not recommended to be applied in liquid fertilizer.

Refer to the Cotoran label for cautions, precautions, and instructions.

Cotoran overlay: Refer to the Cotoran label for cautions, precautions and instructions.

COTTON—Preplant incorporated TRIFLURALIN 4EC and surface applied, pre-emergence Karmex for weed control in cotton grown east of the Mississippi River plus Arkansas, Southeastern Missouri, Louisiana, and Eastern Texas: Preplant soil incorporated applications of TRIFLURALIN 4EC (see above for rates) may be followed by a surface applied, post-plant, pre-emergence application of Karmex 80W. This combination effectively controls all the weeds controlled by TRIFLURALIN 4EC plus many additional weeds (see weeds controlled section). Apply Karmex 80W at 0.6 to 1.5 pounds per broadcast acre to the soil surface after planting but prior to crop emergence. The higher rates are used on heavier soil types. Do not use Karmex 80W on light (sandy or low organic) soils. Do not use on heavy clay soils above 10 percent organic matter. Consult the Karmex label for additional instructions, cautions and precautions.

CUCURBITS—Cantaloupes, Cucumbers and Watermelons—Post-Plant, emerged in Texas only:

Apply TRIFLURALIN 4EC as a directed spray to the soil between the rows and beneath plants which are in the 3 to 4 true-leaf stage. Set incorporation machinery to throw treated soil toward plants in the row. Care should be taken that incorporation machinery does not damage the plants.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1 1/2-1 3/4 pts	1 1/4 to 1 1/2 pts
Fine soils	1 3/4 pts	1 1/2 pts
Soils with 2 to 5% organic matter	1 1/2-2 pts	1 1/2-2 pts
Soils with 5 to 10% organic matter	2 pts	2 pts

GREENS—Turnip Greens Grown for Processing and All Collard, Kale and Mustard Greens:

Apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

MUSTARD—Grown For Seed in Minnesota and North Dakota Only:

Apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

HOPS: (Western U.S. Only.)

Apply and incorporate TRIFLURALIN 4EC while the crop is dormant at a broadcast rate per acre of 1 pint on coarse soils, 1 1/4 to 1 1/2 pints on medium soils and 1 1/2 pints on fine soils and soils with 2 to 10% organic matter.

PEPPERMINT—Established Peppermint and Spearmint: (Western U.S. Only.)

Apply TRIFLURALIN 4EC at a rate per acre of 1 pint on coarse soils, 1 1/4 pints on medium soils, and 1 1/2 pints on fine soils. Use incorporation equipment that will insure thorough soil mixing with a minimum of damage to the established, dormant mint.

OKRA:

Apply and incorporate TRIFLURALIN 4EC before planting.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1 1/2 pts	1 1/4-1 1/2 pts
Fine soils	2 pts	1 1/2 pts
Soils with 2 to 5% organic matter	1 1/2-2 pts	1 1/2-2 pts
Soils with 5 to 10% organic matter	2 pts	2 pts

PEANUTS—Spanish Peanuts Grown in Texas and Oklahoma Only: Apply and incorporate TRIFLURALIN 4EC before planting, at planting or immediately after planting at a broadcast rate per acre of 1 pint on coarse soils. When incorporating after planting, care must be taken not to disturb the seed.

PEAS—Dry (Western U.S. Only.)

Apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

PEAS—English:

Apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

PEAS—Fall Application in Dry Peas and English Peas Grown in Idaho, Oregon and Washington Only:

For dry peas grown in Idaho, Oregon and Washington. Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 1 pint on coarse soils, 1 1/4 to 1 1/2 pints on medium soils, and 1 1/2 pints on fine soils.

PEAS—Southern Peas:

Apply and incorporate TRIFLURALIN 4EC before planting

	Eastern US	Western U.S.
Coarse soils	1 pt	1 pt
Medium soils	1 1/2 pts	1 1/4-1 1/2 pts
Fine soils	2 pts	1 1/2 pts
Soils with 2 to 5% organic matter	1 1/2-2 pts	1 1/2-2 pts
Soils with 5 1/2 to 10% organic matter	2 pts	2 pts

PEAS—TRIFLURALIN 4EC and Avadex BW tank mix for weed control in peas grown in Idaho, Oregon, and Washington:

The tank mix combination of TRIFLURALIN 4EC plus Avadex BW will provide control of wild oats in addition to other annual grasses and broadleaf weeds controlled by TRIFLURALIN 4EC.

Application Rates: Broadcast 3/4 pint of TRIFLURALIN 4EC on coarser textured soils, 1 pint of TRIFLURALIN 4EC on fine soils. Use 1 1/4 quarts of Avadex BW for all soil textures.

Incorporation Directions: Apply the TRIFLURALIN 4EC plus Avadex BW tank mix and incorporate from 3 weeks before seeding up to immediately before seeding. TRIFLURALIN 4EC and Avadex BW must be thoroughly incorporated into the 2 inches of the soil by 2 incorporations. The first incorporation should be made as soon as possible on the day of application. The second incorporation should be made as soon as possible but before seeding. Incorporate with a disc-type implement set to cut 4 inches deep and operate in 2 different directions at 4 to 6 mph or with a field cultivator set to cut 3 to 4 inches deep and operate at 5 mph or more. Shallow incorporation with implements set to cut less than 2 inches may result in erratic weed control.

NOTE: Do not apply to lentils.

Leaf crinkling and delayed maturity of peas may occur, particularly on clay points in the northwest, but this is usually more than offset by a reduction of wild oats. Do not graze livestock on treated crops. Refer to the cautions, precautions and directions on the Avadex BW label.

PEPPERS—Transplants only:

Apply and incorporate TRIFLURALIN 4EC before transplanting. Do not apply after transplanting.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1 1/2 pts	1 1/4-1 1/2 pts
Fine soils	2 pts	1 1/2 pts
Soils with 2 to 5% organic matter	1 1/2 pts	1 1/2-2 pts
Soils with 5 1/2 to 10% organic matter	2 pts	2 pts

POTATOES—Not recommended for use in the state of Maine:

Apply TRIFLURALIN 4EC after planting, up to or immediately following dragoff in the Eastern US or after planting, before emergence on all soil textures, or after the potato plants have fully emerged on coarse and medium soils in the Western US. TRIFLURALIN 4EC is not recommended on muck soils.

Set incorporation equipment so that the bed and furrow will be uniformly covered with a layer of TRIFLURALIN 4EC. If the layer of TRIFLURALIN 4EC treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. Care should be taken so that incorporation machinery does not damage potato seed pieces or elongating sprouts. Cultivation prior to emergence may result in mechanical injury to the elongated potato sprouts.

When applying and incorporating TRIFLURALIN 4EC after potato plants have fully emerged, do not completely cover the foliage with treated soil. Likewise do not completely cover foliage at subsequent cultivations.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1 1/2 pt	1 1/4-1 1/2 pts
Fine soils	2 pts	1 1/2 pts
Soils with 2 to 5% organic matter	1 1/2 pts	1 1/2-2 pts
Soils with 5 1/2 to 10% organic matter	2 pts	2 pts

POTATOES—Split Application in Idaho, Oregon and Washington Only:

On all soils apply and incorporate 3/4 pint of TRIFLURALIN 4EC before planting and 1/2 pint after planting when potato plants have fully emerged. Do not apply to soils containing 2% or more organic matter. Follow incorporation directions listed above. Application to potatoes after planting.

POTATOES—TRIFLURALIN 4EC/Eptam/EPTC Tank-Mix for Potatoes Grown in Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota and Texas only: (Eastern U.S. Only.)

The TRIFLURALIN 4EC/Eptam/EPTC tank-mix effectively controls henbit, black nightshade and nutsedge (nutgrass) in addition to all of the annual grasses and broadleaf weeds listed on the TRIFLURALIN 4EC label. Follow normal TRIFLURALIN 4EC procedures for soil preparation. The TRIFLURALIN 4EC/Eptam/EPTC tank-mix may be applied after planting but prior to crop emergence. In areas where potatoes are normally dragged-off, the TRIFLURALIN 4EC/Eptam/EPTC tank-mix should be applied and incorporated up to or immediately following dragoff at a broadcast rate per acre of 1 pint of TRIFLURALIN 4EC and 1 1/4 pints of Eptam/EPTC 7E on all soil textures or up to the label recommended rate for each herbicide depending on soil

texture and weed problem. TRIFLURALIN 4EC at 1 pint per acre, alone or in combination, should not be used on soils containing 5% or more organic matter. Incorporate immediately after application. Follow normal TRIFLURALIN 4EC procedures for cultivation.

Caution: Read the Eptam/EPTC label before using. Observe all cautions and limitations on labeling of all products used in mixtures. Do not graze or feed forage to livestock from fields treated with the TRIFLURALIN 4EC/Eptam/EPTC tank-mix.

POTATOES—TRIFLURALIN 4EC/Eptam/EPTC Tank-mix: (Western U.S. Only)
Application After Planting—The TRIFLURALIN 4EC/Eptam/EPTC tank-mix effectively controls henbit, nightshade and nutsedge (nutgrass) in addition to all of the annual grasses and broadleaf weeds listed on the TRIFLURALIN 4EC label. Follow normal procedures for soil preparation. The TRIFLURALIN 4EC/Eptam/EPTC tank-mix may be applied after planting, up to or immediately following dragoff at a broadcast rate per acre of 1 pint of TRIFLURALIN 4EC and 1 1/4 pints of Eptam/EPTC 7E on all soil textures or up to the label recommended rate for each herbicide depending on soil texture and weed problem. TRIFLURALIN 4EC at 1 pint per acre, alone or in combination, should not be used on soils containing 5% or more organic matter. Incorporate immediately after application. Follow normal TRIFLURALIN 4EC procedures for cultivation.

Application Before Planting in Washington, Idaho, and Oregon Only—TRIFLURALIN 4EC/Eptam/EPTC may be also applied before planting at a broadcast rate of 1/4 pint of TRIFLURALIN 4EC and 3/2 pints of Eptam/EPTC 7E on all soil textures and incorporated immediately.

Caution: Do not use this tank-mix both before and after planting in the same season. Read the Eptam/EPTC label before using. Observe all cautions and limitations on labeling of all products used in mixtures. Do not graze or feed forage to livestock from fields treated with the TRIFLURALIN 4EC/Eptam/EPTC tank-mix.

SAFFLOWER:

Apply and incorporate TRIFLURALIN 4EC before planting

	Eastern US	Western US
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.
Soils with 10.1 to 20% organic matter		2-3 pts.

SAFFLOWER—FALL APPLICATION:

For safflower grown in Arizona, California, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming: Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 1 1/2 pints on coarse soils; 2 pints on medium soils; and 2 1/2 pints on fine soils.

SOYBEANS:

Apply and incorporate TRIFLURALIN 4EC before planting. Do not plant soybeans deeper than 2 inches.

	Eastern US	Western US
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2-2 1/2 pts.*	2 pts.

* Except charcoal soils in Arkansas, Louisiana and Mississippi. See below.

SOYBEANS—Soils Containing Charcoal in Arkansas, Louisiana and Mississippi: Newly cleared land often contains high organic matter (4 to 10%) and charcoal which result from burning debris. This charcoal and/or organic matter tends to tie up TRIFLURALIN 4EC and reduce its weed control activity. Higher rates of TRIFLURALIN 4EC are therefore necessary for satisfactory weed control. Increased rates can cause crop injury if charcoal or a high percentage of organic matter is not present to tie up some of the TRIFLURALIN 4EC. In the actual windrow or burn row, where a high level of charcoal is present, poor weed control may result even with an increased rate of TRIFLURALIN 4EC.

Apply and incorporate at the following broadcast rates per acre:

Coarse soils	1 1/2 pts.
Medium soils	2 1/4 pts.
Fine soils	3 pts.

SOYBEANS—Fall application:

For soybeans grown in Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri, North Carolina, Oklahoma, South Carolina, Tennessee and Texas: Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 2 pints on coarse and medium soils and 2 1/2 pints on fine soils. For soybeans grown in the Eastern United States other than those states listed above: Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 1 pint on coarse soils; 1 1/2 pints on medium soils; 2 pints on fine soils; 1 1/2 pints on coarse soils with 2 to 5% organic matter; and 2 to 2 1/2 pints on soils with 5.1 to 10% organic matter.

SOYBEANS—Fall panicum control:

For the control of fall panicum in the states of Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia, apply TRIFLURALIN 4EC at the broadcast rate of 2 pints per acre on both coarse and medium soils. Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth or reduced yields may occur under adverse cool, wet weather conditions when TRIFLURALIN 4EC is used according to these special recommendations.

SOYBEANS—More Complete Control of Pigweed and Seeding Johnsongrass In Soybeans Grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, southeastern Missouri, North Carolina, South Carolina, Tennessee and southern Virginia:

For more complete control of pigweed and seeding johnsongrass, TRIFLURALIN 4EC may be applied at a broadcast rate per acre of from 1 to 1½ pints on coarse soils, from 1½ to 2 pints on medium soils and 2 pints on fine soils except in the state of Louisiana where 3 pints per acre are recommended on fine soils.

Precaution: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

SOYBEANS—More Complete Weed and Grass Control in Certain Counties Along the Texas Gulf Coast:

For more complete control of those weeds and grasses listed in the TRIFLURALIN 4EC label in the Texas Gulf Coast Counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, TRIFLURALIN 4EC may be applied up to 2 weeks before planting at a broadcast rate of 1½ pints on coarse soils, 2 pints on medium soils and 3 pints on fine soils.

Precaution: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

SOYBEANS, RED RICE CONTROL—Arkansas, Louisiana, Mississippi and Texas only:

Suppression or partial control of red rice in soybeans can be obtained when TRIFLURALIN 4EC is applied as directed at double the normal rate the first year (not to exceed 4 pints per acre) and at the normal rate the second year. Follow normal TRIFLURALIN 4EC directions for soil preparation and soil incorporation.

APPLICATION: Year 1

Apply and incorporate TRIFLURALIN 4EC the first year any time in the spring before planting at the following broadcast rates per acre:

Coarse soils	2 pts
Medium soils	3 pts
Fine soils	4 pts

Coarse soils with 2 to 5% organic matter	3 pts
Soils with 5.1 to 10% organic matter	4 pts

APPLICATION: Year 2

Apply TRIFLURALIN 4EC the second year at the following normal label broadcast rates per acre:

Coarse soils	1 pt
Medium soils	1½ pts
Fine soils	2 pts

Coarse soils with 2 to 5% organic matter	1½ pts
Soils with 5.1 to 10% organic matter	2-2½ pts

If a combination of high organic matter (4 to 10%) and charcoal are present in the soil, apply TRIFLURALIN 4EC the second year at the following rates labeled for charcoal soils in Arkansas, Louisiana and Mississippi:

Coarse soils	1½ pts per acre
Medium soils	2½ pts
Fine soils	3 pts

*Newly cleared land often contains high organic matter (4 to 10%) and charcoal which result from burning debris. This charcoal and/or organic matter tends to tie up TRIFLURALIN 4EC and reduce its weed control activity. Higher rates are therefore necessary for satisfactory weed control. Increased rates can cause crop injury if charcoal or a high percentage of organic matter is not present to tie up some of the TRIFLURALIN 4EC. In the actual windrow or burn row, where a high level of charcoal is present, poor weed control may result even with an increased rate.

CROP ROTATION

The program for red rice control in soybeans is a 2-year program. Use the rates listed for first year application and plant soybeans. The second year use the normal TRIFLURALIN 4EC rates listed for your soil type and charcoal level and plant only those crops for which TRIFLURALIN 4EC has been registered as a preplant treatment or injury may result. Do not plant rice the second year. Rice may be planted the third year.

PRECAUTIONS

Plant soybeans after early adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury may occur under adverse cool, wet weather conditions early in the season when TRIFLURALIN 4EC is used according to these double-rate recommendations.

SOYBEANS—Rhizome Johnsongrass Control: (Eastern United States and the State of Texas)

Commercially acceptable control of rhizome Johnsongrass can be obtained with a double-rate TRIFLURALIN 4EC program when applied for 2 years in a row.

Soil Preparation—Proper preparation of the soil before application is very important for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the top of the soil. Then follow with a disc before application to cut the rhizomes into small (2 to 3-inch) pieces. This should also destroy any emerged Johnsongrass.

Application—Choose the one application program that best fits your cultural practices.
Spring Application—Apply TRIFLURALIN 4EC any time in the spring before planting for 2 years in a row at a broadcast rate per acre of 2 pints on coarse soils, 3 pints on medium soils, 4 pints on fine soils, 3 pints on coarse soils with 2 to 5% organic matter, and 4 pints on soils with 5.1 to 10% organic matter. OR

Fall Application—Apply TRIFLURALIN 4EC between October 15 and December 31 for 2 years in a row at the same rates as a spring application for the control of rhizome Johnsongrass. OR

Split Application—Apply as directed under both spring and fall applications for 2 years in a row using the following broadcast rates per acre:

	Spring	and	Fall
Coarse soils	1 pt		1 pt
Medium soils	1½ pts		1½ pts
Fine soils	2 pts		2 pts
Coarse soils with 2 to 5% organic matter	1½ pts		1½ pts
Soils with 5.1 to 10% organic matter	2 pts		2 pts

Incorporation—Deep incorporation is essential to good rhizoma Johnsongrass control. Incorporate TRIFLURALIN 4EC thoroughly with a disc set to cut 4 to 6 inches deep and operate in 2 different directions at 4 to 6 mph.

Cultivation—Some Johnsongrass plants will escape. Timely cultivations during the crop season are necessary to obtain commercially acceptable control. Commercially acceptable control will not be obtained with only 1 year of double-rate TRIFLURALIN 4EC use.

Precautions: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

In the season following either the 1 or 2-year treatments, plant only those crops which TRIFLURALIN 4EC has been registered as a preplant treatment or injury may result.

SOYBEANS—Wild Cane Control:

Wild Cane (Shattercane) can germinate from greater soil depth than most other weed seeds. Several "flushes" or germinating times are common in one season. Commercially acceptable control of wild cane can be obtained with increased rates of TRIFLURALIN 4EC.

Land Preparation—Work your land to destroy existing grasses and weed thoroughly mix crop residues into the soil to a depth of 4 to 6 inches.

Application—Apply TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse soils, 2 pints on medium soils, and 2½ pints on fine soils.

Incorporation—Deep incorporation is essential to good wild cane control. Incorporate (mix) TRIFLURALIN 4EC thoroughly with a disc only set to cut 4 to 6 inches deep and operate in 2 different directions at 4 to 6 mph.

Cultivation—Cultivations during the crop season will also contribute to control. **Precaution:** Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

SOYBEANS—TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone Tank-Mix:

The TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix effectively controls additional weeds than controlled by TRIFLURALIN 4EC alone (see Weeds Controlled section). Follow normal procedures for soil preparation. The TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix should be applied from 2 weeks before planting up to planting in 10 to 40 gallons of water with any low-pressure herbicide sprayer equipped with herbicide tips and screens no finer than 50 mesh in nozzle and in-line strainers.

Broadcast Rates Per acre:		Lexone 5½ W P or
	TRIFLURALIN 4EC	Sencor 50 WP
Coarse soils*	1 pt.	½ pound
Medium soils	1½ pts.	¾ pound
Fine soils**	2 pts.	1 pound
		Lexone 4L or
		Sencor 4L
Coarse soils*	1 pt.	½ pt.
Medium soils	1½ pts.	¾ pt.
Fine soils**	2 pts.	1 pt.

*Do not use on coarse soils with less than 1% organic matter.
**Silty clay loam and sandy clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

Do not plant any crop other than soybeans within 4 months after treatment. Follow normal TRIFLURALIN 4EC procedures for incorporation and cultivation.

NOTE: In those areas of the Mid-South where cocklebur is a serious problem, an overlay of Sencor or Lexone may be preferred to the TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix.

Special Precaution: Applied according to directions and under normal growing conditions, the TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix will not harm the treated crop. Over-application may result in crop injury or soil residue. Even application or improper soil incorporation of the TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix can result in erratic weed control or crop injury.

Seeding disease, cold weather, deep planting, excessive moisture, soil pH over 7.5, high salt concentration, or drought may weaken crop seedlings and increase possibility of damage from the TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix. Under these conditions, delayed crop development or reduced yields may result. **Warning:** Observe all warnings and limitations on labeling of all products used in mixtures. Sencor may be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash clothing thoroughly with soap and hot water before reuse. Do not contaminate feed or food. Keep out of reach of children.

Do not use the foliage from soybeans treated with the TRIFLURALIN 4EC/Sencor or TRIFLURALIN 4EC/Lexone tank-mix for feed or forage. Do not contaminate any body of water nor apply to any area not specified on this label. Do not allow sprays to drift onto adjacent desirable plants.

SOYBEANS—TRIFLURALIN 4EC pre-plant followed by Sencor or Lexone as an overlay treatment for weed control in soybeans:

TRIFLURALIN 4EC effectively controls certain annual grasses and broadleaf weeds. See Sencor or Lexone label for additional weeds controlled. Apply TRIFLURALIN 4EC as a preplant incorporated herbicide. As a separate operation, make a single application of Sencor or Lexone as either a band or broadcast spray during planting or as a separate operation after planting, but before the soybeans emerge. Do not spray Sencor or Lexone over the top of emerged soybeans or injury may result.

Use Directions—Follow directions on the TRIFLURALIN 4EC Sencor or Lexone labels for specific instructions regarding each chemical.

Special Precautions—Do not use Lexone or Sencor on Tracy, Semmes, Altona, Vansly or Coker 102 soybeans as these varieties are sensitive to Lexone or Sencor and injury to the crop may result.

Do not use treated vines for feed or forage.

Seed must be planted at least 1½ inches below the soil surface but not more than 2 inches before a Sencor or Lexone application.

Do not apply Sencor or Lexone more than once per season.

Do not replant areas treated with Sencor or Lexone to any crop other than soybeans within 4 months after treatment.

Injury to soybeans may occur if Lexone or Sencor is used on soils having a calcareous surface or pH of 7.5 or higher, or if used in conjunction with soil applied organic phosphate pesticides.

Caution: Read the TRIFLURALIN 4EC Sencor or Lexone labels carefully before using. Note all warnings, precautions and special precautions.

Broadcast Applications Rates	TRIFLURALIN 4EC	SENCOR 50 WP Post-Plant/Pre-Emergence		
		Less than 2% Organic Matter	2 to 4% Organic Matter	Over 4% Organic Matter
Soil Texture*				
Coarse**	1 pt	DO NOT USE	¾ lb	1 lb
Medium	1½ pts	¾ to 1 lb	1 to 1½ lbs	1½ to 2 lbs
Fine	2 pts	1 to 1½ lbs	1½ to 2 lbs	2 lbs
Mississippi Delta	Rate according to soil texture	1½ lbs	1¾ lbs	2 lbs

**Do not apply Sencor to sandy soils or to coarse soils (sandy loam loamy sand) containing less than 2% organic matter

Broadcast Applications Rates	TRIFLURALIN 4EC	LEXONE 50 WP Post-Plant/Pre-Emergence	
		½ to 2% Organic Matter	More than 2% Organic Matter
Soil Texture*			
Coarse**	1 pt	DO NOT USE	¾ lb
Medium	1½ pts	¾ lb	1 lb
Fine	2 pts	1 lb	1 lb
Mississippi Delta	Rate according to soil texture	1½ lbs	1½ lbs

*Do not use Lexone in sand nor on soils with less than ½% organic matter as crop injury may result.

SOYBEANS—TRIFLURALIN 4EC/Amben
 Amben may be applied in a band over the soybean row at planting time in fields where TRIFLURALIN 4EC has been applied as a preplant soil incorporated herbicide. (See Weed Control section for weeds controlled by this treatment.) Or Amben may be applied several days prior to planting as a broadcast tank-mix with TRIFLURALIN 4EC. The tank mixture should be used as a spring preplant soil incorporated treatment. The tank-mix improves broadleaf weed, control of species such as smartweed, velvetleaf, and ragweed. For broadcast treatments, incorporate chemicals immediately and thoroughly to an approximate depth of 2 inches with a disc, field cultivator or similar tool set to cut a depth of 4 to 6 inches. Apply Amben at a rate of 1 gallon (2.0 pounds acid equivalent) per broadcast acre. Apply TRIFLURALIN 4EC at a rate of 1½ pints for medium soils and 2 pints for fine soils. Do not use on muck or charcoal soils. Read and observe all directions and cautions on the Amben label.

SUGAR BEETS:
 Apply TRIFLURALIN 4EC as a broadcast, overlap spray when plants 2 and 6 inches tall. Exposed beet roots should be covered with soil before ton to reduce the possibility of girdling. Set incorporation machinery to soil toward the plants in the row. Care should be taken that incorporation does not damage the sugar beet taproot.

	Eastern US	Western US
Coarse soils	1 pt	1 pt
Medium soils	1½ pts	1½-1½ pts
Fine soils	1½ pts	1½ pts

SUGAR BEETS—Incorporation with a Tine-Tooth Harrow in the State of Colorado, Idaho, Kansas, Montana, Nebraska, Oregon, Utah, Washington and Wyoming Only:
 A properly operated tine-tooth harrow (Flexline or Melroe) can provide incorporation of TRIFLURALIN 4EC for effective weed control in sugar beets. The tine-tooth harrow 2 times over the field in opposite directions at 2 to 6 mph and set the harrow to cut 1 to 2 inches deep. Care should be taken that the tine-tooth harrow does not damage the sugar beet taproot.

SUGARCANE—Plant Cane Only: (Eastern U.S. Only)
 Apply and incorporate TRIFLURALIN 4EC twice a year at a broadcast rate of 2 to 4 pints for all soil textures. Make the application in the fall on fields immediately after the seed pieces are planted. Make the TRIFLURALIN 4EC application in the spring before or shortly after the cane emerges. Loosen beds 2 to 3 inches deep before the spring application. Care should be taken that incorporation machinery does not damage the seed pieces or emerging shoots.

SUGARCANE—Applications up to Layby for Plant Cane or Ratoon Cane in Louisiana or Texas only:
 Apply and incorporate TRIFLURALIN 4EC at a broadcast rate per acre of 4 pints for all soil textures. Make the TRIFLURALIN 4EC application in the spring or shortly after the cane emerges up to layby. Make the TRIFLURALIN 4EC application after the beds have been shaved or false shaved. Loosen rain-packed beds 2 to 3 inches deep before application. Care should be taken so that incorporation machinery does not damage seed pieces or emerging shoots. A roller or bed chopper may be used to incorporate TRIFLURALIN 4EC layby in sugarcane on all soil textures. Follow normal incorporation directions for the roller or bed chopper. Set bed chopper to cut 3 to 4 inches deep and operate at 4 to 6 mph.

SUGARCANE—Ragweed Control in Louisiana only:
 Apply and incorporate TRIFLURALIN 4EC on either plant or ratoon cane at a broadcast rate per acre of 4 pints for all soil textures. Make the TRIFLURALIN 4EC application in the spring from before or shortly after the cane emerges up to the TRIFLURALIN 4EC application after the beds have been shaved or false shaved. Loosen rain-packed beds 2 to 3 inches deep before application. Care should be taken so that incorporation machinery does not damage seed pieces or emerging shoots. A rolling cultivator or bed chopper may be used to incorporate TRIFLURALIN 4EC layby applications in sugarcane on all soil textures. Follow normal incorporation directions for the rolling cultivator. Set bed chopper to cut 3 to 4 inches deep and operate 2 times at 4 to 6 mph.

SUGARCANE—Post-Plant in Hawaii Only For control of most annual grasses, including guineagrass:

Surface apply TRIFLURALIN 4EC after planting (for plant cane) or after harvesting (for ratoon cane), before weeds and cane emerge at a broadcast rate per acre of 6 to 8 pints for all soil textures. In plant cane, the beds should be formed or rolled before application. In ratoon cane, the crop residue should be removed before application. If large amounts of crop residues are present, TRIFLURALIN 4EC will not be effective. Apply just before anticipated rainfall or sprinkle irrigate immediately after application.

SUNFLOWER:

Apply and incorporate TRIFLURALIN 4EC before planting

	Eastern US	Western US
Coarse soils	1 pt.	1 pt.
Medium soils	1½ pts.	1½-1¾ pts.
Fine soils	2 pts.	1½ pts.
Soils with 2 to 5% organic matter	1½-2 pts.	1½-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

TOMATOES:

For Direct-seeded tomatoes apply TRIFLURALIN 4EC at blocking or thinning. Apply as a directed spray to the soil between the rows and beneath the plants and incorporate.

	Eastern US	Western US
Coarse soils	1 pt.	1 pt.
Medium soils	1½ pts.	1½-1¾ pts.
Fine soils	2 pts.	1½ pts.
Soils with 2 to 5% organic matter	1½ pts.	1½-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

For Transplant tomatoes apply and incorporate TRIFLURALIN 4EC before transplanting. Do not apply after transplanting.

	Eastern U.S.	Western U.S.
Coarse soils	—	1 pt.
Medium soils	—	1½-1¾ pts.
Fine soils	—	1½ pts.
Soils with 2 to 5% organic matter	—	1½-2 pts.
Soils with 5.1 to 10% organic matter	—	2 pts.

TREES AND VINEYARDS: (Eastern U.S. only)

For New Plantings of Vineyards, Citrus and Pecan Trees apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse soils; 1½ pints on medium soils; 2 pints on fine soils; 1½ pints on fine soils with 2 to 5% organic matter; and 2 pints on soils with 5.1 to 10% organic matter.

For Non-Bearing Established Plantings of Citrus and Pecan Trees and Bearing Plantings of Grapefruit, Lemon, Orange, Pecan, Tangelo, Tangerine Trees apply TRIFLURALIN 4EC at a broadcast rate per acre of 2 to 4 pints for all soil textures. In these established plantings, apply as a directed spray to the soil around the trees and use incorporation methods not injurious to the trees.

NOTE: If crops are planted between the trees, label directions for those species apply to the area which is interplanted. For continued weed control in citrus apply TRIFLURALIN 4EC 2 times a year at an interval of approximately 4 to 6 months.

TREES AND VINEYARDS: (Western U.S. only)

For New Plantings of Almond, Apricot, Citrus, Nectarine, Peach, Pecan and Walnut Trees apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 pint on coarse soils; 1½ to 1¾ pints on medium soils; 1½ to 2 pints on soils with 2 to 5% organic matter, and 2 pints on soils with 5.1 to 10% organic matter.

For New Plantings of Vineyards apply and incorporate TRIFLURALIN 4EC before planting at a broadcast rate per acre of 1 to 1½ pints on coarse soils; 1½ to 2 pints on medium soils and 3 to 4 pints on fine soils or soils with 2 to 10% organic matter. Do not use more than 2 pints per acre on heat-treated vines.

For Post-Plant Applications on Bearing or Non-Bearing Established Plantings of Almond, Apricot, Grapefruit, Lemon, Nectarine, Orange, Peach, Plum, Prune, Tangelo, Tangerine and Walnut Trees apply TRIFLURALIN 4EC at a broadcast rate per acre of 2 to 4 pints for all soil textures. In these established plantings, apply as a directed spray to the soil around the trees or vines and use incorporation methods not injurious to the trees or vines. Do not apply to vineyards 60 days of harvest.

NOTE: If crops are planted between the trees or vines, label directions for those species apply to the area which is interplanted. For continued weed control in citrus trees, apply TRIFLURALIN 4EC 2 times a year at an interval of approximately 4 to 6 months.

TREES AND VINEYARDS—Rhizome Johnsongrass Control: (Western U.S. only) Commercially acceptable control of rhizome Johnsongrass can be obtained in plant applications in Bearing and Non-Bearing established plantings of Almond, Apricot, Grapefruit, Lemon, Nectarine, Orange, Peach, Pecan, Tangerines and Walnut trees with a TRIFLURALIN 4EC program when applied 2 years in a row.

Soil Preparation—Work the soil thoroughly to bring the rhizomes nearer the surface.

Application—Apply TRIFLURALIN 4EC at a broadcast rate per acre of 4 to 6 pints on coarse soils; 2 to 4 pints on medium soils; and 2 to 4 pints on fine soils. Apply 2 times each year for 2 years in a row. Do not apply to vineyards 60 days of harvest.

Incorporation—Incorporate TRIFLURALIN 4EC thoroughly with a disk or harrow cut 4 to 6 inches deep and operate 2 times at 4 to 6 mph.

Cultivation—Some Johnsongrass plants will escape. Timely cultivation is necessary to obtain commercially acceptable control. Commercially acceptable control will not be obtained with only 1 year of TRIFLURALIN 4EC use.

Precautions: Do not use the 2-quart rate on new plantings as injury may result. Do not interplant orchards or vineyards with other crops. If TRIFLURALIN 4EC is used in vineyards and orchards are diverted to other crop uses, plant only those species which TRIFLURALIN 4EC has been registered as a pre-plant treatment.

10008

TREES AND VINEYARDS—Field Bindweed Control in Vineyards, Almond, Apricot, Grapefruit, Lemon, Nectarine, Orange, Peach, Pecan, Tangelo, Tangerine and Walnut Trees in California only:

For the control of field bindweed in the state of California, apply TRIFLURALIN 4EC at a broadcast rate of 4 pints per acre on all soil textures. TRIFLURALIN 4EC must be applied in the spring with a specially designed spray blade which applies a thin concentrated layer at a soil depth of 4 to 6 inches. The layer of TRIFLURALIN 4EC prevents bindweed shoots from emerging.

Land Preparation—Destroy all weeds and grasses with soil tillage before applying TRIFLURALIN 4EC. This tillage is necessary to prevent trash from interfering with the operation of the spray blade.

Equipment—This operation requires a spray blade capable of running 4 to 6 inches below the surface of the soil. The spray blade should be equipped with nozzles located under the blade and directed so that the TRIFLURALIN 4EC spray will be trapped under the soil which is flowing over the blade as it is pulled through the soil. Use a sufficient number of nozzles with spacing to completely and uniformly apply TRIFLURALIN 4EC underground in a thin horizontal layer.

Application—Apply TRIFLURALIN 4EC in 40 to 80 gallons of water per acre. Operate the spray blade at a depth of 4 to 6 inches.

Precaution: Some soils develop cracks as they dry after rainfall or irrigation. Field bindweed may emerge if the cracks extend through the TRIFLURALIN 4EC layer. Prevent or eliminate cracks by shallow discing or other tillage. Avoid deep tillage which disturbs the subsurface layer. Cultivation or tillage also aids the control of germinating seeds.

WHEAT(WINTER)—TRIFLURALIN 4EC for preplant pre-emergence control of cheatgrass and other weeds in winter wheat grown in Washington, Oregon, Idaho, and Montana:

When applied as directed, TRIFLURALIN 4EC will provide effective pre-emergence control of cheatgrass and a number of other annual grasses and broadleaf weeds controlled by TRIFLURALIN 4EC in winter wheat grown in Washington, Oregon, Idaho, and Montana. The growth, development and yield of winter wheat will not be adversely affected, provided the seed is placed below the zone of soil treated with TRIFLURALIN 4EC.

Broadcast Rates Per Acre:

Apply TRIFLURALIN 4EC any time during a period from 3 weeks up to immediately prior to planting. Broadcast TRIFLURALIN 4EC at the following rates per acre according to soil texture.

Soil Texture	TRIFLURALIN 4EC
Coarse	1½ pts.
Medium	1½ pts.
Fine	2 pts.

Incorporation Directions—Shallowly incorporate TRIFLURALIN 4EC into the soil with a flexible tine-tooth harrow (Flexline, Metroe) set to cut 1 to 2 inches deep. Operate the equipment in 2 different directions at a speed 3 to 6 miles per hour. The first incorporation must be within 24 hours after application. The second incorporation may be done at any time but before planting. Do not till the soil with a disc after the TRIFLURALIN 4EC has been applied and incorporated with a flexible tine harrow.

Seeding Directions—Use only a deep furrow or semi-deep furrow drill that will be sure to place the seed below the zone of soil into which TRIFLURALIN 4EC has been incorporated.

Wheat planted in direct contact with TRIFLURALIN 4EC treated soil may suffer crop injury in the form of delay in emergence and development.

WHEAT (WINTER)—Fallow-soil application of TRIFLURALIN 4EC for weed control in winter wheat grown in Washington and Oregon:

Uniformly applied TRIFLURALIN 4EC at the recommended rate and shallowly incorporated into fallow soil as much as four months ahead of planting time, effectively control cheatgrass and certain annual grasses and broadleaf weeds in winter wheat grown in Washington and Oregon. The growth, development, or yield of winter wheat will not be adversely affected, provided the seed is placed below the zone of soil treated with TRIFLURALIN 4EC with deep or semi-deep furrow-drill.

Broadcast directions and application rates per acre:

Soil Texture	TRIFLURALIN 4EC
Coarse	1½ pts.
Medium	1½ pts.
Fine	2 pts.

Apply any time from May to September prior to the fall planting of winter wheat.

Incorporation—Shallowly incorporate TRIFLURALIN 4EC into the soil with a flexible tine-tooth harrow (also called Flexline or Metroe) set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Thorough incorporation requires two passes of the equipment in different directions over the field. The first pass must be made within 24 hours after application. The second pass may be delayed for several weeks but should be made before seeding. Do not till the soil with a disc after TRIFLURALIN 4EC has been applied with a flexible tine harrow.

Precaution: Use only deep furrow or semi-deep furrow drills. Place seed below the zone of soil into which TRIFLURALIN 4EC has been incorporated. Do not plant wheat directly into the zone of soil treated with TRIFLURALIN 4EC as injury to the crop or a delay in its emergence and development may occur.

NOTICE OF WARRANTY

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of the product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

- * SENCOR is a Reg. T.M. of Parent Co. of Farbenfabriken Bayer GmbH, Leverkusen.
- * KARMEK and LEXONC are Reg. T.M.'s of E.I. duPont de Nemours and Co., Inc.
- * EPTAM is a Reg. T.M. of Stauffer Chemical Co.
- * AMIBEN is a Reg. T.M. of Union Carbide Corp.
- * COTDRAN and CAPAROL are Reg. T.M.'s of Ciba-Geigy Corp.
- * AVADEX is a Reg. T.M. of Monsanto Co.