

34704-259

08/12/2013

1/5

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

AUG 12 2013

Premjit Halarnkar
Registration Manager
Loveland Products
P.O. Box 1286
Greeley, CO 80632-1286

Subject: Submission of amended labeling to add EPA initiated interim plant back restrictions as outlined in 1/16/13 letter EPA Reg. No. 34704-259/Phorate 20 G Submissions dated February 14, 2013

Dear Mr. Halarnkar:

The proposed labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you submit two copies of your final printed label before you release the product for shipment. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

A stamped copy is enclosed for your records. If you have any questions, please contact Marianne Lewis at (703) 308-8043 or lewis.marianne@epa.gov.

Regards,

A handwritten signature in black ink that reads "Venus Eagle". The signature is fluid and cursive, with a long horizontal line extending to the right.

Venus Eagle, PM 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

2/5

RESTRICTED USE PESTICIDE

Due to acute oral, dermal and inhalation toxicity and avian hazards

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certificate.



PHORATE 20 G

Organophosphate Insecticide

ACTIVE INGREDIENT:

Phorate: O,O-diethyl S-[(ethylthio) methyl] phosphorodithioate 20.0%

INERT INGREDIENTS: 80.0%
TOTAL 100.0%

EPA REG. NO. 34704-259

EPA EST. NO. _____

NET WEIGHT 50 LB (22.68 KG)

EXP 02/13

KEEP OUT OF REACH OF CHILDREN

**DANGER
POISON**



PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See Below For Antidote and Other Precautions.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. Call a poison control center or doctor for treatment advice. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-944-8565.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING

NOTE TO PHYSICIANS: Warning symptoms including weakness, headache, tightness of chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers.

Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary.

Probable mucosal damage may contraindicate the use of gastric lavage.

Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Fatal if swallowed, inhaled or absorbed through the skin. Corrosive. Causes irreversible eye damage. DO NOT breathe dust. DO NOT get in eyes, on skin, on clothing. Rapidly absorbed through skin. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to poisoning.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are those made of any water-proof material. If you want more options, follow the instructions for category A on an EPA chemical resistant category selection chart.

Loaders, applicators and other handlers using engineering controls must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks.

In addition to the PPE listed above, loaders must wear

- Chemical resistant gloves.

All other handlers performing tasks such as spill clean up or equipment cleaning for which engineering controls are not feasible must wear:

- Coveralls over long sleeved shirt and long pants,
- Chemical resistant footwear plus socks,
- Chemical resistant gloves,
- Protective eyewear,
- NIOSH-approved dust/mist filtering respirator with MSHA/NIOSH approval number prefix TC-21c or a NIOSH approved respirator with any N,R,P, or HE filter,
- Chemical resistant apron if exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

See Engineering Controls for additional requirements.

Engineering controls statements:

This product is formulated into an EZ Load System that meets the definition of a closed loading system as defined by the Worker Protection Standard for Agricultural Pesticides 40 CFR 170.240 (d)(4). In addition to wearing the required PPE for handlers using engineering controls, loaders must be provided, must have immediately available, and must wear in case of a broken package, spill or equipment breakdown: chemical-resistant apron, chemical-resistant footwear, and a NIOSH-approved dust/mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH approved respirator with any N, R, P, or HE filter.

Applicators must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides for dermal protection. In addition the applicator: must wear the PPE required for handlers using engineering controls, and must either use an enclosed cab that provides equivalent respiratory protection to a dust/mist filtering respirator or wear a NIOSH-approved dust/mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH approved respirator with any N, R, P, or HE filter, must be provided, must have immediately available for use, and must wear if they must exit the cab in the treated area: coveralls, chemical-resistant gloves, chemical-resistant footwear, protective eyewear, and if using an enclosed cab that provides respiratory protection NIOSH-approved dust/mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH approved respirator with an N, R, P, or HE filter. Applicators using enclosed cabs shall:

- take off any PPE that was worn in the treated area before reentering the cab, and
- store all such PPE in a chemical resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under:

EPA. Reg. No. 34704-259

8/12/13

PHORATE 20 G
EPA REG. NO. 34704-259

3/5

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

IMPORTANT! Failure to follow precautions including wearing proper Personal Protective Equipment (PPE) may result in serious or even life threatening poisoning requiring immediate medical attention. The active ingredient of this granular formulation can be absorbed across the skin to cause poisoning.

WORK SAFETY RULES

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposure should be allowed until it has been determined by means of blood tests that the cholinesterase level has returned to normal. Before using this product consult the National Pesticide Information Center for recommendations regarding such blood tests, poisoning management, emergency treatment, and other information regarding the toxicity of phorate. The toll free number for the National Pesticide Information Center is 1-800-858-7378.

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See "FIRST AID" section.

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain (0.6 milligram) tablets for emergency use. If symptoms include blurred vision, stomach cramps or tightness in chest, don't wait for a physician but take two tablets at once. Do not take atropine unless symptoms of poisoning have occurred. See a physician as soon as possible.

ENVIRONMENTAL HAZARDS

This pesticide is very highly toxic to fish and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wastewater or rinseate. Birds and mammals may be killed if granules are not properly covered with soil in all areas of the treated field and in loading areas.

You are advised that use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, COI- 0132C, (W.D. WA). For further information, please refer to the following EPA website: <http://www.epa.gov/espp/wtc/uselimination.htm>.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

IMPORTANT! Read precautionary statements before using this product. Do not use this product on Long Island, NY.

Aerial application is prohibited.

PHORATE 20 G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage.

Incorporation into soil is required.

Only one application per crop per season may be made.

Use on peanuts at pegging is prohibited.

Application of phorate in saturated soils is prohibited. Do not treat while precipitation is occurring or while conditions favor runoff from the treated area.

Do not apply in wet soil conditions that may prevent the equipment from covering pesticide granules.

Under some conditions phorate may have a high potential for runoff into surface water for several days post application. Do not apply in the following situations:

Frequently flooded areas

Areas where intense or sustained rainfall is forecasted to occur within 48 hours.

In the event of a crop failure, any crop on this label can be replanted immediately. All other crops must not be planted in treated fields for 9 months after application.

Use Best Management Practices for minimizing surface runoff in the following areas:

Poorly draining or wet soils with readily visible slopes toward adjacent surface water

Areas over-laying extremely shallow ground water

Areas with in-field canals or ditches that drain to surface water

Areas not separated from adjacent surface waters with vegetated filter strips

Areas over-laying tile drainage systems that drain to surface water

When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local soil conservation service for recommendations in your use area.

In particular, where highly erodible land (HEL) is adjacent to aquatic bodies, a 66 foot buffer/setback area should be left in grass or other natural vegetation.

Do not apply within 50 feet of any drinking water well to minimize potential contamination.

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of ground water contamination.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Each 48 hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.

Exception: if the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls worn over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Chemical-resistant footwear plus socks and protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

TYPICAL APPLICATION RATES FOR VARIOUS ROW SPACINGS

	Rate oz./1,000 Ft. of Row									
	1.6 oz.	2 oz.	4 oz.	6 oz.	8 oz.	10 oz.	12 oz.	14 oz.	16 oz.	17.2 oz.
40"				4.9	6.5	8.2	9.8	11.4	13.1	14.1
38"				5.2	6.9	8.6	10.3	12.0	13.7	14.7
36"				5.4	7.3	9.1	10.9	12.7	14.5	15.6
34"				5.6	7.7	9.7	11.6	13.5	15.4	16.6
32"				6.1	8.2	10.2	12.3	14.3	16.3	17.5
30"				6.5	8.7	10.9	13.1	15.2	17.4	18.7
26"				7.1	9.5	11.8	14.2	16.6	18.9	
24"			5.4	8.2	10.9	13.6	16.3	19.1		
22"			5.9	8.9	11.9	14.9	17.9			
20"			6.5	9.8	13.1	16.3				
16"			8.2	12.3	16.3					
12"		5.4	10.9	16.3						
8"	6.5	8.2	16.3							
6"		10.9								

EQUIVALENT POUNDS PER ACRE

Row Spacing

4/5

PHORATE 20 G
EPA REG. NO. 34704-259

DIRECTIONS FOR USE

Crop	Pests Controlled	Rate PHORATE 20 G	Application	Remarks
BEANS Soiled drilled at planting	Mexican Bean Beetles	4 1/2 to 9 3/8 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 10 lb/A	Distribute the granules evenly in the row to the side of the seed at planting time.	PHI = 60 days The grazing or feeding of treated hay or forage to livestock is prohibited.
	Soil drilled or banded at planting	Leafhoppers, Aphids, Lygus Bugs, Thrips, Mites and Seedcorn Maggots	4 1/2 to 7 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 7.5 lb/A	Do not place PHORATE 20 G granules in direct contact with the seed. Do not make more than one application per crop per season. Granules must be incorporated into the soil.
FIELD CORN: Soil banded incorporated at planting	Corn Rootworms, Mites, Seed Corn Beetle (Iowa and Illinois Only), Reduction of Wireworms (North Central and South eastern States only)	6 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 6.5 lb/A	Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel. Do not place PHORATE 20 G granules in direct contact with seed.	PHI = N/A In-furrow application is prohibited. Do not make more than one application per crop per season. Granules must be incorporated into the soil.
	Soil banded at cultivation	Corn Rootworms Chinch bug nymphs	Apply granules at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply in June as soon as Rootworms activity is observed.	PHI = 30 days Application after cultivation treatment is prohibited. Cultivation application may be made to control corn rootworms or chinch bug nymphs. When made to control chinch bug nymphs, application may only be made in AR, CO, KS, LA, MS, NE, OK, TN and TX and a 30-day PHI/PGI has been established. Do not make more than one application per crop per season. Granules must be incorporated into the soil. Do not graze treated fields for 30 days after application.
SWEET CORN: Soil banded incorporated at planting	Corn Rootworms, Mites, Seed Corn Beetle (Iowa & Illinois only), Reduction of Wireworms (North Central and Southeastern States only)	6 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 6.5 lb/A	Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel. Do not place PHORATE 20 G granules in direct contact with seed.	PHI = N/A In-furrow application is prohibited. Do not make more than one application per crop per season.
	Soil banded at cultivation	Corn Rootworms	Apply granules at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply in June as soon as rootworm activity is observed.	PHI = 30 days Application after cultivation treatment is prohibited. Do not make more than one application per crop per season. Granules must be incorporated into the soil. Do not graze treated fields for 30 days after application.
COTTON: In furrow at planting	Early season Thrips, Aphid, Mites & Leaf Miners	3 ozs. per 1000 ft. of row (minimum 30-inch row spacing or no more than 3.3 lbs./A)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tillth should be favorable for good germination and emergence.	PHI = N/A Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 2 lbs. PHORATE 20 G per acre for hill-dropped cotton. The grazing or feeding of treated hay or forage to livestock is prohibited. Do not make more than one application per crop per season.
		Mites, Thrips, Aphids, Black Cutworms, Leafhoppers, White Flies and Leaf Miners.	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tillth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	PHI = N/A Recommended for areas requiring an extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soil. The grazing or feeding of treated hay or forage to livestock is prohibited. Do not make more than one application per crop per season.
COTTON: Soil incorporated at side-dressing (Irrigated cotton only) (Arizona and California only)	Mites from spring to early summer	9 to 12 ozs. per 1000 ft. of row (minimum 36-inch row spacing or no more than 10.9 lbs./A.) 9 - 10 oz per 1000 ft. of row for 30-inch row spacing or no more than 10.9 lbs./A.	Incorporate granules into soil as a side dress application deep enough to avoid disturbance by future cultivations.	PHI = 60 days Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest. The grazing or feeding of treated hay or forage to livestock is prohibited. Do not make more than one application per crop per season. Granules must be incorporated into the soil.
PEANUTS: In furrow at planting	Thrips, Leafhoppers	5 1/8 ozs. per 1000 ft. of row (minimum 24-inch row spacing) or 6.9 lb/A	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	PHI = 90 days The grazing or feeding of treated hay or forage to livestock is prohibited. Do not make more than one application per crop per season.
POTATOES: In furrow or soil banded at planting	Aphids, Leafhoppers, Leaf Miners, Psyllids, Flea Beetle Larvae and reduction of Flea Beetle Adults.	LIGHT OR SANDY SOILS: 11 1/4 ozs. per 1000 ft. of row (minimum 32-inch row spacing) or 11.25 lb/A HEAVY OR CLAY SOILS: 17 1/4 ozs. per 1000 ft. of row (minimum 32-inch row spacing or 17.25 lb/A	Distribute the granules evenly in the furrow or granules may be banded on each side of the row at planting time. Incorporation into soil is required.	PHI = 90 days Use for Colorado potato beetle control in the Del Marva Peninsula is prohibited. Do not make more than one application per crop per season. Granules must be incorporated into the soil.
		Colorado Potato Beetle (early season control)	11 1/4 ozs. per 1000 ft. of row (minimum 32-inch row spacing) or 11.25 lb/A.	
SORGHUM: Soil drilled or banded at planting	Aphids (Greenbugs)	6 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 6.5 lb/A	Place granules 2 inches to side and 1 inch below the seed at planting or as a 7-inch band over the row just prior to press wheel. Granules must be incorporated into the soil.	PHI = N/A Do not place PHORATE 20 G granules in direct contact with seed. Do not make more than one application per crop per season.
SOYBEANS: Soil drilled or banded at planting	For the early season control of Mexican bean beetle and leafhoppers	9 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 10 lb/A	Distribute the granules evenly in the row to the side of the seed at planting time.	PHI = N/A Do not place granules in direct contact with the seed. The feeding of treated foliage to livestock is prohibited. Crop injury may result when preplant incorporated or preemergence applications of metribuzin herbicides or other soil-applied organophosphate pesticides are used in conjunction with this product.
		Seedcorn maggots	9 ozs. per 1000 ft. of row (minimum 30-inch row spacing) or 10 lb/A	Do not make more than one application per crop per season. Granules must be incorporated into the soil.

5/5

PHORATE 20 G
EPA REG. NO. 34704-259

Pests Crop	Rate Controlled	PHORATE 20 G	Application	Remarks
SUGAR BEETS: Soil drilled or banded at planting	Aphids, Beet Root Maggots, Leafhoppers, Mites, Leaf Miners	4 ½ ozs. per 1000 ft. of row (minimum 20-inch row spacing) or 7.5 lb/A		PHI = 30 days Do not place PHORATE 20 G granules in direct contact with the seed. Do not feed sugar beet tops or silage to dairy cattle. Consult your state experiment station or state extension service for proper timing of application in your area. Do not make more than one application per crop per season. Granules must be incorporated into the soil.
SUGAR CANE: (Florida)	Wireworms	19.5 lb/A	Apply in a 10 to 12 inch band directly on the seed piece and surrounding soil in the open furrow immediately before covering. Cover granules that may be exposed on the end of treated rows, turns and loading areas by deep disking after treating fields. All granules must be incorporated into the soil.	PHI = N/A Use limited to FL. The grazing or feeding of treated forage or hay to livestock is prohibited. Do not make more than one application per crop per season. Granules must be incorporated into the soil.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Do not store opened containers. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength.

SPILLED MATERIAL, block to prevent spreading of spill. Scoop and sweep into a disposable container, whether it occurs indoors, or in the field.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty box into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Offer for recycling if available or dispose of empty box in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances. Do not place hands inside or near the open bottom of the box. **For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY
BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. Box 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR
LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286