

Page 30

8/15

~~300-262704~~  
17/5

Mr. Michael Jackson  
United Agri Products  
410 18th Street  
P.O. Box 667  
Greeley, Colorado 80632

Dear Mr. Jackson:

Subject: Tetrachlorvinphos Registration Standard  
Clean Crop RL-2 Rabon Livestock Dust  
EPA Registration No. 34704-266  
Your Submission Dated March 26, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely,

George T. LaRocca  
Product Manager 15  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

Helbert 4-24-90

APR 30 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.



# RL-2 RABON<sup>®</sup>

## LIVESTOCK DUST

### A READY-TO-USE INSECTICIDE DUST

To control HORN FLIES and LICE and aid  
in the control of FACE FLIES on  
BEEF CATTLE, DAIRY CATTLE and to  
control LICE on SWINE.

ACTIVE INGREDIENT:	By Wt.
Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl)	
vinyl dimethyl phosphate	3.00%
INERT INGREDIENTS:	97.00%
TOTAL	100.00%

may be disposed of on site or at an approved waste disposal facility  
**CONTAINER DISPOSAL:** Completely empty bag into application  
equipment. Then, dispose of empty bag in a sanitary landfill or by  
incineration, or, if allowed by State and local authorities, by burning. If  
burned, stay out of smoke.

### APPLICATIONS

This product is a ready-to-use insecticide dust which can be applied by  
hand or through the use of self-treating dust bags. Do not apply with bare  
hands. **Wear long-sleeved shirt and pants; chemical resistant gloves,  
shoes and socks.** There is no withholding period from last application to  
slaughter.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements

EPA REG. NO. 34704-266

EPA EST. NO. 10543-MO-1

NET WEIGHT 12 1/2 POUNDS

23775

EXP-1Y

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Harmful if swallowed. Do not breathe dust. Do not get in eyes, on skin, or  
on clothing. Wash thoroughly with soap and water after handling and  
before eating or smoking.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff may be hazardous to aquatic  
organisms in adjacent areas. Do not apply directly to water or wetlands  
(swamps, bogs, marshes and potholes). Do not contaminate water when  
disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on  
blooming crops or weeds. Do not apply this product or allow it to drift to  
blooming crops or weeds if bees are visiting the treatment area.

### STATEMENT OF PRACTICAL TREATMENT

If swallowed—Call a physician or Poison Control Center immediately.  
Drink one or two glasses of water and induce vomiting by touching the  
back of throat with a finger. Do not induce vomiting or give anything by  
mouth to an unconscious or convulsing person.

If inhaled—Remove victim to fresh air. Apply artificial respiration if indi-  
cated.

If on skin—Remove contaminated clothing and wash affected areas  
with soap and water.

If in eyes—Flush eyes with plenty of water. Call a physician if irritation  
persists.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent  
with its labeling. Take Time. Observe All Label Directions.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or  
disposal.

**STORAGE:** Store in a cool, dry place in original container.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product

### ANIMAL INSECT REMARKS

ANIMAL	INSECT	REMARKS
Beef Cattle	Horn Flies	Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.
Dairy Cattle	Face Flies	
Cattle	Lice	

### HAND DUSTING

### SELF-TREATING DUST BAG

**FORCED USE—**Put dust in cotton cloth or double  
burlap bags or use pre-packed weather-proof cattle  
dust bags and hang in barn door exits or alleyways  
leading from animal buildings, salt or mineral  
blocks or watering holes. Protect cloth or burlap  
bags from weather.

**FREE CHOICE USE—**Use the same dust bags as  
above but place in loafing sheds, holding pens,  
feedlots, near watering holes or other areas where  
cattle gather. The free choice aids in the control of  
lice.

Swine	Lice	<b>HAND DUSTING—</b> Apply 3-4 oz. of dust by conven- tional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.
-------	------	--

### NOTICE OF WARRANTY

Platte warrants that this product conforms to the chemical description on  
the label thereof and is reasonably fit for the purposes stated on such  
label only when used in accordance with the directions under normal use  
conditions. It is impossible to eliminate all risks inherently associated  
with the use of the product. Crop injury, ineffectiveness, or other un-  
intended consequences may result because of such factors as weather  
conditions, presence of other materials, or the manner of use or applica-  
tion, all of which are beyond the control of Platte. In no case shall Platte  
be liable for consequential, special or indirect damages resulting from  
the use or handling of this product. All such risks shall be assumed by the  
buyer. Platte makes no warranties of Merchantability or fitness for a  
particular purpose nor any other express or implied warranty except as  
stated above.

U.S. Patent No. 3777716

FORMULATED FOR

PLATTE CHEMICAL COMPANY, INC.

150 SO. MAIN STREET

FREMONT, NEBRASKA 68025

**BEST AVAILABLE COPY**