


34704-276

02/09/2004

1/2



LIVESTOCK 1%

RABON® DUST

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under 34704-276

A READY-TO-USE INSECTICIDE DUST

Controls **HORN FLIES** and **LICE** and aid in the control of **FACE FLIES** on **BEEF** and **DAIRY CATTLE**.

Controls **LICE** on or around **SWINE**.

Controls **CHICKEN MITES, LICE** and **NORTHERN FOWL MITES** on or around **POULTRY**.

Controls certain **FLIES, LICE, TICKS** and **GNATS** on or around **HORSES**.

Control **TICKS** and **FLEAS** on or around **DOGS** and **CATS**.

ACTIVE INGREDIENT:

Tetrachlorvinphos: (z)-2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	1%
INERT INGREDIENTS	99%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements

EPA REG. NO. 34704-276

EPA EST. NO. 34704-NB-1

NET WEIGHT _____ LBS. (_____ KG)

IHT

U.S. Patent No. 3777716

EXP01Y02

FIRST AID

If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 -20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Call a poison control center or doctor for treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.

NOTE TO PHYSICIAN VETERINARIANS: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE):

Handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves (such as Barrier Laminate, Nitrile, Butyl and/or Neoprene), shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff may be hazardous to aquatic organisms in adjacent areas. For terrestrial uses, do not apply directly to water, or to where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE & DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place in original container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE.

APPLICATIONS

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

No interval is required between treatment and slaughter or between treatment and use of milk for human consumption.

Consult a veterinarian before using this product on debilitated, medicated, aged, pregnant or nursing animals.

LIVESTOCK 1% RABON® DUST
EPA REG. NO. 34704-276

2/2

ANIMAL	INSECT	APPLICATION DIRECTIONS
Beef Cattle & Dairy Cattle	Horn Flies, Face Flies & Lice	<p>Self-Treating Dust Bags: Place this product in cotton cloth or double burlap bag or any commercially available weather-proof Cattle dust bag. Protect cloth or burlap bags from weather.</p> <p>Hang the bag so the bottom of the bag will hang four to six inches below the topline of the cattle. For reduction of face flies, the bags must be located so the cattle will use them daily and must be hung to insure the faces of the cattle will be dusted.</p> <p>Regularly inspect dust bags for proper functioning and adequate dust supply. Repair or replace damaged or malfunctioning bags promptly. Recharge with dust as necessary.</p> <p>Forced Use: Suspend bags in gateways or lanes through which the animals pass daily for water, feed, or minerals. Bags may also be suspended in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in milking barn exits.</p> <p>Free Choice: Suspend bags in loafing sheds, holding pens, feedlots, near watering holes, or other areas where cattle gather daily.</p> <p>Hand Dusting: Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the tailhead, back, neck and poll, and to the face. Rub in lightly to carry the dust beneath the hair, but do not apply with bare hands. Use dust mitt or other protective glove. Repeat as necessary.</p>

ANIMAL	INSECT	HAND DUSTING
Swine	Lice	Apply 9-12 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestation, both individual animals and bedding may be treated. Three lbs. of dust should be applied per 150 sq. ft. of bedding. Do not apply with bare hands.

POULTRY

CHICKEN MITES, LICE, NORTHERN FOWL MITES

Type Housing	Applications	Remarks
Wire cages	1 lb./100 birds Plunger, rotary type duster or shaker can	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days. Do not apply with bare hands. Wire rungs and corners should also be duster treated.

NOTE: For Northern Fowl Mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.

Floor

Management	Application	Remarks
Litter	3 lbs./100 sq. ft. Plunger or rotary type duster	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	6 lbs./100 birds	Use box about 2 ft. by 2 ft. by 1 ft. deep.
Roost Paint	3 lbs./100 ft.- make thick slurry by mixing 1 lb. of dust with 1 pint of water - continually stir.	Treat roosts thoroughly, particularly the cracks and crevices.

ANIMAL	INSECT	HAND DUSTING
Horses	Face flies, Stable flies, House flies, Horn flies, Horse flies, Deer flies, Lice, Ticks, Gnats and Fly Larvae	Apply approximately 6 oz. by shaker can, hand duster, grooming brush or dust mitt. Cover upper portions of back and neck and to the face, mane and tail for added control of face flies; rub or brush in lightly to carry dust beneath the hair. Do not apply with bare hands. Repeat as necessary. For premise dusting, apply to barn or stall area floors where manure accumulates, to control fly larvae and flies in the stable area.
Horse Premises		
Dogs, Cats	Ticks and Fleas	Dust powder liberally over the animal and rub thoroughly through the hair coat to skin. Apply also to feet, between the toes and on the legs for ticks. Fleas will die quickly and can be combed from hair in a few hours if desired. Ticks are somewhat more difficult to kill and can be combed out of pet's hair a day or two after application. Do not apply to kittens and puppies under 12 weeks old. Do apply with bare hands. Occasional dusting in and around sleeping quarters and other areas will help free area of ticks and fleas. Dust bedding as well.
Kennels, Dog Houses	Ticks and Fleas	

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.

419 18TH STREET, GREELEY, COLORADO 80631-5852