

Reg no 34704-304

PM-16

DIRECTIONS FOR USE

ALFALFA AND CLOWERS: Alfalfa weevil, aphids, grasshoppers, potato leafhopper, and spittlebugs, apply 1 1/2 to 2 pints of malathion per acre in sufficient water to give good coverage. Make through coverage application when larvae is small. Repeat as necessary. When crop is in bloom, apply malathion only in the evening or early morning, when bees are not working in the field or are not hanging on the outside of the hives. Repeat application as necessary. Do not apply within 7 days of harvest or pasturing.

TOBACCO: Aphids, spray at the rate of 1 1/2 pints per acre in sufficient water to make full coverage. Repeat as necessary. Do not apply within 7 days of harvest.

CORN AND GRAIN SORGHUMS: Aphids, apply at the rate of 1 1/2 pints per acre in sufficient water for good coverage. Make full coverage application and repeat as necessary.

SMALL GRAINS: Grain aphids and green bugs on barley, wheat, rye and oats, apply at the rate of 1 1/2 pints per acre in sufficient water for good coverage. Do not apply within 7 days of harvest.

LIVESTOCK AND POULTRY

BEEF CATTLE: Lice, ticks, hornflies, apply a complete coverage spray at the rate of 1 to 1 1/2 gallon per 100 gallons of water. For lice and ticks repeat application at two week intervals if needed. For hornflies repeat at two week intervals, during fly season. Do not treat calves under one month of age.

SWINE: For lice and sarcoptic mange, apply at the rate of 1 gallon in 100 gallons of water. Thoroughly spray all animals, bedding and walls. Repeat if needed. Avoid contamination of feed, feed containers and watering troughs.

NET CONTENT 1 GALLON



E.P.A. Reg. No. 4876-51

MALATHION 50

EMULSION CONCENTRATE

For Control Of
ALFALFA WEEVIL, TOBACCO APHIDS,
SMALL GRAIN GREEN BUGS,
POULTRY and LIVESTOCK INSECTS

ACCEPTED

| | |
|-----------------------------------|-----|
| ACTIVE INGREDIENT | 50 |
| Malathion* | |
| INERT INGREDIENTS | 50 |
| TOTAL | 100 |
| * O O - Dimethyl phosphorothioate | |

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 4876-51

CAUTION: Keep out of reach of Children
See side panel for additional cautions

Manufactured By

"AG" SUPPLY COMPANY

DIRECTIONS FOR USE

POULTRY: Chickens, Geese, Ducks and Turkeys: Direct Application. (fowl mites and poultry lice apply 2 tablespoons per 1 gallon of water. Apply one gallon of mixture to 100 to 150 birds. Repeat in 4 to 8 weeks or as necessary.

CHICKEN BODY LICE and SHIRT LICE: Make a solution containing 64 oz. in 15 gallons of water (1 oz. to 2 gallons), dip three to four inches of the tail into the solution, to thoroughly wet the vent and surrounding areas. Repeat in 7 to 12 days or as needed.

POULTRY HOUSES: Spray for fowl mites, red mites, poultry lice and flies, mix 4 to 5 tablespoons per 1 gallon of water, liberally spray walls, ceiling, roosts and nests, being sure to spray into all cracks and crevices. Repeat as necessary. Avoid contaminating feed, feed troughs, and water fountains. Mixture may be brushed on as a roost paint at the rate of 1 pint per 150 feet of roost.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid getting into eyes, on skin or clothing. Avoid contamination of feed or food stuffs. If poison occurs get prompt medical aid. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of waste. Apply this product only as specified on this label.

DO NOT USE, POOP, SPILL OR STORE NEAR FEED FRAME.

Do not reuse empty container. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of smoke.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.