

34704-306 PM 15

103

~~184007~~

FEB - 5 1987

Ms. Diana G. Williams
Platte Chemical Company
419 8th Street
P.O. Box 667
Greeley, CO 80632

Dear Ms. Williams:

Subject: Amendment - Label Update and Change in Product Name
Clean Crop 'D-Louse' Animal Insecticide
EPA Registration No. 34704-306
Your Letter Dated August 5, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. Present the Active Ingredient as:

Coumaphos [O,O-Diethyl O-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) phosphorothioate].

Coumaphos is now an acceptable common name.

2. Change "Keep out of lakes, streams, and ponds" to "Do not apply directly to water."

The product name has been changed from Crown D-LOUSE to Clean Crop 'D-Louse' Animal Insecticide.

91945:Newkirk:LR-15:KENCO:1/27/87:1/31/87:sj:vo

CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

2083

2

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

P.S. In the table showing recommended uses, revise the heading "Parasites" to "Pests", or "Insects", or "Insect Pests".

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dusts. Avoid contact with eyes, skin, and clothing.

STATEMENT OF PRACTICAL TREATMENT

- If swallowed** - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to unconscious person.
- If inhaled** - Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention if victim displays signs of poisoning.
- If on skin** - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes** - Flush eyes with plenty of water. Call a physician immediately.

To Physician: Prolonged exposure will result in cholinesterase depression. Atropine sulfate is antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE:

Store in a cool, dry place.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Completely empty the container by shaking and tapping sides and bottom to loosen clinging particles, thereby dispensing the entire contents of the can only as directed by the label. Dispose of container in sanitary landfill or by incineration if allowed by state and local authorities. If burned, stay out of smoke.

IMPORTANT: Read these entire Directions and a Warranty before using this product.

Hold container at a distance from the area to be treated that will ensure even distribution of the dust.

RECOMMENDED APPLICATIONS

ANIMAL	PARASITE	REMARKS
Cattle (Dairy & Beef)	Horn Flies Lice	DIRECT APPLICATION Apply not more than 2 oz. Dust evenly into the hair on head, neck, shoulders, back. Repeat as necessary. Interval is required between treatments. Use of meat or milk as food.
Swine	Lice	DIRECT APPLICATION Apply not more than 1 oz. of a uniform coat to the skin on back. Repeat as necessary. Interval is required between treatments. Use of meat as food. BEDDING TREATMENT Apply 2 oz. of a uniform coat to square feet of fresh, dry bedding. Repeat as necessary, but not more than once every 10 days. DIRECT APPLICATION AND BEDDING TREATMENT In severe infestations both individual animal and the bedding may be treated as directed above.
Horses	Horn Flies	DIRECT APPLICATION Apply not more than 2 oz. Dust evenly into the hair on head, neck, shoulders, back. Repeat as necessary.

*NOTE: 1 oz. of this material equals approximately 100 Tablespoons.

USE RESTRICTIONS

For external insecticidal use only on above specified animals. Do not contaminate feed troughs, water, or water utensils. Provide ventilation while dusting. Use caution when spraying in areas where avoid exposure to eyes. Do not apply to sick, stressed, or convalescent animals.

WARRANTY

Platte warrants that this product conforms to the chemical composition on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of the product. Crop loss, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025

BEST AVAILABLE COPY