

CAUTION

Harmful if swallowed. If swallowed, do not induce vomiting, call a physician immediately. Avoid contact with eyes and skin. In case of skin contact, wash with soap and water. Avoid inhalation of spray mist. Avoid contamination of food and feed.

Do not contaminate water used for irrigation or domestic purposes.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use, pour, spill or store near heat or open flame.

Important: Read the entire Directions For Use and the Conditions of Sale and Warranty before using this product.

DIRECTIONS FOR USE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store below temperature of 32°F. Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATIONS

Uniformly spray one gallon of this product mixed in ten gallons of water over 1,000 sq. ft. of soil surface. Thoroughly wet all foliage to insure contact action. Best results are obtained when applications are made during or after weed emergence while growth is succulent. Where vegetation is tall and dense, use 1½ gallons of this product in ten gallons of water per 1,000 sq. ft. For control of hard-to-kill perennial weeds and grasses such as johnsongrass, bindweed, and wild carrot, use 3 to 4½ gallons per 1,000 sq. ft. Although this herbicide has considerable activity through foliar contact, much of its activity is through the roots and, therefore, adequate rainfall to move the chemical into the root zone is necessary.

Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed-free soil surface.

Care must be taken to keep this herbicide away from desirable flower beds, lawns and shrubs. Do not apply over roots of valuable trees or woody ornamental plants. Only apply herbicide outside the dripline of trees. Avoid spray drift to desirable plants.

CONDITIONS OF SALE AND WARRANTY

Since weather and other conditions under which this product may be used will vary, the Seller makes no warranty express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025



SEVIN[®]

brand Carbaryl Insecticide

4 OIL

PAI-12
EPA # 34704-313
PAI-12
LF 1/3/10

| | |
|---|-------------------|
| ACTIVE INGREDIENT: | BY WEIGHT: |
| Carbaryl (1-naphthyl methylcarbamate) | 46.10% |
| INERT INGREDIENTS: | 53.90% |
| TOTAL | 100.00% |

CONTAINS 4 POUNDS CARBARYL PER GALLON
*This product contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Below For Additional Precautionary Statements.
EPA REG. NO. 34704-313
EPA EST. NO. _____
NET CONTENTS 1 GALLON
* SEVIN is the registered trade mark of
Rhône-Poulenc Nederland B.V. for carbaryl insecticide.

25322 01Y90

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL IF SWALLOWED. Avoid breathing of spray mist. Do not take internally. Avoid contact with skin, eyes, and clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except under the forest canopy. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEE CAUTION

This product is highly toxic to bees exposed to direct treatment or residue on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Contact a physician or poison control center immediately. Do not induce vomiting. Never give anything by mouth to an unconscious or convulsing person. See "NOTE FOR PHYSICIAN," below.

If in eye: Flush eyes with plenty of water. Get medical attention if irritation persists.

If on skin: Wash thoroughly with soap and water.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase-inhibiting drugs.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), socks and shoes. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift

expose workers or other persons. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written or oral warnings must include the following information:

CAUTION

(Insert area or field description) treated with carbaryl on (insert date of application.) Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to carbaryl and describe his condition.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL INFORMATION

This product is a dispersion of finely ground technical carbaryl in a non-aromatic, low volatile oil. It is designed for air application as a low volume or ultra low volume spray. It may also be applied by ground mist blower or cold fogging, or diluted with diesel fuel, kerosene or #2 fuel oil. Do not apply this product through any type of irrigation system.

READ THIS LABEL BEFORE USE. USE IN STRICT ACCORDANCE WITH DIRECTIONS AND CAUTIONS.

PRE-HARVEST AND GRAZING USE INFORMATION AND LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when this product is used in accordance with label directions. If this product is used as directed, treated forage may be grazed or used as feed for dairy and meat animals without causing illegal residues in meat or milk. This product may be applied up to and including the day of harvest or grazing of forage crops, unless otherwise specified.

Do not use reclaimed irrigation water from crops treated with carbaryl on upland crops for which carbaryl tolerances are not established.

PLANT RESPONSE PRECAUTION

To avoid possible injury to tender foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days.

SPRAY PREPARATION

This product thickens during storage. Roll or agitate drums or recirculate product in bulk tanks before use to reduce viscosity and restore homogeneity. This product may be held prior to use and transferred in any materials of construction suitable for use with fuel oil, kerosene or diesel oil. Protect from water. Before and after use, flush nurse tanks, pumps, lines, hoses and entire spray systems with diesel fuel, kerosene, or #2 fuel oil until clean. Water, scale, rust and other residue must be removed from pumps, mixing and spray systems before use.

COMPATIBILITY

THIS PRODUCT IS NOT COMPATIBLE WITH AND SHOULD NOT BE MIXED WITH WATER, ALCOHOL OR AROMATIC SOLVENTS, but can be diluted with aliphatic oils (diesel fuel, kerosene or #2 fuel oil). Compatibility with other pesticides has not been thoroughly tested. **DO NOT USE**

(continued on page 2)

SEVIN 4 OIL
EPA REG. NO. 34704-313

(continued from page 1)

IN TANK MIX COMBINATIONS UNLESS YOUR EXPERIENCE INDICATES THAT THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN PLANT INJURY OR MECHANICAL DIFFICULTY. For best mixing when preparing diluted product, add diluent oil to mix tank before adding proper volume of this product. Provide complete recirculation of diluted suspensions once every 24 hours to maintain uniformity.

APPLICATION

On all crops use sufficient gallonage to obtain thorough and uniform coverage. Calibrate equipment to deliver the desired spray volume. Flat fan nozzles may be used to apply both undiluted and diluted product. Rotary atomizing and hollow cone nozzles are not recommended for applying this product undiluted, but can be used if 1 to 1 dilution is made. Use of a high-volume 50-mesh in-line strainer is suggested. Use of screens behind nozzles is not recommended. Apply by air undiluted, or diluted with kerosene, diesel fuel or #2 fuel oil. At temperatures below 50°F (10°C) dilution of 4 volumes of this product with 1 volume of diluent oil is recommended to assure uniform flow and spray distribution. Dilutions greater than 1 to 1 by volume are not recommended; higher dilutions will reduce residual properties. Dilute 1 to 1 by volume when applying with ground equipment. Apply only when weather conditions are favorable. Wind and rising air currents may cause undesirable spray drift and reduce insect control.

INSECT CONTROL

Apply when insects or their damage appear. To maintain control repeat at 7 to 14 day intervals or as necessary unless a shorter interval is specified below. Where a dosage range is indicated, use lower rate on young plants and early instars and higher rate on mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control.

This product does not control spider mites. If spider mites are a problem, use a registered miticide.

Regional differences have been noted in the susceptibility of certain strains of fall armyworm to this product. If local experience indicates inadequate control, use an alternative pesticide.

Follow directions for most effective use. DOSAGES LISTED REFER TO QUARTS OF THIS PRODUCT PER ACRE, NOT TO VOLUME OF FINISHED SPRAY PER ACRE. Refer above for instructions on dilution.

ALFALFA AND CLOVERS INSECT CONTROL

For grasshoppers: Apply 1/2 to 1 1/2 quarts of this product per acre. To avoid possible injury to tender alfalfa foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days. Do not apply to alfalfa within 3 days of harvest.

CORN (field, pop) INSECT CONTROL

For corn rootworm beetles: Apply 1 quart of this product per acre as needed at first silking if populations are at economic levels and/or apply later when needed to control adult population buildup to reduce larval damage to next year's crop. Do not make more than two applications.

For western bean cutworms: Apply a single application of 2 quarts of this product plus 2 quarts diesel fuel, kerosene or #2 fuel oil per acre when an average of 1 in 7 plants have western bean cutworm egg masses or newly hatched larvae and 90 to 100% of the tassels have emerged. Application after 100% of the silks have emerged will reduce effectiveness.

For Japanese beetle: Apply 1 quart of this product per acre when adult beetles are present.

For grasshoppers: Apply 1/2 to 1 1/2 quarts of this product per acre. Lower rate is suggested for nymphs on small plants. Use 1 1/2 quarts when grasshoppers or foliage are mature and greater coverage is required.

FOREST INSECT CONTROL

For control of gypsy moth, fall and spring cankerworms, saddled prominent, forest tent caterpillar, elm sawfly, oak leaf roller complex and Japanese beetle: Apply 1 quart of this product per acre when larvae are in early instars and leaves are at least 1/2 grown, or when adult Japanese beetles are present.

For spruce budworm and western spruce budworm: East of the Rocky Mountains apply 1/2 to 1 quart of this product per acre when a majority of the larvae are in the third through sixth instars. Lower rate is suggested for heavy spruce budworm infestations where 2 applications are necessary for adequate foliage protection. In the Rocky Mountains and west of the Rocky Mountains apply 1/2 to 1 quart of this product per acre when a majority of the larvae are in the third through fifth instars.

For Douglas-fir tussock moth: For maximum foliage protection apply 2 quarts of this product per acre when larvae are in first and second instars. Application to later instars will provide control but with reduced foliage protection.

For population control and foliage protection of high value trees using ground mist blower application: Apply 0.8 pint of spray mixture containing equal volumes of this product and diesel fuel, kerosene or #2 fuel oil per 20 to 30 foot tree (0.2 lb. ai/tree) when larvae are in early instars. Cover foliage thoroughly, but avoid runoff.

PASTURE, RANGELAND AND WASTELAND INSECT CONTROL

For grasshoppers: Apply 1/2 to 1 1/2 quarts of this product per acre. Lower rate is suggested for nymphs on small plants or sparse vegetation. Use higher rate when grasshoppers are mature or vegetation is thick.

For range caterpillar: Apply 1/2 to 1 quart of this product per acre. Use the lower rate when larvae are in early instars and the higher rate for later instars.

For Japanese beetle: Apply 1 quart of this product per acre when adult beetles are present.

For black grass bugs: Apply 1/2 to 1 1/2 quarts of this product per acre. Use lower rate for control of immature bugs in sparse vegetation. Use higher rate for mature bugs or when vegetation is thick.

WHEAT INSECT CONTROL—(DO NOT USE IN CALIFORNIA.)

Do not apply within 21 days of harvest of grain. No time limitation on green wheat used as pasture or forage.


For grasshoppers: Apply 1/2 to 1 1/2 quarts of this product per acre. Lower rate is suggested for nymphs on small plants. Use higher rate when grasshoppers are mature or vegetation is thick.

For cereal leaf beetle: Apply 1 quart of this product per acre. Application is effective against eggs, larvae and adults.

For armyworm and fall armyworm: Apply 1 to 1 1/2 quarts of this product per acre.

NOTICE

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of the product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025