

Tacket

J. Allen Dunlap III  
Platte Chemical Company  
419 18<sup>th</sup> Street  
P.O. Box 667  
Greeley, CO 80632

Dear Mr. Dunlap:

Subject: Revised Labeling - Signal Word WARNING  
Clean Crop C-O-C-S Copodust  
EPA Registration No. 34704-320  
Your Submission Dated November 3, 1994

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



for Cynthia Giles-Parker  
Product Manager (22)  
Fungicide Herbicide Branch  
Registration Division (7505C)

Enclosure

PROPOSED  
BEL



# C-O-C-S Copodust

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:

Nov 25 1994

34704-320

## FUNGICIDE

ACTIVE INGREDIENT	By Wt.
† Copper expressed as metallic .....	6.00%
INERT INGREDIENTS .....	94.00%
TOTAL	
	100.00%

† From copper oxychloride and basic copper sulfate.

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See Below For Additional Precautionary Statements  
And Directions For Use.

EPA REG. NO. 34704-320

EPA EST. NO. \_\_\_\_\_

NET WEIGHT \_\_\_\_\_ LBS. (\_\_\_\_\_ KG)

4307

EXPWPS10G94

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### WARNING

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray dust.

#### Personal Protective Equipment:

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eyewear. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering controls statements:

When handlers use closed systems, enclosed cabs, or a tractor in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (c) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

- Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by disposal of equipment washwaters.

### STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink plenty of water. Do not induce vomiting or give anything by mouth to an unconscious person.

If in eyes: Call physician. Hold eyelids open and flush gently with a steady stream of water for 25 minutes.

If inhaled: Remove victim to fresh air. Apply artificial respiration if not breathing.

If on skin: Wash affected areas with plenty of soap and water. Seek medical attention if irritation occurs.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (40 CFR part 170). This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. If a label contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals, the requirements on this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, waterproof gloves, shoes plus socks and protective eyewear.

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep out of reach of children and animals.

Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

This fungicide is a product containing Copper Oxychloride Sulfate, which is a neutral insoluble copper. It may be applied with airplane or ground application equipment.

### FRUITS AND NUTS

**ALMONDS:** To control Brown Rot and Shot Hole Fungus apply 40 to 50 pounds per acre. Apply in pre-bloom and petal fall period.

**APRICOTS:** To control Shot Hole Fungus use 40 to 50 pounds per acre.

**BEST AVAILABLE COPY**

# C-O-C-S Copodust

EPA REG. NO. 34704-320

Apply as a fall treatment and in spring at red bud blossom. To control Brown Rot use 40 to 50 pounds per acre and apply in pre-blossom and blossom period.

**PEACHES AND NECTARINES:** To control Peach Blight and Peach Leaf Curl use 40 to 50 pounds per acre. Apply as a fall treatment and in spring at bud swell. To control Brown Rot and Peach Blight use 40 to 50 pounds per acre. Apply in pre-blossom, blossom and petal fall periods.

**CHERRIES:** To control Brown Rot and Leaf Spot use 40 to 50 pounds per acre. Apply in popcorn, bloom and petal fall periods.

**APPLES AND PEARS:** In pre-blossom and blossom periods for the control of Fire Blight and Scab use 15 to 30 pounds per acre. All blossoms must be protected ahead of periods of high humidity. Number of applications to be determined by length of the bloom period. For Scab control after the blossom period use a suitable fungicide which does not contain copper. Do not use on Anjou or Comice varieties of pears.

**WALNUTS:** To control Walnut Blight use 20 to 50 pounds per acre. Apply in pre-bloom and post-bloom periods.

**CITRUS (California):** For prevention of Brown Rot use 50 pounds per acre. Apply during the winter months. Treat the ground around the tree and the skirts of the trees to a height of 3 to 4 feet. Give particular attention to covering the fruit. Treat inside as well as outside, covering the trunks and skirts thoroughly. Retreatment may be necessary. Do not disturb soil after treatment. In areas with a history of severe copper damage treat the skirts no more than 2 feet high. Do not apply in areas where copper injury is known to occur or where fumigation with hydrogen cyanide gas is practiced.

**BUSHBERRIES (BLACKBERRIES, RASPBERRIES, BOYSENBERRIES, LOGANBERRIES):** To control Leaf and Cane Spot and Yellow Rust use 50 pounds per acre. Two applications are necessary. Apply first application when new laterals are leafing out, the second when flowers begin to open. A post-harvest application after pruning but prior to the fall rains is of benefit in controlling the above diseases.

**FILBERTS:** To control Bacterial Blight use 50 pounds per acre. Two applications are necessary. Apply first application in August, the second when half of the leaves have dropped. If the fall is unusually wet, apply a third application when most of the leaves have dropped. Apply dust when trees are wet with dew.

## VEGETABLES AND FIELD CROPS

FOR DISEASE CONTROL ON THE FOLLOWING CROPS USE 25 TO 35 POUNDS PER ACRE AT 7 TO 10 DAY INTERVALS TO PROTECT NEW GROWTH THROUGHOUT THE GROWING SEASONS.

**MELONS:** For control of Downy Mildew, Leaf Blight, Scab and aids in control of Bacterial Wilt.

**CARROTS:** For control of Leaf Blight.

**CUCUMBERS:** For control of Downy Mildew, Leaf Blight, Scab and aids in control of Bacterial Wilt.

**EGGPLANT:** For control of Late Blight.

**LIMA BEANS:** For control of Downy Mildew and Pod Spot.

**PEPPERS:** For control of Late Blight.

**LETTUCE:** For control of Downy Mildew.

**POTATOES:** For control of Early and Late Blight.

**SQUASH:** For control of Downy Mildew, Leaf Blight, Scab and aids in control of Bacterial Blight.

**SUGAR BEETS:** For control of Leaf Spot.

**TOMATOES:** For control of Late Blight, Early Blight and Septoria Leaf Spot.


**CELERY:** For control of Celery Blight.

**HOPS:** For control of Downy Mildew As a crown treatment for Early Downy Mildew control apply in the early spring.

## NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION. ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER

373  
EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.



FORMULATED FOR  
PLATTE CHEMICAL CO.  
150 SO. MAIN STREET      FREMONT, NEBRASKA 68025-5697