

34704-326

6/13/2008

1 of 7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUN 13 2008

Mr. John Tice
Manager Registrations
Loveland Products
P.O. Box 1286
Greeley, CO 80632-1286

RE: Notification to change Primary Brand Name: "C-O-C-S WDG"
EPA Registration Number: 34704-326
Date of Submission: May 2, 2008

Dear Mr. Tice:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated May 2, 2008, for the product, Clean Crop C-O-C-S Wettable. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in cursive script, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

STG



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 34704-326	2. EPA Product Manager Tony Kish	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) CLEAN CROP C-O-C-S WETTABLE	PM# PM-22	
5. Name and Address of Applicant (Include ZIP Code) Loveland Products, Inc. P.O. Box 1286 Greeley, Colorado 80632-1286 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

NOTIFICATION

JUN 13 2008

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Change of the Primary Brand Name of this product to: "C-O-C-S WDG".

This notification is consistent with the provisions of PR Notice 98-10 and EPA Regulations at 40 CFR 152.46 and not other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms or PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 50 lbs		5. Location of Label Directions <input type="checkbox"/> _____	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name John T. Tice	Title Manager Registrations	Telephone No. (Include Area Code) 970-347-1484
Certification		6. Date Application Received (Stamped) _____ _____ _____ _____ _____ _____
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.		
2. Signature 	3. Title Manager Registrations	
4. Typed Name John T. Tice	5. Date May 2, 2008	

3087



Performance

Quality

Value

May 2, 2008

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504-P)
U. S. Environmental Protection Agency
2777 S. Crystal Drive, Room S-4900,
Arlington, VA 22202-4501

RE: Clean Crop C-O-C-S Wettable, EPA Reg. No. 34704-326, Change of Primary Brand Name.

Dear Sir or Madam:

Loveland Products, Inc. is submitting an application to permanently change the Primary Brand Name of the product listed above. Loveland Products would like to change the name to "C-O-C-S WDG". Enclosed you will find the Notification Form (8570-1) with appropriate certification statement and two copies of the current approved label with the new name.


If you have any questions, please feel free to call or contact me at 970-347-1484 or email at John.Tice@uap.com.

Sincerely,

John Tice
Manager Registrations
Loveland Products, Inc.

Enclosures (2)

JUN 13 2008



C-O-C-S WDG

FOR ORGANIC PRODUCTION

This product meets the Standards set forth by The National Organic Program (NOP) final rule 7 CFR Part 205.

FUNGICIDE

ACTIVE INGREDIENTS:	
Copper oxychloride*	74.8%
Basic copper sulfate*	14.2%
INERT INGREDIENTS	11.0%
TOTAL	100.0%

*Metallic copper equivalent 50%.

KEEP OUT OF REACH OF CHILDREN WARNING—AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See Below For Additional Precautionary Statements And Directions For Use.

EPA REG. NO. 34704-326

EPA EST. NO. 45002-MX-2

NET WEIGHT 20 LBS. (9.07 KG)

IHT

06G06

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin, or on clothing. Avoid breathing dust (vapor or spray mist). Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves, shoes plus socks and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not allow rinsate from cleaning of equipment or disposed material to enter surface or ground water.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:

1-800-301-7976.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves, shoes plus socks and protective eyewear.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

C-O-C-S WDG is a dry Copper material designed to be used as a spray either alone or in combination with a spreader or sticker. Do not use in undiluted form. Do not apply this product through any type of irrigation system.

C-O-C-S WDG should not be applied in a spray solution having a pH less than 6.5 or greater than 9.0 as phytotoxicity or loss of effectiveness may occur. Environmental conditions such as extended period of wet weather, acid rain, etc., which alter the pH of the leaf surface may also affect the performance of this product resulting in possible phytotoxicity or loss of effectiveness.

Mixing Instructions: Fill the spray tank one-fourth to one-third full with clean water. Start agitation (proper agitation creates a rippling or rolling action on the liquid surface). Add C-O-C-S WDG at the recommended rate. Mix thoroughly. Add water to fill spray tank. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture. Thorough coverage is required. Apply when the disease is reported in the area. Repeat at 7 to 10 day intervals or as required to maintain effective control.

The following instructions are based on general applications. Except as specified begin applications before or at first sign of disease and repeat at 7 to 10 day intervals throughout the growing season to protect new growth. When a rate range is given, apply higher doses when climatic conditions most favor disease(s).

ORNAMENTALS

Amounts listed are rates per 100 gallons of water. Application to blooming plants may cause browning of petals. C-O-C-S WDG has demonstrated excellent plant safety; however, not all cultivars have been tested. **Before treating a large number of plants of a particular cultivar, treat a few plants and observe prior to full scale applications.**

CULTIVAR	DISEASE	RATE PER 100 GALS.	REMARKS
Azalea	Bud Blight, Twig Blight	4 lbs.	Begin application when disease is reported in area.
Maple	Tar Leaf Spot	4 lbs.	Safe with 2 pounds hydrated lime per 100 gallons of water.
Oak (White)*	Anthraxnose	4 lbs.	Safe with 2 pounds hydrated lime per 100 gallons of water.
Pachysandra*	Canker, Leaf Spot, Twig Blight	3 lbs.	Safe with 3 pounds hydrated lime per 100 gallons of water.
Vinca*	Canker, Dieback	4 lbs.	Begin applications when disease is first reported in area.
Sycamore	Sycamore Anthracnose (Sycamore Blight)	3 lbs.	Begin applications when buds begin to swell. Repeat about 7 to 14 days later at early leaf stage.
Flowering Plants* (Dahlias Nasturtiums Peonies Roses)	Botrytis Blight, Leaf Spots Leaf Spots Anthraxnose, Botrytis Blight Black Spot, Leaf Spots, Rust	2½ lbs.	Begin applications before diseases appear. For Black Spot on roses, begin when leaf buds swell in early spring. Repeat at 7 to 10 day intervals and after rains.

*Not for use in California

VEGETABLES & FIELD CROPS

C-O-C-S WDG is adapted to most types of spray equipment. Unless otherwise stated, apply the following rates of C-O-C-S WDG: For dilute high volume spray, use a minimum of 25 gallons of water per acre; for concentrate ground sprays apply a minimum of 5 gallons per acre; and for aerial spraying, a minimum of 3 gallons per acre should be applied.

CROP	DISEASE	RATE PER ACRE	REMARKS
Beans (dry and succulent)	Cercospora Leaf Spot, Bacterial Blight, Anthracnose, Downy Mildew	2-4 lbs.	Begin application when plants are 5-6 inches high and repeat as needed.
Beets (table)	Downy Mildew, Leaf Spot	3-4 lbs.	Begin applications prior to onset of disease.
Brassica Leafy Vegetables (Broccoli, Brussels Sprouts, Cabbage, Cauliflower)	Downy Mildew, Black Rot	3-4 lbs.	Begin applications after transplants are set in field or after emergence of seedlings. Caution: Reddening of older leaves may occur on Broccoli and flecking may occur on wrapper leaves of cabbage.
Carrots	Leaf Blight, Leaf Spot	2-4 lbs.	Repeat applications are necessary to protect new growth.
Celery	Blight, Downy Mildew	2-4 lbs.	
Cucurbits (Cucumbers, Melons, Squash [Summer & Winter], Pumpkins, Watermelon)	Downy Mildew, Leaf Spots, Anthracnose	2-4 lbs.	Also aids in the control of Bacterial Wilt.
Eggplant, Peppers	Downy Mildew, Alternaria, Phytophthora Blight, Anthracnose	2-4 lbs.	
Lettuce (Head and Leaf)	Downy Mildew	1-3 lbs.	Under adverse weather conditions flecking of wrapper leaves may occur.
Onions	Downy Mildew, Purple Blotch	3-4 lbs.	Apply when plants are 4-8 inches high. If disease infection is severe, increase dosage up to 7 pounds on larger plants.
Potatoes	Early Blight, Late Blight	1½-4 lbs.	
Spinach	Downy Mildew, Leaf Spot	1-3 lbs.	Caution: Leaf flecking may occur under adverse weather condition.

6097

CROP	DISEASE	RATE PER ACRE	REMARKS
Strawberries	Leaf Spot, Bacterial Blight	2-4 lbs.	Begin applications when plants are established and continue as needed. Apply in a minimum of 100 gallons per acre. Stop applications if any signs of phytotoxicity begin to appear.
Sugarbeet	Cercospora Leaf Spot	2-4 lbs.	May be tank mixed with sulfur to enhance control.
Tomatoes	Early Blight, Late Blight, Septoria Leaf Spot	2-4 lbs.	If Late Blight infection is severe, increase dosage up to 7 pounds on large plants.
	Bacteria Spot	2-6 lbs.	Apply every 5-7 days after true leaves appear.
Vegetable Seedlings	Damping off	4 lbs./100 gals. or 2 oz./3 gals.	For damping off in cold frames, greenhouses, etc., begin treatment when seedlings emerge. Spray plants and surrounding soil lightly but thoroughly. Do not use on cabbage or related seedlings.
Wheat, Barley*	Helminthosporium Spot Blotch, Septoria Leaf Blotch	1-2 lbs.	Make first application at early heading and follow with second spray 10 days later.

*Not for use in California

TREE FRUIT & VINE CROPS

Rates listed are pounds per 100 gallons of water and/or pounds per acre. Unless otherwise noted, C-O-C-S WDG may be applied by air or concentrate applications. For concentrate ground spray, use equivalent rates of C-O-C-S WDG in a minimum of 25 gallons of water per acre. For air application, use equivalent ground rates in a minimum of 10 gallons of water per acre. When used in combination with oil, observe the precautions on the oil label. Spray for thorough coverage at indicated times.

CROP	DISEASE	RATE OF PRODUCT PER 100 GALS.	PER ACRE	REMARKS
Almonds	Coryneum Blight	4 lbs.	12-16 lbs.	Apply as a dormant/delayed dormant spray up to bud swell. May be combined with 2 gal. dormant oil per 100 gallons of water.
	Coryneum Blight, Brown Rot	4 lbs.	12-16 lbs.	Apply pre-bloom and petal fall.
Apples	Fireblight*	2-4 lbs.		Make one application between silver tip to green tip. (Discontinue at this rate when green tip exceeds 1/2 inch as phytotoxicity may occur.)
	Fireblight	1/4 lbs.	1 lb.	Bloom spray: Use 1/4 pound when 20% of blossoms are open and repeat when 75% are open.
	European Canker	4-5 lbs.		Apply post harvest in fall of year. Apply as dilute spray. Combine with dormant oil at 1/2 to 1 gallon per 100 gallons of water.
Apricots	Coryneum Blight, Brown Rot	4 lbs.	5-12 lbs.	For shothole, apply as a dormant spray up to red bud. May be combined with dormant oil at 2 gallons per 100 gallons water or at 3-6 gallons dormant oil per acre concentrate or air. For Brown Rot, apply in pre-blossom, blossom and petal fall periods.
Caneberries (Boysenberries, Blackberries, Youngberries, Loganberries, Red and Black Raspberries, Dewberries)	Leaf and Cane Spot, Yellow Rust	4-6 lbs.	4-6 lbs.	For Leaf and Cane Spot: Apply when new laterals are leafing out and repeat when flowers begin to open. A post-harvest spray after pruning is also beneficial. Apply in a maximum of 150 gallons per acre.
Cherries*	Dead Bud, Brown Rot, Leaf Spot	4-6 lbs.	8-12 lbs.	Dead Bud: Apply in fall before heavy rains and again in spring before bud swell. May be combined with 1 pint per 100 gallons water dormant spray oil. Brown Rot and Leaf Spot: Apply in popcorn, bloom and in petal fall periods.
Cherries (Tart)*	Cherry Leaf Spot	1 1/2 lbs.		Apply as a dilute spray. Combine with 3 pounds hydrated lime per 100 gallons of water to reduce copper injury.
Cherries (California)	Brown Rot, Leaf Spot	4-6 lbs.	7.5 lbs.	Apply in popcorn, bloom and petal fall periods.
Citrus Fruits (Citrus Citron, Grapefruit, Kumquat, Lemons, Limes, Oranges, Tangelos, Tangerines)	Melanose*, Scab*, Greasy Spot*	1 1/4-2 lbs.	4-12 lbs.	Apply as pre-bloom and post-bloom sprays. For Melanose and Scab, repeat 4 weeks later. Do not apply after fruit is 3/4 inch in diameter. Do not apply in areas where copper injury is known to occur.
	Brown Rot (California and Arizona Only)	1-2 lbs.	4-10 lbs.	Begin applications in fall before or just after first rains. In severe Brown Rot season a second application may be necessary. Spray as overall or skirt spray insuring that ground around tree receives spray coverage. Give particular attention to covering fruit. Treat inside as well as outside, spraying trunks and skirts thoroughly. Do not disturb soil after treatment. Apply a minimum of 100 gallons per acre ground concentrate and 20 gallons of water per acre by air. When danger of copper injury is severe, spray mix may be safened by adding 1/3 to 1 pound hydrated lime per 1 pound fungicide.
Filberts*	Bacterial Blight	4 lbs.	12-16 lbs.	Apply as a post-harvest spray. In seasons of heavy rainfall apply a second spray when 75% of the leaves have dropped. One (1) pint superior type oil per 100 gallons of water may be added.
Grapes	Black Rot, Downy Mildew, Powdery Mildew		2 lbs.	Mildews: Begin applications at or near bud break. Thoroughly spray both upper and lower leaf surfaces. Repeat as needed. Black Rot: Apply when buds break and repeat when shoots are 10 inches long, when blossoms fall and berries reach the size of a BB shot. Foliage injury may occur on copper sensitive varieties. Where danger of copper injury is severe, add 4 pounds of hydrated lime in a minimum of 125 gallons of water per acre to safen spray mixture.
Hops*	Downy Mildew		4-6 lbs.	Apply soon after training vines (when growth is approximately 18 inches long) in 100 gallons of water per acre. Apply as a directed spray using adequate spray volume for good coverage.
Olives	Olive Leaf Spot	4 lbs.		Apply as a full coverage spray after harvest and prior to rainy season. Under heavy disease pressure, a second application may be necessary.

*Not for use in California

CROP	DISEASE	RATE OF PRODUCT PER		REMARKS
		100 GALS.	ACRE	
Peaches and Nectarines	Bacterial Spot, Peach Blight, Peach Leaf Curl, Brown Rot	4 lbs.	12-16 lbs.	For Leaf Curl, Peach Blight and Bacterial Spot: Apply as a dormant spray up to bud swell. May be combined with dormant oil at 2 gallons per 100 gallons of water or at 3-6 gallons dormant oil per acre concentrate or by air. For Brown Rot: Apply in pre-blossom, blossom and petal fall periods.
Pears* and Quince*	Fireblight	2-4 lbs. ¼-½ lb.	1-2 lbs.	For dormant to bud swell, apply as a full cover spray. Apply at 10 bloom and repeat at 3-5 day intervals until all late bloom is over.
	Pseudomonas, Anthracnose	4 lbs.	12-20 lbs.	Apply after harvest and before rains fall and again in the dormant to delayed dormant period. Apply in a minimum of 100 gallons per acre by ground and 20 gallons per acre by air.
Pears and Quince (In California)	Fireblight	¼-½ lb.	1-1½ lbs.	For Fireblight: Apply at 10 bloom and repeat at 3-5 day intervals until all late bloom is over.
	Pear Blight	4 lbs.	12 lbs.	For Pear Blight: Apply during bloom and other periods favorable to blight infections.
	Pseudomonas, Anthracnose	4 lbs.	2-20 lbs.	Apply after harvest and before rains fall and again in the dormant to delayed dormant period. Apply in a minimum of 100 gallons per acre by ground and 20 gallons per acre by air.
Plums and Prunes	Brown Rot, Shot Hole	4 lbs.	6-12 lbs.	For Shot Hole: Apply as a dormant spray up to bud swell. May be combined with dormant oil at 2 gallons per 100 gallons water or with 3-6 gallon dormant oil per acre concentrate or by air. For Brown Rot: Apply in pre-blossom, blossom and petal fall periods.
Walnuts	Walnut Blight	3-4 lbs.	12-16 lbs.	Apply in each of the following growth stages: pre-bloom, bloom (1% and 10-20% bloom) and post-bloom. When rain threatens, additional applications made immediately before or after rains are important.

*Not for use in California

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80632-1286

Product of Mexico