

Code: 30027

C-O-C-S® 50-V

C-O-C-S 50-V is a fixed copper material designed to be used as a spray either alone or in combination with spreader and sticker. Apply C-O-C-S 50-V to the control of those diseases against which it is available copper fungicide may be used.

DIRECTIONS FOR USE

Fill the spray tank half full of water. With the sprayer running, add in about the required amount of C-O-C-S 50-V. Afterwards add the required amount of an insecticide if it is desired to use it. Finish filling the tank with water and apply at once.

FOR DISEASE CONTROL ON THE FOLLOWING CROPS BEGIN APPLICATION AS NECESSARY AND REPEAT AT 7 TO 10-DAY INTERVALS TO PROTECT NEW GROWTH THROUGHOUT THE GROWING SEASON. USE 3 TO 4 POUNDS OF C-O-C-S 50-V IN 50 TO 100 GALLONS OF WATER PER ACRE.

- CARROTS - Leaf Blight.
- CHESTNUT - Colony Blight.
- CUCUMBER, PEPPERS - Late Blight.
- GRAPES - Downy Mildew. As a disease treatment for Early Downy Mildew control apply in the early spring.
- LEAF BEANS - Downy Mildew, and Pod Spot.
- MELONS, CUCUMBERS - Downy Mildew, Leaf Blight, Leaf and stem blight in control of Bacteria Wilt.
- ONIONS, LETTUCE - Downy Mildew.
- POTATOES - Early and Late Blight.
- SQUASH - Downy Mildew, Leaf Blight, leaf and stem in control of Bacterial Wilt.
- SWEET PEAS - Leaf Spot.
- TOMATOES - Late Blight, Early Blight and Septoria Leaf Spot.

FUNGICIDE	
EPA REGISTRATION NO. 279-1077 2A	EPA EST. 279-CA-2
ACTIVE INGREDIENTS	BY WT.
†Copper expressed as metallic	30.00%
INERT INGREDIENTS	70.00%
	100.00%
†From Copper Oxide Chloride Sulfate	
C-O-C-S - FMC Trademark	

SHIPPING OFF: For the control of Downy Mildew, and certain other diseases, it will be better by spraying the plants with 50 to 100 gallons of water. Apply during the week and the plants should be kept well watered. 4 to 7-day intervals, beginning ground level. The plants live ground. DO NOT use this for control of Downy Mildew.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. Avoid inhaling dust. Avoid skin contact. Keep away from domestic animals, food and feedstuffs.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the area treated. Do not apply when run-off is likely to occur. Apply this product only as specified on the label.

Remove equipment and DISPOSE OF WASTES by burying in non-erosion land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of streams.

For Emergency Assistance Call - 714-726-2766

PREPARATION, EQUIPMENT, LABELING (CONTAINER LABELING): For use of C-O-C-S 50-V and 1 to 1 1/2 pounds of water. Apply during the week and the plants should be kept well watered. 4 to 7-day intervals, beginning ground level. The plants live ground. DO NOT use this for control of Downy Mildew.

Containers Should Not Be Reused: Tanks of Soda or Gas or other tanks used in the following manner. Emptying the tanks and other containers used in the production of this material when such use is contrary to label instructions. Directions and Recommendations. Mixing and method of application with other chemicals. Other influencing factors in the control of the disease and a Loss of Product. FMC's recommendations are based upon tests of this product beyond the guarantee, expressed or implied, or the results to be obtained. Directions or established procedures. Damage. Buyer's or user's or breach of warranty or negligence damages not exceeding the value include incidents of product.



FMC Corporation
Agricultural Chemical Division
MIDDLEPORT, N.Y. MARIETTA, GA.
PESQUERA, CA. RICHMOND, VA.

ACCEPTED
AUG 6 1976
U.S. Dept. of Agriculture
Federal Insecticide, Fungicide, and Rodenticide Act
EPA 79-1072

BEST AVAILABLE COPY

C-O-C-S® 50-V

Net Weight
20 Pounds

designed to be used in
with sprayer and sticker.
no disease against which
is not.

USE
with the sprayer running.
C-O-C-S 50-V. A fungicide
only if it is desired to use
it as a fungicide.

STARTING CROPS BEGIN APPLI
7 7 2-DAY INTERVALS
IN THE GROWING SEASON.
1/4 TO 1/2 GALLONS OF

treatment for Early Blight
and Spot.
Leaf Blight, Cork and other

and other leaf spots.

and other leaf spots.

FUNGICIDE
EPA REGISTRATION NO. 279-1077 ZA EPA EST. 279-CA-2
ACTIVE INGREDIENTS: BY WT. 30.00%
(Concentrated as material) 50.00%
INERT INGREDIENTS: 180.00%
From Copper Dichloride Sulfate
C-O-C-S - FMC Trademark

CAUTION

KEEP OUT OF REACH OF CHILDREN

Hazardous if inhaled. Avoid inhaling dust. Avoid skin contact. Keep
away from domestic animals, food and feedstuffs.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes,
ponds and streams. Do not contaminate water by cleaning of
equipment or disposal of wastes. Do not apply when weather
conditions favor drift from the areas treated. Do not apply
where runoff is likely to occur. Apply this product only as
specified on the label.
Wash equipment and DISPOSE OF WASTES by burying in non-
crop land away from water supplies. Empty bags should be
buried with wastes or burned. Keep out of ponds.

For Emergency Assistance Call - 714-726-3740

ACCEPTED
AUG 6 1976
Under the Federal Insecticide,
Fungicide and Rodenticide Act.
EPA REG. NO. 279-1077

DAMPING OFF for the control of Damping-Off of tomatoes,
peppers, and certain other crops in hot beds, coldframes, and
greenhouses, it will be helpful to supplement seed treatment
by spraying the plants with C-O-C-S 50-V at the rate of 4
pounds in 50 to 100 gallons of water per acre. The suspension
should be kept well agitated. The applications should be made at
4 to 7 day intervals, beginning when the plants break through the
ground. Spray the plants lightly but thoroughly, covering the
ground. DO NOT use this on Cabbage and related crops for the
control of Damping-Off.

GRAPEFRUIT, TANGELO, LEMONS, LIMES, ORANGES, TAMARINDS,
TANGELOS (California) for prevention of Brown Rot use 1 pound of
C-O-C-S 50-V and 1 to 1 1/2 pounds of hydrated lime per 100 gallons
of water. Apply during the winter months. Spray the ground around
the tree and the skirts of the trees to a height of 3 to 4 feet. Give
particular attention to covering the fruit. Treat inside as well
as outside, spraying inside and skirts thoroughly. 4 to 6 gallons
per tree is usually adequate. Irrigation may be necessary. Do not
disturb soil after treatment. In areas with a history of severe
canker damage, spray the skirts no more than 2 feet high. Do not
apply in areas where canker injury is known to occur or where
fungicide with hydrogen cyanide gas is practiced.

Dealers Should Sell in Original Packages Only
Terms of Sale or Use on purchase of this product buyer and user
agree to the following conditions:
Reversely FMC makes no warranty, expressed or implied,
concerning the use of this product other than indicated on the
label. Except as so warranted, the product is sold as is. Buyer
and user assume a trial of use and of handling, and of storage
of this material under such use and of handling and of storage
is contrary to label instructions.

Directions and Recommendations Follow directions carefully.
Timing and method of application, weather and crop conditions,
mixture with other chemicals not specifically recommended, and
other influencing factors in the use of this product are beyond
the control of the seller and are assumed by buyer at his own risk.
Use of Product: FMC's recommendations for the use of this pro-
duct are based upon tests believed to be representative. The use of
this product being beyond the control of the manufacturer, no
guarantee, expressed or implied, is made as to the effects of such
or the results to be obtained if not used in accordance with
directions or established safe practice.

Damages Buyer's or User's exclusive remedy for damages for
breach of warranty or negligence shall be limited to direct
damages not exceeding the purchase price paid and shall not
include incidental or consequential damages.

Printed in U.S.A.

BEST AVAILABLE COPY