

CAPTAN 10 DUST

Fungicide

For Agricultural or Commercial Use Only.

ACTIVE INGREDIENTS:	By Wt.
*Captan	9.77%
Related Derivatives	0.23%
INERT INGREDIENTS:	90.00%
TOTAL	100.00%
*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide	
* CLEAN CROP is a Registered TM. of United Agri Products, Inc.	

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

PRECAUCION AL USUARIO: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

EPA REG. NO. 34704-342 EPA EST. NO.

NET WEIGHT 50 POUNDS

1741

EXP-20

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes irreparable eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician.
If swallowed: Call physician or Poison Control Center immediately. Drink plenty of water. Do not induce vomiting. Do not give anything by mouth to an unconscious person.
If inhaled: Remove victim to fresh air and breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.
If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). DRA and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply the product in such a manner as to directly or indirectly injure through drift, sprays, contact or other persons. The area being treated must be vacated by unexposed persons. Wear protective clothing and shoes when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and shoes before eating, drinking, or smoking. Do not eat, drink, or smoke while using this product. Do not wear a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron. A dust mask must be worn when mixing and loading this product. Footwear and handwear must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and shoes with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

During application only, from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit a long-sleeved shirt and long-legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while sitting.

RE-ENTRY STATEMENT

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Consequently, post-reentry information at site of application. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be treated without specific protective clothing. Period of time this must be vacated and appropriate actions to take in case of accidental exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given whenever it is known to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

DANGER

Area treated with Captan on (insert date). Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn (insert State Department of Agriculture re-entry interval, if more restrictive).

In case of accidental exposure to persons, body or skin, wash the skin thoroughly with soap and water. Remove contaminated clothing and shoes before rinsing. If in eyes, flush with plenty of water. Call a physician if inhaled, remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

STORAGE AND DISPOSAL

PERIODIC STORAGE: Use for use or storage in or around the house. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After part of use, hold and not back bags, clamp and close tightly. Do not put contents or dust material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact. Store in a cool, dry place and keep out of reach of children and unexposed persons. Contain spills. To contain spill: If liquid, soak surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Storage and Disposal Continued

PESTICIDE DISPOSAL: Pesticides will not be acutely hazardous. Improper disposal of excess pesticide, spray material, or waste is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Apply to give thorough coverage to upper and lower leaf surfaces. May be applied by ground or air. Observe days interval between application and harvest indicated by number in parentheses following crop.

Almonds (A): Brown Rot, Shot Hole, Scab, Leaf Blight—Use 25 to 50 pounds per acre. Apply at petal-fall, bloom, petal-fall stages and up to 5 weeks after petal-fall. When applied after petal-fall, do not feed almonds hulls to dairy animals or animals being finished for slaughter.

Apples (A): Apple Scab—Use 20 to 80 pounds per acre. Apply in prebloom, bloom, petal-fall, and first cover periods. For secondary infection use 11 to 25 pounds per acre and apply in cover periods. Black Rot, Black Rot, Black Rot, Black Rot—Use 11 to 37½ pounds per acre and apply in cover periods. Black Rot, Black Rot, Black Rot, Black Rot—Use 20 to 37½ pounds per acre. Apply in late cover and preharvest periods.

Apples (B): Brown Rot, Jacket Rot—Use 50 pounds per acre. Apply at red bud stage and repeat at 75 percent petal-fall.

Cherries (C): Cherry Leaf Spot—Use 30 to 40 pounds per acre. Apply in pink, full bloom, petal-fall and first cover periods. Brown Rot—Use 11 to 50 pounds per acre. Apply in pink, full bloom, petal-fall, shock, and cover periods.

Grapes (B): Botrytis Rot (Bunch Rot)—Use 40 to 50 pounds per acre. Apply twice before bloom, and once just after bloom. Repeat periodically to make 3 cover applications before bunches close. Apply to every row and direct dust toward the clusters. To be effective, dust must be applied before rain. Whenever rain threatens apply an additional dust. Applications made following rainfall have been of very little benefit in controlling Botrytis. Ground application equipment is preferred.

Grapes: Mold control for fruit on drying trays in fields—*Penicillium* spp., *Aspergillus* spp., *Botrytis* spp.—Use 10 pounds per acre. Apply to both trays for protection of fruit and prevention of bloom rot. Apply to both trays for even coverage. On dry damp weather, make a second application after trays are turned. Apply the dust by hand or ground dusters only. Do not apply within 5 days of a preharvest application of captan.

Peaches (B), Nectarines (B): Brown Rot, Peach Scab—Use 50 pounds per acre. Make 2 to 3 prebloom applications. Repeat at petal-fall making 3 to 4 cover sprays as necessary. Make preharvest application where necessary.

Pears (B): Pear Scab (except Petite Northwest)—Use 20 to 30 pounds per acre. Make first applications during early flower and petal-fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Rusting may be produced on Boar Pears. Do not use on D'Anjou Pears.

Plums (B), Prunes (B): Brown Rot—Use 20 to 50 pounds per acre. Apply at red bud and 75 percent bloom, full bloom, petal-fall and in cover sprays as necessary. Thorough coverage is essential for maximum control.

Strawberries (B): Botrytis Rot—Use 20 to 40 pounds per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat applications weekly. Severe infection conditions: Continue applications through harvest period, treating sprayed area after each picking.

Cantaloupes (B), Cucumbers (B): Downy Mildew, Angular Leaf Spot, Anthracnose—Use 20 to 25 pounds per acre on young plants, increasing to 30 to 35 pounds per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 7 day intervals until danger of infection is past.

Potatoes (B): Early Blight, Late Blight, Damping-Off—Dust at 7½ ounces per bushel cut seed pieces (1½ cut or ¼ pound to 100 pounds cut seed pieces). Apply in conventional container or by dusting attachment over cut. Cut pieces should be treated within 8 hours of cutting. If planting is to be delayed more than 1 to 2 days the treated pieces should be stored for 2 days in open crates before bagging.

Tomatoes: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—Young Plants—Use 25 to 30 pounds per acre. Mature Plants—Use 35 to 50 pounds per acre. Begin applications at first sign of disease. Repeat at 5 to 7 day intervals.

Ornamentals: Roses—Black Spot, Camellias—Rust and Leaf Spot—Chrysanthemums—Flower Blight—Use 25 to 30 pounds per acre. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

Box and Greenhouse Bench: The utmost pre-planting for Damping-Off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrub and tree), many flowers and lawns. Seedlings—Use 1½ to 2 pounds per 1000 square feet (50 to 60 pounds per acre). Walk into upper 3 to 4 inches of soil before planting.

Only the applicator is permitted to be in the greenhouse during application of captan. Open vents to greenhouse during application and for at least 1 hour after application.

When planting in captan-treated soil must wear chemical resistant gloves.

Block: Botrytis Rot—Use 40 to 50 pounds per acre. Apply when disease appears. Repeat at weekly intervals as needed.

Turf and Lawns: Brown Patch, Leaf Spot, Damping-Off, Root Rot—Use 1 to 2 pounds per 1000 square feet. Begin treatment before infection occur. Repeat at 7 to 10 day intervals throughout summer. Do not graze treated areas. Do not feed clippings to livestock.

Dealers should sell in Original Package Only

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions.

Warranty: Platte makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user shall be at all risk of use under handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other factors influencing factors in the use of this product are beyond the control of the seller and are assumed by the user at his own risk.

Use of Product: Platte recommends the use of this product as a seed treatment and as a fungicide to be reliable. The use of this product beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established best practices.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET FREMONT, NEBRASKA 68015