

DIRECTIONS

This product should be applied as a coarse (large droplet) spray-in spot treatment of localized surface areas only.

Roaches and Waterbugs Remove all dishes, food and utensils from the area to be sprayed. Make spot applications to infested roach haunts — behind sinks, into cracks and crevices, the underside of all drawers, baseboards, pipeholes in walls, and cracks of kitchen cupboards. Inspect and treat other areas in the house where infestation is evident. Pay particular attention to bathrooms, cracks and crevices in closets, dresser drawers and frames.

Ants Spiders Silverfish, Crickets Earwigs, Firebrats, Centipedes, Millipedes, Sowbugs and Springtails Spray areas where these pests are found or may hide including dark corners of rooms and closets, cracks and crevices, along baseboards and door sills and frames, behind and beneath sinks, stoves, refrigerators, and cabinets, and around garbage cans, plumbing and other utility installations. Pests driven out of hiding places should be sprayed directly. Spray ant trails, nests, and points of entry. Repeat treatment as required.

Carpet Beetles Make localized application to areas of the floor and baseboards. Spray directly into cracks and under carpets where insects may be found. Treat crevices, cracks and infested areas of shelving in closets. Repeat as necessary.

Flea Dog Ticks Spray infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of pets and localized areas of floors and floor coverings. Repeat as necessary. Bedding should be replaced in animal quarters after treatment. Do not treat animals.

Fleas Mosquitoes and Clover Mites Use only on outdoor surfaces. Spray outside surfaces of windows and door frames and other areas where pests may enter the home. Flies and mosquitoes coming to rest on treated surfaces will be killed. Repeat treatment when effectiveness diminishes.



READY-TO-USE

RESIDUAL

ALSO KILLS: SPIDERS • WATERBUGS • SILVERFISH

ACTIVE INGREDIENTS:

0,0-Diethyl 0-(2, isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....0.5001
 Pyrethrins.....0.0521
 *Piperonyl Butoxide, Technical.....0.2601

INERT INGREDIENTS:

Petroleum Distillate.....99.1801
 100.0001

*Equivalent to 0.2001 of (butylaurthyl) (6-propylpiperonyl) ether and 0.0521 of related compounds.

3561 West 105th St.

Cleveland, Ohio 44111

CAUTION

Harmful if swallowed; combustible; may be absorbed through skin; avoid contact with eyes, skin, and clothing. Avoid breathing vapors or spray mist. Keep away from food, feedstuffs, and domestic water supplies. Do not smoke while using. Wash thoroughly after handling. **DO NOT POUR, USE, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** If swallowed do not induce vomiting. Call a physician immediately.

NOTE TO PHYSICIAN: Contains petroleum distillate. Active ingredients contain a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

Do not allow children or pets in treated areas until surfaces are dry. Do not spray air in rooms or use on plants. Apply as a surface spray only as specified on this label. Do not allow spray to contact food, feedstuffs, or water supplies. Thoroughly wash dishes and food handling utensils if contaminated with this product. Remove pets, birds, and cover fish bowls before spraying. Do not apply this insecticide to asphalt tile or foam backed carpeting.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning equipment or disposal of wastes. Do not reuse empty container. Destroy empty container by perforating or crushing and burying in a safe place away from water supplies.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

EPA Reg. No. 36272-6-8408
 EPA Est. No. 36272-OH-1

Net Contents:

171

24036

PHOSDRIN is a registered T.M. of Shell Chemical Co.

02769 A

953
53

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store in or around the home.

Do not store below 20°F. (-7°C).

Do not use or store near heat, open flame or hot surfaces.

Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**SEE ATTACHED BOOKLET FOR COMPLETE DIRECTIONS FOR USE.
DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.**

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Platte makes no warranty, express or implied, concerning this use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: Platte's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, express or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established sale practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

EPA REG. NO. 34704-343 EPA EST. NO. _____
NET CONTENTS _____ GALLONS

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
188 SO. MAIN STREET FREMONT, NEBRASKA 68925



POISON

6

MEVINPHOS MIXTURE, LIQUID RQ NA2783

ing aphids, use 1 quart per acre, but do not treat within 4 days of harvest.

MELONS (Including Cantaloupe, Honeydew Melons, Muskmelons, Watermelons) (1): Aphids—Use 1/4 to 1/2 pint per acre. Cabbage Looper, Dipterous Leafminer (adult), Leafhoppers, Lygus Bugs, Mites, Salt-Marsh Caterpillar, False Chinch Bugs, Climbing Cutworms, Grasshoppers—Use 1/2 to 1 pint per acre. To control Rindworms (Cabbage Looper, Cutworms, Salt-Marsh Caterpillars, Tobacco Budworms) on watermelons apply 1/2 pint per acre.

OKRA (1): For control of Aphids use 1/4 to 1/2 pint per acre. For control of Climbing Cutworms, Corn Earworms, Green Stink Bug, Mites and Velvet Bean Caterpillar—Use 1/2 to 1 pint per acre.

ONIONS (Including Green Onions) (1): Thrips, Climbing Cutworms—Use 1/2 to 1 pint per acre.

PARSLEY (5)*: For control of Aphids apply 1 to 2 pints (0.5 lb. to 1.0 lb. a.i./acre) as a foliage application using a minimum of 40 gallons of water per acre by ground application or 10 to 15 gallons of water per acre by air application. For light infestation use lower dosage of 1 pint (0.5 lb. a.i./acre). *For hard to kill insects and increased populations use 2 pints (1.0 lb. a.i./acre). Do not treat later than 8 days before harvest for higher doses. Begin application when insects first appear and repeat as often as necessary to maintain control. Do not treat more than 3 times between harvests.

PEAS (Including Vines) (1): Aphids—Use 1/4 to 1/2 pint per acre. For control of Grasshoppers, Leafhoppers, Mites and Climbing

Use 1 quart per acre, but do not treat within 4 days of harvest.

SUMMER SQUASH (1): For control of Aphids use 1/4 to 1/2 pint per acre. For control of Cabbage Looper (adult), Leafhoppers, Lygus Bugs, Mites, False Chinch Caterpillar, Climbing Cutworms, an

TOMATOES (Outdoor) (1): Aphids—Use 1/4 to 1/2 pint per acre. For control of Grasshoppers, Leafhoppers and Mites—Use 1/2 to 1 pint per acre.

TURNIPS (3): For control of Aphids—Use 1/4 to 1/2 pint per acre. For control of Cabbage Looper, Imported Cabbageworm, Leafhoppers, Mites, False Chinch Caterpillar, Climbing Cutworms, an

GREENHOUSE: Diluted product should be used in greenhouse (close all doors, windows and

barricade all entrances, post warning signs and precautions are necessary to prevent un-

authorized persons from entering the treat-

ment. Must wear a full-face mask of a type found on the market for protection. Apply any time the greenhouse is closed for two hours without endangering

persons. Such as early morning, cloudy day. If application is made during the day, keep it

tightly for at least two hours, then ventilate thoroughly for an hour before resuming regular work. If made in the late afternoon, the house can be

On the following morning, ventilate for one hour to render the air safe for regular work.

LETTUCE (Including Leaf, Bibb and Romaine) (10): Aphids, Corn Earworms, Climbing Cutworms, Dipterous Leafminer (adult), Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Mites, Lygus Bugs, Salt-Marsh Caterpillars, False Chinch Bug and Thrips—Use 1 1/4 to 2 1/2 pints per 50,000 cu. ft.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Platte makes no warranty, express or implied, concerning this use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: Platte's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, express or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

