



RESTRICTED USE PESTICIDE

Due to Very High Acute Toxicity to Humans and Residue Effects on Avian, Mammalian and Aquatic Species For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

PHOSDRIN® 4.0 MISCIBLE

Insecticide

ACTIVE INGREDIENTS:	BY WEIGHT
† Mevinphos (2-Carbomethoxy-1-methylvinyl dimethyl phosphate, alpha isomer)	28.50%
† Related Compounds	19.00%
INERT INGREDIENTS:	52.50%
	TOTAL
	100.00%

† Equivalent to 47.5% or 4 pounds active ingredients per gallon.

KEEP OUT OF REACH OF CHILDREN
DANGER  **POISON**
PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada en su idioma.

See Side Panel For Additional Precautionary Information.

24038

• PHOSDRIN is a Registered T.M. of Shell Chemical Co.

02R89 A

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
DANGER**

Poisonous if Swallowed, Inhaled or Absorbed Through Skin or Eyes. Rapidly absorbed through skin! Do not get in eyes, on skin or on clothing. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to mevinphos poisoning. Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Do not contaminate or store near feed and foodstuffs.

USE ONLY WHEN WEARING THE FOLLOWING PERSONAL PROTECTIVE EQUIPMENT DURING MIXING/LOADING, APPLICATION, REPAIR AND CLEANING OF MIXING, LOADING, AND APPLICATION EQUIPMENT, AND DISPOSAL OF THE PESTICIDE: protective suit of one or two pieces that covers all parts of the body except head, hands, and feet; chemical-resistant gloves; chemical-resistant shoes (or chemical-resistant shoe coverings or chemical-resistant boots); goggles or face shield; hood or wide brimmed hat; NIOSH or MSHA approved respiratory protection device.

IF MIXING/LOADING IS PERFORMED USING A CLOSED SYSTEM, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE: Long-sleeve shirt; long-legged pants; chemical-resistant gloves; chemical-resistant apron; shoes and socks. Goggles or face shield must be worn when the system is under pressure. All other protective clothing and equipment required for use with open systems must be available nearby.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE: Long-sleeve shirt and long-legged pants; shoes and socks. Chemical-resistant gloves must be available in the cab or cockpit and must be worn during entry to and exit from the application vehicle. All other protective clothing and equipment required for use during application must be available in the cab and must be worn when exiting the cab into treated area. When used for this purpose, contaminated clothing may not be brought back into the cab unless in an enclosure such as a plastic bag. REMEMBER — THIS CLOTHING IS INADEQUATE TO PROTECT YOU DURING REPAIR AND CLEANING OF APPLICATION EQUIPMENT AND EARLY REENTRY TO TREATED AREAS! REFER TO PROTECTIVE CLOTHING

AND EQUIPMENT REQUIREMENTS ABOVE.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water and contact a physician immediately. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or blotting.

AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving job — do not wear contaminated clothing. Personal and protective clothing worn during work must be stored and laundered separately from household articles. Clean or launder protective clothing after each use. Respirators must be cleaned and filters replaced according to instructions included with the respirator. Protective clothing and protective equipment that becomes heavily contaminated or drenched with mevinphos must be destroyed according to State and local regulations. **HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.**

The National Pesticide Telecommunication Network is available for recommendations regarding poisoning management, emergency treatment, and other information regarding the toxicity of mevinphos. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7378.

IF HANDLED INDOORS provide mechanical exhaust ventilation. Keep all unprotected persons, children, livestock, and pets away from treated area or where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away, see the First Aid (Practical Treatment) section of this label.

FIRST AID TREATMENT

If swallowed: Call a physician, or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

If in eyes: Wash at once in running water for at least 10 minutes.

Antidote

NOTE TO PHYSICIAN: Phosdrin is a cholinesterase inhibitor. Patient should be under observation for 48 hours. Atropine is antidotal. Give 4 mg intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. 2-PAM is also antidotal and may be administered in conjunction with atropine.

For Emergency Assistance Call
CHEMTREC 800-424-9300.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Drift and runoff may be

hazardous to aquatic life if applied directly to water bodies. Do not contaminate washwaters. This product is highly toxic to blooming crops and to blooming crop treatment area.

PHYSICAL

Do not use or store near

DIS

It is a violation of Federal law to use this product in a manner inconsistent with its label.

RE

Reentry into treated areas and peach orchards after application, unless label for early reentry is prohibited for 48 hours, unless the protective reentry is worn.

FOR EARLY REENTRY: SPRAYS HAVE DRIED, wear protective clothing.

FOR EARLY REENTRY: HAVE DRIED, wear protective clothing.

part of the body and gloves; chemical-resistant shoes or chemical-resistant boots. Written or oral warning must be in treated areas or fields that have clothing, period of time to take in clothing are given, warning understood by workers, reason to believe the workers. Written warning.

Area treated with this pesticide is toxic to fish and wildlife. Drift and runoff may be hazardous to aquatic life if applied directly to water bodies. Do not contaminate washwaters. This product is highly toxic to blooming crops and to blooming crop treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store in or around the home.

Do not store below 20°F, (-7°C).

Do not use or store near heat, open flame or hot surfaces.

Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**SEE ATTACHED BOOKLET FOR COMPLETE DIRECTIONS FOR USE.
DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.**

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Platte makes no warranty, express or implied, concerning this use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.


Use of Product: Platte's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, express or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

EPA REG. NO. 34704-343 EPA EST. NO. _____

NET CONTENTS _____ GALLONS

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
139 SO. MAIN STREET FREMONT, NEBRASKA 68025



POISON

6

MEVINPHOS MIXTURE, LIQUID RQ NA2783

ars, Mites and Mexican Bean Beetle—Use $\frac{1}{4}$ to 1 pint per acre.
BEEFS (Including Beef Cattle)—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre.
CABBAGE (Including Broccoli, Cauliflower, Cabbage, Cabbage Worm, Greenhouse, Leafhopper, Depressive Leafminer (adult), Felted Cabbage Bug, Grasshopper, Imported Cabbageworm, Mite, Leafhopper, Mite, Salt-Marsh Caterpillar)—Use $\frac{1}{4}$ to 1 pint per acre.
BROCCOLI (1); CABBAGE (1): For control of Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Cabbage Looper, Imported Cabbageworm, Greenhouse, Leafhopper, Depressive Leafminer (adult) and Lygus Bug—Use $\frac{1}{4}$ to 1 pint per acre. For hard to kill insects, including aphids, use 1 quart per acre, but do not treat within 3 days of harvest.
BROCCOLI SPINETS (2), CHRYSEID (2), COLLARED (2), KALE (2): For control of Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Cabbage Looper, Imported Cabbageworm, Greenhouse, Leafhopper, Salt-Marsh Caterpillar, Mite, Cabbage Caterpillar, Depressive Leafminer (adult) and Lygus Bug—Use $\frac{1}{4}$ to 1 quart per acre. For hard to kill insects, including aphids, use 1 quart per acre, but do not treat within 7 days of harvest.
CABBAGE (2); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Leafhopper, Lygus Bug, Mite, Cabbage Looper, Depressive Leafminer (adult), Cabbage Caterpillar, Salt-Marsh Caterpillar—Use $\frac{1}{4}$ to 1 pint per acre.
CELERY (2): Aphid—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Depressive Leafminer (adult), Lygus Bug, Salt-Marsh Caterpillar, Leafhopper, Cabbage Looper and Mite—Use $\frac{1}{4}$ to 1 pint per acre. For hard to kill insects, including aphids, use only be necessary to 2 pints per acre, but do not apply within 5 days of harvest.
CORN (Flint, Sweet and Popcorn) FOR GRAIN ONLY (1); Aphids—Use $\frac{1}{4}$ to 1 pint per acre.
CUCUMBERS (Cucumber) (2); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Grasshopper, Leafhopper, Mite—Use $\frac{1}{4}$ to 1 pint per acre.

On the following evening, ventilate for one hour to restore the air safe for regular work.
LETTUCE (Including Leaf, Bibb and Romaine) (10); Aphids, Corn Earworm, Cabbage Caterpillar, Depressive Leafminer (adult), Cabbage Looper, Imported Cabbageworm, Greenhouse, Mite, Lygus Bug, Salt-Marsh Caterpillar, Felted Cabbage Bug and Thrips—Use $1\frac{1}{2}$ to 2 pints per 15,000 sq. ft.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Platte makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use under handling and/or storage of the material when such use under handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other information factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: Platte's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product under varying the control of the manufacturer, no guarantee, express or implied, is made as to its effects or that the results to be obtained if not used in accordance with directions or instructions are given.

Damage: Buyer or user is advised remedy for damage by breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price and shall not include consequential or consequential damages.

EDDPLANT (2), PEPPERS (2); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Grasshopper, Leafhopper and Mite—Use $\frac{1}{4}$ to 1 pint per acre.
LETTUCE (2): Aphid—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Corn Earworm, Cabbage Caterpillar, Depressive Leafminer (adult), Cabbage Looper, Imported Cabbageworm, Greenhouse, Mite, Lygus Bug, Salt-Marsh Caterpillar, Felted Cabbage Bug and Thrips—Use $\frac{1}{4}$ to 1 pint per acre. For hard to kill insects, including aphids, use 1 quart per acre, but do not treat within 4 days of harvest.
MELONS (Including Cantaloupe, Honeydew Melons, Watermelon, Muskmelon) (10); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Cabbage Looper, Depressive Leafminer (adult), Leafhopper, Lygus Bug, Mite, Salt-Marsh Caterpillar, Felted Cabbage Bug, Cabbage Caterpillar, Greenhouse, Mite, Salt-Marsh Caterpillar, Tobacco Budworm on melons apply $\frac{1}{4}$ pint per acre.
ORCH (1): For control of Aphids use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Cabbage Caterpillar, Corn Earworm, Green Stem Bug, Mite and Velvet Bean Caterpillar—Use $\frac{1}{4}$ to 1 pint per acre.
ONIONS (Including Sweet Onions) (1); Thrips, Cabbage Caterpillar—Use $\frac{1}{4}$ to 1 pint per acre.
PUMPKIN (2): For control of Aphids apply 1 to 2 pints (0.5 to 1.0 lb. a.i./acre) as a foliage application using a maximum of 40 gallons of water per acre by ground application or 10 to 15 gallons of water per acre by air application. For light infestation use lower dosage of 1 pint (0.5 lb. a.i./acre). For hard to kill insects and increased populations use 2 pints (1.0 lb. a.i./acre). Do not treat later than 5 days before harvest for regular crops. Spraying application which results first appear and repeat as often as necessary to maintain control. Do not treat more than 3 times between harvest.
PEAS (Including Snap Peas) (7); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Grasshopper, Leafhopper, Mite and Cabbage

Caterpillar—Use $\frac{1}{4}$ to 1 pint per acre.
PODICOSES (1); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Grasshopper, Leafhopper, Mite—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre.
SQUASH (4): For control of Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Cabbage Looper, Imported Cabbageworm, Greenhouse, Leafhopper, Mite, Depressive Leafminer (adult), Cabbage Caterpillar, Salt-Marsh Caterpillar and Felted Cabbage Bug—Use $\frac{1}{4}$ to 1 pint per acre. For hard to kill insects, including Aphids—Use 1 quart per acre, but do not treat within 7 days of harvest.
WATERMELON (1); For control of Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Cabbage Looper, Depressive Leafminer (adult), Leafhopper, Lygus Bug, Mite, Felted Cabbage Bug, Salt-Marsh Caterpillar, Cabbage Caterpillar, and Greenhouse—Use $\frac{1}{4}$ to 1 pint per acre.
WATERMELON (2); Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. Grasshopper, Leafhopper and Mite—Use $\frac{1}{4}$ to 1 pint per acre.
TUBEROSES (2): For control of Aphids—Use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For control of Cabbage Looper, Imported Cabbageworm, Greenhouse, Leafhopper, Mite, Felted Cabbage Bug and Depressive Leafminer (adult)—Use $\frac{1}{4}$ to 1 pint per acre.
GREENHOUSE: Omit product should be applied in a closed greenhouse unless all doors, windows and ventilators. Lock or otherwise of entrance, post washing signs, and take whatever precautions are necessary to prevent unattended humans and domestic animals from entering the treated area. The operator must wear a full-face mask of a type found adequate for maximum protection. Apply any time the greenhouse ventilation can remain closed for two hours without endangering crops from high temperatures, such as early morning, cloudy days or late afternoon. If application is made during the day, keep the greenhouse closed tightly for at least two hours, then ventilate the enclosure thoroughly for an hour before resuming regular work. If application is made in the late afternoon, the house can be kept closed at night.

RESTRICTED USE PESTICIDE

Due to Very High Acute Toxicity to Humans and Residual Effects on Fish, Invertebrates and Aquatic Species for retail sale it can only be used by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Strict supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.



PHOSDRIN® 4.0 MISCIBLE Insecticide

ACTIVE INGREDIENTS: BY WEIGHT
 Disulfoton (2-Carbamothio-1-methylthio-1-ylmethyl) phosphorothioic acid (ester) 26.00%
 Thionyl Chloride 15.00%
 Inert Ingredients 59.00%
 TOTAL 100.00%

1000 gallons of 47.5% or 4 pounds active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
 DANGER POISON
 PELIGRO**

PRECAUTIONAL MEASURE: If water is too high, no use until products have been in contact to help into sufficient applications.

EPA REG. NO. 34784-343

PHOSDRIN is a Registered Trademark of Shell Chemical Co.

INTENTIONALLY LEFT BLANK

INTENTIONALLY LEFT BLANK

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
 100 SO. MAIN STREET FREMONT, NEBRASKA 68021

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **DANGER**

Poisonous if Swallowed, Inhaled or Absorbed Through Skin or Eyes. Possibly absorbed through skin. Do not get in eyes, on skin, or on clothing. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to respiratory poisoning. Do not breathe or get in eyes, on skin, or on clothing. Do not breathe vapors. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Keep all unprotected persons out of spraying areas or nearby where there may be danger of drift. Do not contaminate or store near food and feed-stuffs.

USE ONLY WHEN WEARING THE FOLLOWING PERSONAL PROTECTIVE EQUIPMENT DURING MIXING/LOADING, APPLICATION, REPAIR AND CLEANING OF MISTS, LOADS, AND APPLICATION EQUIPMENT, AND DISPOSAL OF THE PESTICIDE: protective suit of one or two pieces that covers all parts of the body except head, hands, and feet; chemically-resistant gloves; chemically-resistant shoes for chemical-resistant shoe coverings or chemically-resistant boots; goggles or face shield; hood or wide brimmed hat; NIOSH or MSHA approved respiratory protection device.

IF MIXING/LOADING IS PERFORMED USING A CLOSED SYSTEM, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE: Long-sleeve shirt; long-legged pants; chemically-resistant gloves; chemically-resistant shoes; and goggles. Goggles or face shield must be worn when the system is under pressure. All other protective clothing and equipment required for use with open systems must be supplied by workers. Only coverings must be worn if there is reason to believe that certain coverings cannot be understood by workers. Worker coverings must include the following information:

DANGER

Area treated with alachlor has been tested. Do not enter without appropriate protective clothing until insect data have been obtained and reentry restricted as per the label. In case of accidental exposure see FIRST AID TREATMENT found on the alachlor product label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **SPILLAGE:** Do not store in or around the home. Do not use containers below 20°F (7°C).

Do not use if more than 1 year, open flame or hot surfaces. Store in original container only. Store in a cool, dry place and out of direct heat. Carefully re-cork containers. After correct use, replace lids and store tightly. Do not put concentrate or other material into food or drink containers. Do not contaminate other pesticides, herbicides, fertilizers, soil, or food by storage or disposal. In case of spill, avoid contact, contain area and keep out animals and unauthorized persons. Control spills.

To contain spill, use surrounding area or absorb with sand, soil, or other non-hazardous material. Place discarded package in a holding container tightly covered.

PESTICIDE DISPOSAL: Pesticide wastes are highly hazardous. Complete disposal of excess pesticide, spray material, or residue is a violation of Federal law if these wastes cannot be disposed of by use according to label instructions, consult your State Pesticide or Environmental Control Agency, or the nearest State Waste Management or the nearest EPA Regional Office for guidance.

Do not use if you are pregnant or nursing. **IF A PREGNANT IS PERFORMED USING AN ENCLOSED CAB OF JOYSTICK, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE:** Long-sleeve shirt and long-legged pants; shoes and socks. Chemically-resistant coverings must be available in the cab or tractor and must be worn during application. Protective clothing and equipment required for use during application must be available in the cab and must be worn when exiting the cab and treated area. When used for this purpose, contaminated clothing may not be brought back into the cab unless in an enclosure such as a stall. **REMEMBER - THIS CLOTHING IS INADEQUATE TO PROTECT YOU DURING REPAIR AND CLEANING OF APPLICATION EQUIPMENT AND EARLY REENTRY TO TREATED AREAS REFER TO PROTECTIVE CLOTHING AND EQUIPMENT REQUIREMENTS ABOVE.**

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water and consult a physician immediately. Always wear gloves, shoes, and socks with socks and water before spraying, cutting, mowing, or tilling. **AFTER USE:** Before removing gloves, wash them with soap and water. Wash all work clothes and shoes. Shower using soap and water. Wash only clean clothes when leaving job - do not wear contaminated clothing. Personal and protective clothing used during work must be stored and handled separately from household clothes. Clean or launder protective clothing after each use. Pesticides must be stored and then replaced according to instructions provided with the pesticide. Protective clothing and protective equipment that becomes heavily contaminated or damaged with alachlor must be destroyed, or, if necessary in State and local regulations, HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED. The National Pesticide Telecommunication Network is available for recommendations regarding pesticide management, emergency

Storage and Disposal, CONTAINER DISPOSAL: Empty containers must be recycled, or otherwise disposed of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not use or reuse metal containers. Plastic: These may be recycled. These after recycling or reconditioning, or painted or placed in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Not for use or storage in or around the home. Mix with water by vigorous agitation, agitate the mixture while spraying. If allowed to stand, re-agitate before use. Do not use undiluted form. Apply when insects first appear and repeat application as often as necessary to maintain control. Apply in a complete volume of water to good uniform coverage of foliage. For application by aircraft, apply this product with 3 to 10 gallons of water per acre. **HUMAN FLAGGERS ARE PROHIBITED** during aerial application of this product unless a written release is obtained. **FRUIT AND NUT CROPS:** Use recommended dosage in application, water to obtain thorough spray coverage, but do not exceed recommended maximum gallons for this crop. If appropriate ground application is employed, do not apply less than 10 gallons of spray per acre, and avoid damage. If the product is applied to cover crops per acre, do not exceed 1/2 lb. per acre. **FOLIAR AND YIELD-TIME CROPS:** Use 10 to 125 gallons of spray per acre when using ground application equipment. Avoid drift of the product to crops for which it is not registered or to crops for which its use may be regulated but application would be closer to harvest than recommended. **NOTE:** Consult your State Department of Agriculture or local Agricultural Extension Service for information on the time interval between applications into a new harvest to the treated field.

treatment, and other information regarding the toxicity of alachlor. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7376.

IF HANDLED INDOORS provide mechanical exhaust ventilation. Keep of unprotected persons, children, livestock, and pets away from treated area or where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away, see the First Aid (First Aid) Treatment section of the label.

FIRST AID TREATMENT

If swallowed: Call a physician, or Poison Control Center. Drink one or two glasses of water and induce vomiting by spuching back of throat with finger, or if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

If in eyes: Wash at once in running water for at least 15 minutes. Continue.

NOTE TO PHYSICIAN: Pesticide is a cholinesterase inhibitor. Patient should be under observation for 48 hours. Administer atropine. Give a mg intravenously or intramuscularly. Repeat every 15 minutes until patient is fully atropinized. 2-PAM is also available and may be administered in conjunction with atropine.

For Emergency Assistance Call CHEMTRIC 800-858-0808.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Drift and runoff may be hazardous to aquatic organisms in neighboring waters. Do not apply directly to water or to watersheds, bays, streams, and wetlands. Do not contaminate water when disposing of equipment and washwaters.

This product is highly toxic to some important to direct treatment on flowering crops or weeds. Do not apply this product or allow it to drift to flowering crops or weeds while bees are active, unless the treatment is a weed.

PHYSICAL OR CHEMICAL HAZARDS

Do not use if more than 1 year or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Reentry into treated citrus groves, grape vineyards, and nut orchards and peach orchards is prohibited for 24 hours (4 days after the end of application), unless the appropriate reentry statement on the label for early reentry is used. Reentry into all other treated areas is prohibited for 48 hours (2 days) after the end of application, unless the appropriate reentry statement on the label for early reentry is used.

FOR EARLY REENTRY INTO TREATED AREAS BEFORE SPRAYS HAVE DRIED, wear of protective clothing specified on the label for an application.

FOR EARLY REENTRY INTO TREATED AREAS AFTER SPRAYS HAVE DRIED, wear protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemically-resistant gloves; chemically-resistant shoes for chemically-resistant shoe coverings or chemically-resistant boots.

Wetland or soil coverings must be given to workers who are expected to be in treated areas or in an area about to be treated with this pesticide. Wetlands include wetlands which contain waters of rivers or fields that may not be covered without specific permit or covering, ground of land that may be covered and appropriate means to take in case of accidental discharge. When soil coverings are given, coverings shall be given in a language as clearly un-

Observe days interval between last application and harvest indicated by number in () following the crop.

Fruit and Nut Crops: **APPLES (1); PEACHES (1); PEARS (1);** For control of Aphids and Mites use 1/4 to 1/2 pint per 100 gallons water. Do not exceed 1,200 gallons of finished spray per acre on apples and pears when 1/4 pint is used. For control of Greenhouseflies, Lygus Bugs and Red-Banded Leafhoppers—Use 1/4 to 1/2 pint per 100 gallons water. Do not exceed 500 gallons of finished spray per acre on apples and pears when 1/4 pint is used. Do not exceed 750 gallons finished spray per acre on peaches or pears. For control of other insects, use high relative humidity and high temperatures. May cause injury to golden retriever, dogwood and related varieties.

CHERRIES (Bear Only) (2); For control of Aphids and Mites—Use 1/4 to 1/2 pint per 100 gallons water. Do not exceed 750 gallons finished spray per acre.

GRAPEFRUIT (1); LIME (1); ORANGES (1); Aerial—Use 1 to 2 pints per acre. Apply in 200 or more gallons of water per acre. Fruiting Leafhopper, Orange Thrips, Omomyzid Leafhopper—Use 1 quart per acre. Apply in 500 gallons of water per acre. Western Tuleweed Leafhopper, Citrus Curculionid, Variegated Citrusworm, Pink Geomysid Caterpillar—Use 2 quarts per acre. Apply in 1,200 gallons of water per acre. Allow at least 7 days between applications.

GRAPES (2); For control of Aphids—Use 1/4 to 1/2 pint per acre. For control of Red-Banded Leafhopper, Mites, Red-Banded Leafhopper and Lygus Bugs—Use 1 to 2 pints per acre. At the 2 pint dosage do not treat within 5 days of harvest.

RASPBERRIES (2); Aerial—Use 1/4 to 1/2 pint per 100 gallons of water. Mites, Leafhoppers, Fruiting Leafhopper, Orange Thrips—Use 1/4 to 1/2 pint per 100 gallons of water. Apply no more than 200 gallons of water per acre.

STRAWBERRIES (1); Aerial—Use 1/4 to 1/2 pint per 100 gallons of water. Greenhouseflies, Strawberry Leafhopper, Sanjosean

Scale, Lygus Bug—Use 1/4 to 1 pint per 100 gallons of water (200 gallons per acre). For hard to kill insects, using sprays and water, use 1 quart per 100 gallons water, but do not treat within 2 days of harvest. Do not exceed 100 gallons finished spray per acre.

WILDMINTS (1); Aerial—Use 1/4 to 1/2 pint per 100 gallons of water. Mites, Greenhousefly, Leafhopper, Orange Thrips, Fruiting Leafhopper, Western Tuleweed Leafhopper—Use 1/4 pint per 100 gallons of water. By ground, apply a minimum of 200 gallons of finished spray per acre. By air, apply 1 to 1 1/2 pints per acre in a minimum of 75 gallons of water. Do not apply more than 2 1/2 quarts (1,200 gallons finished spray) per acre.

Field Crops: **ALFALFA (1); CLOVER (1);** Aerial, Alfalfa Caterpillar—Use 1/4 to 1/2 pint per acre. Greenhouseflies, Leafhoppers, Citrus Curculionid, Mites, Alfalfa Wooded Leafhopper, Lygus Bugs—Use 1/4 to 1 pint per acre.

CORN (Feed, Sweet and Popcorn) FOR FORAGE ONLY (1); Aerial, Sweet Corn—Use 1/4 to 1 pint per acre.

BUSHED GREENS (2); TURNIP TOPS (2); Aerial—Use 1/4 to 1/2 pint per acre. Caterpillar Leafhopper, Imported Citrusgreening, Potato Chewing Bug, Depressed Leafhopper, Lygus, Greenhousefly, Leafhoppers, Mites—Use 1/4 to 1 pint per acre.

PEA VINES (For Forage Only) (1); Aerial—Use 1/4 to 1/2 pint per acre. Greenhouseflies, Leafhoppers, Mites, Caterpillar Leafhopper—Use 1/4 to 1 pint per acre.

SODIUMSALT (For Forage and Stock) (2); Aerial—Use 1/4 to 1/2 pint per acre. Corn Earworm and Western Corn Rootworm—Use 1/4 to 1 pint per acre. Fall Armyworm—Use 1 pint per acre.

Vegetable Crops: **ARTICHOKES (2);** Aerial—Use 1/4 to 1/2 pint per acre. Aerial—Use 1/4 to 1/2 pint per acre. **BEANS (1);** Use 1/4 to 1/2 pint per acre. Greenhouseflies, Leafhopper