

COPY

26/099/301  
5 | 17

J. Allen Dunlop III  
Platte Chemical Company  
419 18th Street  
P.O. Box  
Greeley, CO 80632

Dear Mr. Dunlop

Subject: Label Revision  
Clean Crop Botran 6 Sulphur 25 Dust  
EPA Registration No. 39704-404  
Your Submission Dated March 1, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

SL

Susan Lewis  
Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

55182:I:A-4:Stone:R-15:KENCO:03/20/90:04/20/90:CL:sw:vo:dd:dg

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

# BOTRAN<sup>®</sup> 6 SULFUR 25 DUST

## Fungicide

ACTIVE INGREDIENTS:	By Wt.
2,6-Dichloro-4-nitroaniline .....	6.00%
Sulfur .....	25.00%
INERT INGREDIENTS: .....	69.00%
TOTAL .....	100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

EPA REG. NO. 34704-404  
EPA EST. NO. 279-CA-1

**NET WEIGHT 50 POUNDS**

\*BOTRAN is a Registered T.M. of NOR-AM Chemical Co.

02G87 AA

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if inhaled. Avoid breathing dust. Remove contaminated clothing and wash before reuse. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

## STORAGE AND DISPOSAL

**STORAGE:** Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## NOTES TO USER

Do not use for dried fruit.

Do not use on germinating seeds or annual seedlings except as suggested.

Applications of materials containing sulfur should not be made just prior to or during periods of excessively high temperatures, or within 2 weeks of an oil spray.

When growing crops for processing, consult processor before applying sulfur.

Do not allow this material to drift onto sulfur sensitive crops.

## APPLICATIONS

For Powdery Mildew control apply when disease first appears and repeat as necessary. For other disease control see the directions below. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

**Blackberry (7), Boysenberry (7), Raspberry (7), (Pacific Northwest Only):** Fruit Decay (Botrytis Rot), Powdery Mildew—Use 50 pounds per acre. Use a maximum of 4 applications. Begin application just prior to bloom and continue at 7 day intervals.

**Grapes (1):** Bunch Rot (Botrytis Rot), Storage Rot, Powdery Mildew—Use 30 pounds per acre. For Bunch Rot and Storage Rot control, begin application while clusters are still loose and repeat applications at two week intervals. Ground application preferred.

MAR 26

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered for the use on

Leaf Lettuce (Greenhouse Type): Blight (Botrytis Rot, Sclerotinia)—Use 30 pounds per acre. Apply first application at thinning. Repeat application 7 days later.

Leaf Lettuce (Greenhouse Type): Blight (Botrytis Rot)—Use 30 pounds per acre. Apply dust 10 days following transplanting. Repeat applications when plants are half mature. Do not apply to wilted plants or seedlings.

Peaches (1), Nectarines (1): Brown Rot Blossom Blight (Monilinia), Powdery Mildew—Use 60 pounds per acre. Apply at pink bud and full blossom. Fruit Decay (Monilinia), Brown Rot (Rhizopus), Powdery Mildew—Use 60 pounds per acre. Apply 18 days, 10 days and 1 day prior to harvest. Do not allow animals to graze treated areas.

Snap Beans (2): White Mold (Sclerotinia sclerotiorum), Powdery Mildew—Use 50 pounds per acre on pole varieties. On bush varieties use 37 pounds per acre. Begin applications when disease is anticipated and continue at 7 day intervals during periods favorable to development of disease. Do not feed treated forage to livestock. Sulfur may injure certain varieties. Check tolerance to sulfur prior to application.

Sweet Cherries (1): Blossom Blight—Use 50 pounds per acre. Apply at Popcorn Blossom, Full Bloom, Petal Fall. Fruit Decay (Monilinia), Brown Rot (Rhizopus)—Use 50 pounds per acre. Apply 10 days and 1 day prior to harvest. Do not allow animals to graze treated areas.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions.

**Warranty:** Platte makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** Platte's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, express or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift or runoff from target areas.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**REENTRY STATEMENT**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long-legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**CAUTION**

Area treated with BOTRAN® 6 SULFUR DUST on (insert date). Do not enter treated areas for 24 hours unless appropriate protective clothing is worn (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

Oral warnings must include the same information that is in the written warnings.

**FORMULATED FOR  
PLATTE CHEMICAL COMPANY, INC.  
150 SO. MAIN STREET    FREMONT, NEBRASKA 68025**