

ACCENT
Cyanide
1987
EST. 1974

CAPTAN 80 WDG

Fungicide

For Agricultural or Commercial Use Only.

ACTIVE INGREDIENT:	By Wt.
Captan	78.22%
Related Derivatives	1.78%
INERT INGREDIENTS:	20.00%
TOTAL	100.00%

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

* CLEAN CROP is a Registered TM of United Agri Products, Inc.

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO

PRECAUCION AL USUARIO: Si usted no lee Ingles, no use este producto. hasta que le etiqueta le haya sido explicada ampliamente.

EPA REG. NO. 34704-431 EPA EST. NO. _____

NET WEIGHT 50 POUNDS

16910

EXP. 2G

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing.

STATEMENT OF PRACTICAL TREATMENT

If In eyes: Flush with plenty of water. Call a physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg whites, protein solutions, or dextrose if not available, drink large quantities of water. Avoid alcohol. Do not give anything by mouth to an unconscious person. Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

The pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and ponds). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleanup of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply the product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

Maintainers when mixing and loading this product must wear a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron. A dust mask must be worn when mixing and loading the product.

Feldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

During application only from a branch with a completely enclosed cap or from an aerosol with an enclosed nozzle a long-sleeved shirt and long-legged pants must be worn. Chemically resistant gloves must be available in the car or cockpit and worn while mixing.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

RE-ENTRY STATEMENT

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Correspondingly post reentry information. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated so that they know when and how to enter a treated area and what personal protective equipment and clothing to wear. If field must be entered to take care of accidental exposures, written or oral warnings must be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warning cannot be understood by workers. Written or oral warnings must include the following information:

DANGER

Area treated with Captan 80 WDG on (insert date). Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn (insert State Department of Agriculture's reentry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes flush with plenty of water. Call a physician. Inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Not for use or storage in or around the house. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not pour concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact with skin and keep out animals and unprotected persons. Confine spills.

To confine spill: If liquid, dig a surrounding area or absorb with sand, cat litter or commercial clay if dry material, cover to prevent dispersal. Place damaged package in a holding container (empty containers).

Storage and Disposal Continued

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray residue, or rinsates is a violation of Federal Law. If pesticide wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning if burned, stay out of smoke.

Do not treat areas while unprotected humans or domestic animals are present in the treatment area.

Do not apply this product through any type of irrigation system.

MIXING INSTRUCTIONS: Fill the spray tank one-fourth to one-third full with clean water. Start agitation (propeller creates a rippling or roiling action on the liquid surface). Add Captan 80 at the recommended rate. Mix thoroughly. Add water to fill spray tank. Continue sufficient agitation during mixing and spraying operations to ensure a uniform spray mist.

Note: For cancer sprayer applications, the same quantity (per acre) of CAPTAN 80 WDG, as would normally be applied when spraying oil, may be applied to each acre in sufficient water for coverage.

Except as specified, begin applications before or at the first sign of disease. Repeat applications as necessary to maintain control but observe use limitations. With the exception of orchard crops, use the lower rate on small young plants with low disease potential. Use the higher rates on larger (or mature) plants or when disease potential is moderate to severe.

For orchard crops, use the higher rates as disease potential increases.

Almonds: Brown Rot, Shot Hole, Scab, Leaf Blight—1½ lbs per 100 gals spray. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if nuts are not fed to dairy animals or animals being flushed for slaughter.

Apples: Primary Scab Infestation—1½ lbs per 100 gals spray. Apply at 5 day intervals as needed, to maintain cover in prebloom, bloom, petal fall and test cover sprays. Secondary Apple Scab: Brixia Fruit Spot, Flyspeck, Bitter Root, Black Root, Bonnyphora, Black Psa, U. S. Eye Rot, Boryta Rot—¾ to 1½ lbs per 100 gals spray at 10 to 14 day intervals in second and later cover sprays up to start of harvest.

CAPTAN 80 WDG should not be used in combination with or closely following or in alternation with established sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Stayman, Baldwin, King, etc., as severe foliage injury and defoliation may occur.

Folage injury may at times occur in Red Delicious, Winesap and other sensitive varieties of apples in early season sprays.

Apricots: Brown Rot (Taeg Blight), Jacket Rot—1½ lbs per 100 gals spray. Apply at red bud stage, bloom, and repeat at 75 percent petal fall.

Cantaloupe, Cucumber: Anthracnose, Downy Mildew—1½ lbs per 100 gals spray or apply 2½ lbs per acre in water to cover. Apply at first sign of disease. Repeat at 5 to 7 day intervals or as necessary.

Cherries: Cherry Leaf Spot (West Only), Brown Rot, Brixia Rot—1½ lbs per 100 gals spray. Spray at prebloom, bloom, petal fall.

Grapes: Downy Mildew, Black Rot—1½ lbs per 100 gals spray. Apply just before bloom, repeat immediately after bloom. Depending upon susceptibility of grape variety, from 1 to 3 sprays may be needed at 7 to 10 day intervals. Dead Arm Disease (Northeast)—1½ lbs per 100 gals spray. Make test application when new growth is 4" to 6" long, repeat when growth is 8" to 10" long.

Peaches, Nectarines: Brown Rot, Scab, Coryneum (Shot Hole)—1½ lbs per 100 gals spray. Apply at pink bud, full bloom and petal fall stages. Repeat at 10 to 14 day intervals after petal fall. Also make late fall applications for Coryneum Blight.

Pears: Pear Scab (Except Pacific Northwest)—Primary Infestation—1½ lbs per 100 gals spray. Apply during early flower stage and petal fall stage. Secondary Infestation—1½ lbs per 100 gals spray during cover sprays. Severe Scab Infestation—1½ lbs per 100 gals spray. Additional treatments may be necessary. Rusting may be produced on Bosc Pears. Do not use on D'Angelo Pears.

Pruines, Plums: Brown Rot—1½ lbs per 100 gals spray. Apply at Green Tip or popcorn stage. Repeat at bloom and 75% petal fall stages. Also repeat late in season as necessary. Prune Russet Scab (Lacy Scab in Calif.)—1½ lbs per 100 gals spray. Apply at full bloom.

Strawberries: Brixia Rot—1½ to 2 lbs per 100 gals spray (200 diluted gals per acre or equivalent to 2½ to 3½ lbs per acre in water to cover). Apply when new growth starts in spring before fruit starts to form. Repeat weekly. Severe Infestation—Continue through harvest period, treating shaded areas after each picking.

Tomatoes: Early Blight, Late Blight, Gray Leaf Spot, Anthracnose, Septoria Leaf Spot—1½ lbs per 100 gals spray or apply 2½ to 3½ lbs per acre in water to cover. Severe Infestation—2½ lbs per 100 gals spray or apply 3½ to 5 lbs per acre in water to cover. Apply at first sign of disease. Repeat at 5 to 7 day intervals or as necessary to keep diseases under control.

Turf, Lawn, Brown Patch, Leaf Spot, Damp-off, Root Rot—1½ lbs per 100 gals spray. Apply 10 gals to each 1000 sq ft of turf or apply \$5.00 per acre in water for coverage. Apply prior to infection. Repeat at 7 to 10 day intervals as necessary. Do not graze treated areas or feed clippings to livestock.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered unfit for sale, use or consumption.

Compatibility: CAPTAN 80 WDG may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline materials such as hydrated lime, which reduces the fungicidal activity. Do not use this product in combination with sprays or with all-in-one formulations of parasitic products. Do not use this product on forage closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixtures.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions.

Warranty: Plaintiff makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer to his own risk.

Use of Product: Plaintiff's recommendations for the use of this product are based upon its belief to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025