



34704-443 11 26 90

PYRENONE[®]

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:	By Wt.
Technical Piperonyl Butoxide*	10.0%
Pyrethrins	1.0%
INERT INGREDIENTS**:	89.0%
TOTAL	100.0%

*Equivalent to 8.0% (butylcarbityl) (6-propylpiperonyl) ether and 2.0% Related Compounds.

**This product contains aliphatic petroleum compounds.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Below For Additional Precautionary Statements.

EPA REG. NO. 34704-443

EPA EST. NO. _____

NET CONTENTS 1 GALLON

8200 [®] PYRENONE is a Registered T.M. of the Fairfield American Corp. 12789

CAUTION

Remove pets, birds, and cover fish aquariums before spraying.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in a cool, dry place. Keep container closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATIONS

GRAIN ELEVATORS, FEED STORAGE, FARM BUILDINGS, FOOD PROCESSING PLANTS, INDUSTRIAL USES

Flour Beetles, Grain Weevils, Rice Weevils, Caddisflies, Grain Beetles, Carpet Beetles, Grain Moths (Accessible or Exposed Stages). Before attempting to spray, clean thoroughly interior and exterior of all handling equipment and completely remove all accumulations of dust, grain, and other infested food products from bins and storage areas.

SURFACE SPRAYING

Dilute 1 to 9 with water and spray all surfaces where insects crawl or rest, on walls, in cracks and crevices, niches, partitions, on shelves under equipment, and to all areas where insects harbor. Apply as a coarse wetting spray, at the rate of 1 gallon diluted emulsion per 750 sq. ft. of wall, ceiling, floor, partition or equipment surfaces. Repeat cleaning operations and spray treatments as needed. Do not apply on surfaces which could be stained by water.

Cockroaches, Waterbugs, Ants, Silverfish, Cheese Mites, and Certain Spiders: Dilute 1 to 9 with water and apply diluted emulsion at the rate of 1 gallon per 750 square feet of surface to be sprayed. Spray thoroughly into all hiding places, entrances and trails of these insects, into cracks in floors, partitions and walls, and under and behind equipment, shelves, cabinets and counters. Repeat treatments at regular intervals or as often as necessary to keep insects under control.

SPACE TREATMENT

To kill the accessible exposed adults of granary weevil, rice weevil, confused flour beetle, cadelle, saw toothed grain beetle, angoumois grain moth and grain mites, fog with Aerosol Generator using 1 pint of Undiluted Concentration to 40,000 cubic feet. Do not remain in treated areas. Ventilate the area after treatment is completed.

CONTINUED

2/3

PYRENONE® EMULSIFIABLE CONCENTRATE, cont'd.

LIVESTOCK USES

HORSES, CATTLE, GOATS, HOGS, AND SHEEP—Lice: Dilute 1 to 1000 with water. Apply as a thorough coating spray or by dipping to wet all parts of body. Usually 1½ to 3 gallons is required for horses or cattle and 2 to 4 qts. for sheep, goats, and hogs. Repeat as needed. Sheep Ticks: Dilute 1 to 49 with water. Apply as a spray or by dipping to wet all portions of body. Repeat as needed.

POULTRY HOUSES—Chicken Mites and Bedbugs: For chicken mites dilute 1 to 19 with water. Apply liberally and thoroughly as a driving spray with particular attention to perch poles, nests and cracks in floors and walls. Spray litter thoroughly. Repeat treatment when needed. For Bedbug control dilute 1 to 9 parts of water.

CATTLE AND HORSE—Horse Flies, Stable Flies, and Deer Flies: Dilute 1 to 9 with water. Apply as a spray at the rate of about 1 qt. per animal to wet over entire body. Spray each animal completely with particular attention to the legs and underline. Spray applications on animals to be protected should be made routinely every 3 days.

CATTLE AND HORSES—Horn Flies and Mosquitoes: Dilute 1 to 19 with water. Apply as a spray at the rate of about 1 qt. per animal to wet hair over entire body. Spray each animal completely with particular attention to the legs and underline. Spray applications on cattle to be protected should be made on a once a week schedule.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

**FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025**

Reviewed. Registrations to be in accordance with
and Golden State 2
to West Coast

