

# MILL SPRAY

A Ready To Use Fog or Spray For  
Mills, Elevators, Terminals, Farm Storage,  
Flat Storage, Warehouses, Feed & Seed Stores  
and Food Plants

## ACTIVE INGREDIENTS:

Piperonyl butoxide technical\* ..... 1.50%

Pyrethrins ..... 0.15%

INERT INGREDIENTS:\*\* ..... 98.35%

TOTAL ..... 100.00%

\*Equivalent to 1.20% (butylcarbityl) (6-propylpiperonyl) ether and 0.30% of Related Compounds.

\*\*Contains petroleum distillate.

KEEP OUT OF REACH OF CHILDREN  
CAUTION

See side panel for additional precautionary statements.

EPA REG. NO. 34704-444

EPA EST. NO. \_\_\_\_\_

NET CONTENTS \_\_\_\_\_ GALLONS

FORMULATED FOR  
PLATTE CHEMICAL CO.

150 SO. MAIN STREET      FREMONT, NEBRASKA 68025

EXP02Y91

It in eyes: Flush with plenty of water and get prompt medical attention.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

## PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE MIXTURE! Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Personnel should use clothing and equipment consistent with good pesticide handling.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

This product is a ready to use spray for the control of certain insect pests of stored grain, stored food and food products. It kills on contact the exposed (or accessible) stages of the Granary Weevil, Rice Weevil, Confused Flour Beetle, Cadelle, Saw Toothed Grain Beetle, Indian Meal Moth, Angoumois Grain Moth, Mediterranean Flour Moth, Red Flour Beetle, Yellow Mealworms, Dark Mealworms, Cigarette Beetle, Drugstore Beetle, Clover Mites, Cheese Mites, Houseflies, Mosquitoes, Gnats, Fruit Flies, Spiders, Wasps, Hornets, Silverfish, Sandflies, Small Flying Moths and Grain Mites.

For best results processing equipment and storage areas should be thoroughly cleaned before spraying. Waste grain, dust, dirt and debris should be removed and destroyed so the spray will come into direct contact with as

spraying directly on and around the base where insects hide and walk over them.

For extended protection and control in warehouses and food product storages, spray heavily cracks, crevices, recesses and hiding and breeding areas as well as walls, floors, shelves and between stacked cartons or containers.

**SPACE SPRAY:** For control of flying, running and crawling stages of insects listed on this label, apply this product in a good machine adjusted to deliver a fine mist. If possible, keep doors and windows closed for 30 minutes after application. Against flying insects, direct the spray toward the upper portions of the enclosures, filling the room with mist and using 2 to 4 ounces of material to each 1000 cubic feet of space. In the control of active stages of running and crawling pests, direct the mist into all places commonly inhabited by these insects and spray liberally all protected hiding and breeding places.

**CAUTION:** Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

Do not remain in treated areas. Ventilate the area after treatment is completed.

**VAPOR SPRAY:** This product is especially applicable for use as a fog or vapor treatment, in special electric, steam or compressed air equipment. Use in accordance with the manufacturers' directions for the type of applicator selected. Most commonly used applicators are designed to give best results when the material is used at 1 pint to each 8000-15,000 cubic feet of space. In all cases, dosage should be sufficient to fill enclosure with fog or vapor.

**GRAIN SURFACE TREATMENT:** As a Grain Surface Spray, level the grain surface and spray at the rate of 1 gallon to 750 square feet. Application can be made before and after fumigation in the event fumigation is required. Repeat every 4 to 6 weeks or as infestation reappears. Apply with a good hand or electric sprayer for uniform distribution.

To get adequate insect kills with Surface Sprays, results depend entirely on the amount of insecticide deposited on areas with which insects will come into contact.

## NOTICE

Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

BEST AVAILABLE COPY