



VAPONA[®] 1% LIVESTOCK SPRAY

READY TO USE — FOR USE ON BEEF AND DAIRY CATTLE — CONTROLS HOUSEFLIES, FACE FLIES, HORN FLIES, STABLE FLIES AND MOSQUITOES

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl Dimethyl Phosphate* ..	0.92%
Related Compounds*	0.08%
INERT INGREDIENTS**:	99.00%
TOTAL	100.00%

*Equivalent to 1% Dichlorvos and Related Compounds.
**This product contains Aliphatic petroleum hydrocarbons.

KEEP OUT OF THE REACH OF CHILDREN WARNING

See Side Panel For
Additional Precautionary Statements

EPA REG. NO. 34704-445

EPA EST. NO. _____

NET CONTENTS 1 GALLON

VAPONA is a Registered T.M. of Shell Chemical Co.

8195

12799

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

MAY BE FATAL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. Do not use in areas where food or feed crops are growing. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

PRODUCT INFORMATION

This product is specially prepared for uses recommended in the control of certain insects attacking livestock (Houseflies, Horn Flies, Stable Flies and Mosquitoes).

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BEEF and DAIRY CATTLE: Flies (Houseflies, Horn Flies, Stable Flies, Face Flies and Mosquitoes): Apply 1 to 2 fluid ounces per animal as a mist spray daily with a hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply in excess of 2 fluid ounces per animal per day.

ANIMAL BUILDINGS (Dairy Barns, Milk Sheds, Shelter Sheds and Horse Barns): For control of Flies, Gnats and Mosquitoes. **SURFACE Spray:** Dilute this product with equal parts of diesel fuel. Apply dilute mixture as a coarse wet spray at the rate of 1 quart per 1000 square feet for milk sheds, stables, livestock barns, loafing sheds, pig pens; treat around (doorways, feed storage rooms, alleyways and window sills.

In POULTRY HOUSES: Apply to manure, window sills, outside of penned enclosures, and on floors of feed storage rooms. Also controls roaches, fleas, ants, sowbugs and brown dog ticks. Repeat as necessary. Do not treat animals or humans.

FLY MAGGOTS: In maggot breeding areas in garbage dumps and manure piles, dilute 1 to 2 pints of this product in equal parts of diesel fuel and apply 1 to 2 quarts of the 0.5% dilution per 100 square feet.

FOG or MIST: Use 1 pint of this product undiluted per 8000 cubic feet of space. Animals can be left in building provided they have not received a direct application within eight hours. Before applying spray reduce air movement as much as possible by closing doors, windows and other openings.

IMPORTANT: Do not contaminate water, feed or foodstuffs, milk or milking utensils.

CONTINUED

2/2

VAPONA® 1% LIVESTOCK SPRAY, cont'd.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but neither this warrant nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

**Not Reviewed. Registrant claims to be in accordance with
Registration of 1% Vapona Livestock Spray, Phase 2
Response, Form 212, under the Inert Chemicals Status.**

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025

PLATTE CHEMICAL CO.
FREMONT, NEBRASKA