NOTICE OF PESTICIDE

(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

WAME OF PESTICIDE PRODUCT

Clean Crop Chlorpyrifes 1.14G

Insecticide + Fertilizer

MAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Platto Chemical Company P.O. Box 667 Greeley, CO 80632

ı

NOTE: Changes in labeling formula differing in aubstance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the digistrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pest-leide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
- 2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 34704-449."
 - b. In your Environmental Hazards paragraph, change "Keep out of any body of water" to "Do not apply directly to water." Also, add "Do not contaminate water when disposing of equipment washwaters."
 - c. Revise the If In Eyes section to read:

Plush with plenty of water. Call a physician if irritation persists.

- 3. Please submit storage stability data for this product withis 15 months from the date of this Registration Notice.
- 4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Dennis H. Edwards, Jr. Product Manager (12) Insecticide-Rodenticide Branch Registration Division (TS-767C)

Enclosures

CLEAN CROP

CHLORPYRIFOS 1.14G INSECTICIDE + FERTILIZER

Lawn Insect Control Kills:

Ants, Armyworms, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Hyperodes Weevils, Sowbugs (Pillbugs), Sod Webworms and White Grubs (Larvae of Japanese Beetle and European Chafer) in Home Lawns and Other Recreational Areas & Ornamental Turf.

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-

1.14% pyridyl)phosphorothioate3..... INERT INGREDIENTS:

98.86%

EPA Reg. No. 34704-

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN (12 pt.) CAUTION (18 pt.)

00T161088

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS - CONSERVED UNDER FOR No. CAUTION

Under the Polici Insecticità. Fungicide, and Posicitative Act an amended, for the posticide 34704-449

May be harmful if swallowed. Do not take internally. Do not dust. Avoid contact with eyes, skin or clothing. Wash contaminated clothing before reuse. Wash thoroughly after handling and before eating or smoking. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry. Do not contaminate feed, foodstuffs, and water supply.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with plenty of moap and water. Wash clothing before reuse.

IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician immediately.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention immediately.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. *Keep out of any body of water. Do not apply where runoff is likely to .. occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal.of waster Apply this product only as specified on this label.

> FORMULATED FOR PLATTE CHEMICAL COMPANY, INC. 150 SO. MAIN STREET FREMONT, NEBRASKA 68025

> > BEST AVAILABLE COPY

CLEAN CROP CHLORPYRIFOS 1.14G + FERTILIZEF

45

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

STORAGE: Store product in its or ginal container in a cool, dry,

locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product

may be disposed of on site or at an approved waste disposal

lfacility.

CONTAINER DISPOSAL: Completely empty bag into application equipment.

Then dispose of empty bag in a sanitary landfill or by incineration,

lor, if allowed by State and local authorities, by burning. If burned

lstay out of smoke.

SOD WEBWORMS, CHINCH BUGS, ANTS, FLEAS, CRICKETS, GRASSHOPPERS CUTWORMS, BROWN DOG TICKS, SOWBUGS, EARWIGS

(HOUSEHOLD: Do not reuse empty container. Wrap and put in trash.____!

SURFACE FEEDING INSECTS: For control of Chinch Bugs, Sod Webworms, Fleas, Ants, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Grasshoppers, and Sowbugs, apply 40 lbs. to uniformly cover 20,000 sq. ft. of turf (20 lbs. treats 10,000 sq. ft.). Refer to Spreader Settings. Apply in the spring when insects first appear troublesome. Allow 3-4 days for maximum control and retreat as needed, but not more often than one treatment in seven (7) days. Additional treatments at 30 to 60 day intervals may be needed for continued control of Sod Webworms. After application water the area in order to move the granules into the zone of insect infestation. Do not allow children or pets on the area until the grass is dry.

CONTROL WHITE GRUBS AND OTHER ROOT FEEDING INSECTS

For control of Root Feeding Insects such as Billbug Larvae and the "White Grub" larvae of Japanese Beetle, European Chafer, Southern Chafer and Hyperodes Weevil apply 40 lbs. to treat 10,000 sq.\ ft. of lawn (20 lbs. treats 5,000 sq. ft.). Apply the granules uniformly over the area and immediately follow the application with a thorough watering to wash the insecticide into the soil where the insects are feeding. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Most areas also have a feeding period in the spring, or grubs are harder to control them. Consult your local Cooperative Agricultural Extension Service for proper time to treat in your area. Contest will be improved by removing heavy thatch buildup (over 1/2 to 3/4 inch) before application. For the Hyperodes (Turfgrass) Weevil apply in mid-April and again in mid-May at the same rate of 40 lbs. per 10,000 sq. ft. Consult your Agricultural Extension Service for the best-time to treat in your area. Keep children and pets off treated areas until insecticide has been washed into the soil and grass is dry.



POOR INSECT CONTROL WILL RESULT IF YOUR SPREADER APPLIES LESS GRANULES THAN RECOMMENDED....MAKE SURE TO APPLY THE TOTAL POUNDS PER SQUARE FEET AS SHOWN IN THE FOLLOWING CHART:

SPREADER SETTINGS

		Root Feeding	Surface Feeding
		Insects: Per	Insects: Per
		10,000 sq.ft.	20,000 sq.ft.
		4.0 lbs./	2.0 lbs./
Spreader	Apply	1,000 sg. ft.	1,000 sq. ft.
Scott	40 pounds	#5 1/2	#4
Cvclone	40 pounds	#4 1/2	#3 3/4

When using a 20 lb. bag: Spreader settings will be the same as the 40 lb. bag but it will treat only half the amount of sq. ft. as the 40 lb. bag. These settings are based on a "One Time Over" application.

NOTICE

Seller warrants that the product conforms to the chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



